




EVALUATION AND COMPARISON OF REGIONAL CLIMATE MODELS OVER THE IBERIAN PENINSULA

C.Sánchez de Cos, C.Jiménez, J.M. Sánchez-Laulhé, J.M.Sánchez
AGENCIA ESTATAL DE METEOROLOGÍA, SPAIN (csanchez@inm.es)



ABSTRACT

Different relations between parameters involved in both water and energy land surface budgets are computed from daily ERA-Interim data for the months of July and November (representative of the dry and wet season) in the period 1989-2008 over an area within the Iberian Peninsula covering most of Tajo and Guadiana basins (from 40.5N to 37.5N, and from 7.0W to 2.0W). The main objective of this work is to use the obtained relations for the evaluation and comparison of regional climate models (RCMs) participating in the ENSEMBLES project. This approach was first proposed by Betts (2004) for comparing and evaluating global climate models. He proposed the assessment of model surface components as a system with widely connected components. In this way, models are compared among themselves and evaluated against observational data. The work is mainly focused on the goodness of the representation of physical surface processes and their feedbacks. The obtained relationships among different parameters are therefore considered as imposed restrictions by physical processes which can be used to evaluate RCMs.

OBJECTIVES

- 1) To analyze in an area of the Iberian Peninsula the behavior of the ERA-Interim model in terms of daily water and surface energy balances to establish some relationships that should meet the climate models in this area.
- 2) In a next step we will check if these relations obtained are well represented in the models of the ENSEMBLES project.
- 3) If these relationships are satisfied in a similar way on the models of the ENSEMBLES project we will obtain a greater robustness for future projections for the SRES A1B scenario for the period 2001-2050.

INTRODUCTION

- The climate processes can be represented in mathematical terms based on physical laws.
- The climate system of the land surface exerts controls on the amplitudes of the variables in all time scales.
- The changes in the radiative flux at the Earth's surface affect the surface heat and moisture.
- The comparison of predictability and uncertainty of the models indicates that the latter are relatively more important in most of the European region.
- It is necessary to know the goodness of the models to reproduce the reference climate in different regions and thus to begin the task of construction of probabilistic scenarios.



Fig.1-The yellow quadrilateral is the model approximation to the Tajo and Guadiana river basin shown in the left (beige-shaded)




Fig.3-Schematics of surface energy and water fluxes. FCS, FCL, LW and SW stand for surface sensible and latent heat flux, surface longwave and shortwave radiation flux, respectively. E, P, and R stand for evaporation, precipitation and runoff. The surface water balance and the surface energy balance equations.

METHODOLOGY

The methodology used by Alan K. Betts (2004) is followed for the understanding of the linkages and feedbacks between physical processes in the models, so that different models can be compared with each other and with data. ERA-Interim model data (grid 0.5X0.5) are used for the period 1989-2008, in an inland area of the Iberian Peninsula which covers roughly the Tajo and Guadiana basins for July and November. Daily data are chosen since the surface climate variations can be represented with considerable accuracy by means of the average daily state variables and average daily flows. The work is made with the months of July and November as representative of the dry season and wet season respectively in the area of calculation, but sometimes we also use data from May because there is a huge drought in July in this area and this does not allow us to draw conclusions of the connections between some parameters. The mean lifting condensation level height is a good approximation to mean cloud base in the daytime (Hnub): $Hnub = 122(T-Td)$. The index moisture soil SMI is computed for the first 0-7cm soil layer as $SMI = (SM-0.171)/(0.323-0.171)$; SMI is the model soil water fraction, 0.171 the model soil permanent wilting point and 0.323 is the model field capacity. The evaporative fraction (FE) = $FCL/(FCL+FCS)$.

RESULTS




Fig.4-Scatterplots for the selected area of the Iberian Peninsula of Hnub as a function of SMI

Fig.5-Surface LWnet as a function of cloud base, total cloud cover, the diurnal range of radiometric skin temperature and soil moisture index (0-7cm layer) for July and Nov for the selected area of the Iberian Peninsula

Table 1-What happens when LWnet decreases?

Fig.6-(top) Surface energy balance terms SWnet, LWnet, sensible heat FCS, and latent heat FCL as a function of mean cloud-base Hnub for Jul and Nov, and (bottom) radiation fluxes as a function of total cloud cover

Fig.7-As in Fig.5, but SWnet against LWnet, total cloud cover, sensible heat flux and cloud base

Table 2-What happens when SWnet decreases?

Fig.10-As Fig.5, but for sensible heat flux against maximum skin temperature and SWnet, and mean cloud-base height against SWnet and sensible heat flux for May, July and November.

Fig.11-Scatterplots for the selected area of Hnub and evaporative fraction partitioned into two ranges of Fnet as a function of SMI

CONCLUSIONS




Fig.11-Surface LWnet as a function SMI, TCC, Tsmmax-Tsmmin for Jul and Nov for the selected area of the Iberian Peninsula for ERA-Interim data(1989-2008) (left-hand) and for the Madeira River basin (right-hand, obtained by Betts 2004) for ERA-40 data(1972-2002)

Fig.12-The coupling of the soil moisture, cloud base, cloud cover, radiation fields, sensible and latent heat fluxes. We can see that there are strong feedbacks on the daily time scale between the cloud field and the surface energy budget. This means that cloud and the land surface components of a model must be evaluated as a tightly coupled system, not as independent components.

Fig.13-If we assume that net longwave radiation flux LWnet increases the change they will experience the other terms is indicated in the table (increases in red, decreases in blue)

- > We have found out several relationships that meet the ERA-Interim model within the Iberian Peninsula
- > The conclusions for our work area with ERA-Interim data largely coincide with those obtained by Betts for the Madeira River basin with data from ERA40.
- > We have verify that the variables and fluxes are indeed interconnected.

REFERENCES

• Barker T., Bonville, I., Bernsten, J., E. Bognar, P. R. Bosch, R. Dave, O. R. Davidson, B. S. Fisher, S. Gupta, K. Hashino, G. J. Haug, S. Hain, R. Ribes, S. Kobayashi, M. D. Levine, D. L. Martin, O. Moser, B. Metz, L. Meyer, G.-J. Nabuurs, A. Najjar, N. Nakicenovic, R. H. Rogner, J. Roy, J. Sahay, R. Schock, P. Shukla, R. E. H. Sims, P. Smith, D. A. Tjipk, D. Urge-Vorsatz, D. Zhou, 2007. TECHNICAL SUMMARY IN CLIMATE CHANGE 2007: MITIGATION. CONTRIBUTION OF WORKING GROUP III TO THE FOURTH ASSESSMENT REPORT OF THE INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE (B. Metz, O. R. Davidson, P. R. Bosch, R. Dave, L. A. Meyer (eds)), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

• Betts Alan K., UNDERSTANDING HYDROMETEOROLOGY USING GLOBAL MODELS, 2004, DOI: 10.1175/BAMS-85-11-1673.

• Betts Alan K., LAND-SURFACE-ATMOSPHERE COUPLING IN OBSERVATIONS AND MODELS, 2009, DOI: 10.3894/JAMES.2009.14

• Brunel, M., Orosco, M. J., Castro, M., Galán, P., López, J. A., Martín, J. J., Pastor, A., Petros, G., Ramos, P., Rodríguez, J., Rodríguez, E., Sora, I., Torres, L., 2009. GENERACION DE ESCENARIOS REGIONALIZADOS DE CAMBIO CLIMÁTICO PARA ESPAÑA. Agencia Estatal de Meteorología (AEMET), Ministerio de Medio Ambiente, y Medio Rural y Marino. ISBN: 978-84-8320-470-2.

• Christophorou, Daniel Lutz, and Urs Beyerle, THE SOIL-PRECIPITATION FEEDBACK: A PROCESS STUDY WITH A REGIONAL CLIMATE MODEL, 1998, Volume 12, 722-741.

• Dimoyez Paul A. and Fawong J. Zeng, THE SENSITIVITY OF SURFACE FLUXES TO SOIL WATER CONTENT IN THREE LAND SURFACE SCHEMES, 2000, Journal of Hydrometeorology, volume 1, 121-134.

• FS DOCUMENTATION: CYSRI. OPERATIONAL IMPLEMENTATION, 3 June 2008.

• Koster RD, Domeyda PA, Guo Z, Bonan G, Chen E, Cox O, Gordon CT, Keefer S, Kwiatkowsky E, Lawrence D, Liu P, Lu CH, Malyshev S, McAvaney B, Mitchell K, Mocko D, Ohi T, Oleson K, Pitman A, Sud YC, Taylor CM, Vasegh D, Vasic R, Xue Y, Yamada T, GLACE Team, REGIONS OF STRONG COUPLING BETWEEN SOIL MOISTURE AND PRECIPITATION, Science 20 August 2004, Vol.305 no.5887, pp. 1138-1140, DOI: 10.1126/science.1100217.

• Lorenz, E.N., 1963. DETERMINISTIC NONPERIODIC FLOW, Journal of the Atmospheric Sciences, 20, 130-141.

• Neftci, F., Fast, I., Hasse, H., Casasnovas, L., THE STREAM ENSEMBLES PROJECTIONS OF FUTURE CLIMATE CHANGE, ENSEMBLES Theoretical Report No.3, April, 2008.

• Palmer, T.N., 2001, A NONLINEAR DYNAMICAL PERSPECTIVE ON MODEL ERROR: A PROPOSAL FOR NON-LOCAL STOCHASTIC-DYNAMIC PARAMETRIZATION IN WEATHER AND CLIMATE PREDICTION MODELS, Quarterly Journal of the Royal Meteorological Society, 127, 279-304.

• Trenberth, K.E., & Dai, S., 2005, RELATIONSHIPS BETWEEN PRECIPITATION AND SURFACE TEMPERATURE, Geophys. Res. Lett., 32, L14703, DOI: 10.1029/2005GL022750.

• Velloso, P., THE ROLE OF THE LAND SURFACE IN THE CLIMATE SYSTEM, ECMWF/2002, Meteorological Training Course Lectures Series.

• Velloso, P., L. D. Luna, C. Fra, S. I. Semivertova, and C. Sospit, PREDICTABILITY AND UNCERTAINTY IN A REGIONAL CLIMATE MODEL, J. Geophys. Res., 109(D18), 4586, DOI: 10.1029/2002JD002810, 2003.

• Wilks, D.S., 2006, STATISTICAL METHODS IN THE ATMOSPHERIC SCIENCES, ISBN 13: 978-0-12-751966-1.