

Climatic observations and instrumental reconstructions: development of high-quality climatic gridded products

Sixto Herrera², Juan Javier Miró Pérez⁴, Pere Quintana-Seguí⁵, Julián Gonzalo⁶, José Antonio Ruiz-Arias⁷, Jose Carlos González-Hidalgo⁸, José Antonio Guijarro³, Jose Antonio López¹

1 AEMET, Agencia Estatal de Meteorología, Spain

2 Meteorology Group. Dept. Applied Mathematics and Comp. Sci. Universidad de Cantabria. Spain

3 State Meteorological Agency (AEMET), meteorology group, Spain

4 Departament de Física de la Terra i Termodinàmica, Facultat de Física, Universitat de Valencia, Spain

5 Observatori de l'Ebre (OE), Universitat Ramon Llull – CSIC, Roquetes, Spain

6 Grupo SECF Fitoclimatología y Cambio Climático, Instituto Universitario de Investigación en Gestión Forestal Sostenible - Universidad de Valladolid, Palencia, Spain

7 Solar Radiation and Atmosphere Modeling Group, Physics Department, and Centre for Advanced Studies in Energy and Environment, University of Jaén, Spain

8 Department of Geography, Zaragoza University, Spain

Introduction

Observations, studies of feedback processes, and model simulations are the basis of our knowledge and understanding of the climate system, as has been shown by the Working Group I of IPCC's Fifth Assessment Report (IPCC, 2013). In order to improve and adapt observational datasets to the actual needs of the scientific and impact studies communities, as well as to the needs of policy makers, many international programs have been setup to collect data and information used to study climate (e.g. Copernicus, formerly GMES: Global Monitoring for Environment and Security), and to provide climate services (e.g. Copernicus and GFCS: Global Framework for Climate Services). The main focus of some of these projects is to prepare and deliver climate information that meet users' needs (WMO, 2011). Within these initiatives, a set of Essential Climate Variables (ECVs) (GCOS, 2010a; Bojinski et al., 2014) have been identified, based on their relevance to characterize the climate system and its changes, feasibility and cost effectiveness. Also, guidance and best practices have been defined to obtain and support the generation of long-term, high-quality and traceable ECV datasets (GCOS, 2010b).

In this framework, several projects have grown in the last decade to improve the temporal and spatial coverage of our observational networks (e.g. EURO4M or UERRA Projects), process the raw measurements

to isolate the climatic signal (e.g. Action Cost HOME or MeteoMet), develop products (e.g. gridded datasets) useful for the different communities, and include them in the assimilation process of the reanalysis products (e.g. UERRA Project). Despite the improvement of our observations and the development of very high quality observational datasets (see Guijarro et al., this volume, for more details), high-resolution gridded observational products that capture the temporal and spatial diversity of climatic variables are increasingly demanded by the climate analysis and impact communities. Several products have been developed within the activities of different national (e.g. Portugal: Belo-Pereira et al., 2011; Romania: Birsan and Dumitrescu, 2014; Dumitrescu and Birsan, 2015; the Alpine region: Isotta et al., 2014; or Germany: HYRAS precipitation database, Rauthe, 2013) or international (e.g. E-OBS, Haylock et al., 2008; van den Besselaar et al., 2011) projects by applying some interpolation process to the raw observations.

This work describes the main advances and initiatives since the previous CLIVAR-Spain assessment (Pérez and Boscolo, 2010) on the development of climatic gridded products emphasising the studies affecting the Iberian Peninsula.

Development of gridded datasets

As described in the previous section, the high-

quality observational datasets usually comprise of a limited number of time series non-homogeneously distributed, locally representative and, in most cases, spanning different time periods. However, datasets with different features are needed for different purposes, and some post-processing is needed to build adequate products. High-resolution gridded datasets built from quality controlled observational datasets are increasingly demanded for climate analysis and the impacts communities, and several products have been developed in the last few years in response to this demand (e.g. E-OBS in Europe or MOPREDAS, MOTEDAS, SAFRAN or Spain02, among others, in Spain).

Due to the high climatic variability and complex relief of Spain, the available international products (e.g. E-OBS, WATCH or WFDEI) are not able to reproduce properly the different Spanish climate regimes (Herrera et al., 2012; Herrera et al., 2015; Bedia et al., 2013). Many regional and national gridded datasets have been built in the last decade using different approaches according to the needs of the scientific and impact communities, leading to a wide range of products which will be summarized in this section.

Focusing on the spatial resolution, Ninyerola et al. (2007) developed a dataset with a 200-m spatial resolution for the Iberian Peninsula of monthly and annual climatologies of precipitation, radiation and temperatures, which has been extensively used in ecological modelling. Gonzalo et al. (2010) built a similar dataset to develop the phytoclimatic diagnosis of the Peninsular Spain, but using different interpolation method and explicative variables, and with lower spatial resolution (1 km).

At a regional scale, Vicente-Serrano et al. (2003, 2007 and 2010) built climatic maps of monthly precipitation, temperatures and fog in the Ebro Valley and Aragón at a 1 km spatial resolution, considering several interpolation methods. Garzón-Machado et al. (2014) built a climatophilous potential natural vegetation map with a spatial resolution of 25 m for La Palma Island, in the Canary Islands. Ruiz-Arias et al. (2011, 2015 and 2016) developed solar radiation gridded datasets for Andalusia (1km) and Peninsular Spain and Balearic Islands (10km), using this one to assess the solar radiation of the WRF Model. However, the aim of most of these studies was comparison between different interpolation methods and, as a result, most of these datasets are distributed only under request.

Despite the very high spatial resolution of the previous datasets, time series of monthly, daily or intra-daily data are necessary in many cases (e.g. for trend analysis). In this sense, González-Hidalgo et al. (2011 and 2015) developed monthly datasets of precipitation (MOPREDAS) and temperatures (MOTEDAS) for mainland Spain to be used for trend analysis, model validation and downscaling

purposes, covering 1945-2005 and 1951-2010, respectively. Both datasets have a spatial resolution of 0.1°, and have made use of a dense observational network of 2670 stations for precipitation and 1358 for temperature from the Spanish Meteorological Agency (AEMET).

Herrera et al. (2011, 2012 and 2016) developed a series of high-resolution daily precipitation and temperature gridded datasets (Spain02) for the peninsular Spain and the Balearic Islands. To this aim, 2756 and 250 quality-controlled stations from the Spanish Meteorological Agency (AEMET) were used to build the different existing versions. In particular, version v2 (0.2° regular grid) is a gridded dataset with local representativeness, appropriate to local and extreme events analysis (Herrera et al., 2012), whilst version v4 includes three different resolutions matching Euro-CORDEX grids (0.11°, 0.22° and 0.44° in rotated coordinates). This version provides areal representative (AA) values (by averaging from an auxiliary 0.01° grid) in products AA-2D, AA-3D and AA-OK, following the notation used in Herrera et al. 2016, and it is an appropriate product to validate regional climate models (RCM). Point representative values are still provided in version v4 in product OK (Ordinary Kriging). Version v3 was built considering quality controlled stations with long time series of precipitation (at least 40 years with at most 10% of missing data per year) in order to obtain an appropriate dataset for trend analysis. The different versions of Spain02 are freely distributed for research purposes at the AEMET climate services portal. Version v2 is the reference dataset used in the development of the scenarios for the national program for regional climate change (PNACC-2012). Version v4 is one of the national gridded datasets considered in the framework of the COST Action VALUE (<http://www.value-cost.eu/>) used to validate regional climate models of Euro-CORDEX, the European branch of the CORDEX initiative.

In addition to the Spanish gridded datasets, Belo-Pereira et al. (2011) developed a dataset for Portugal (PT02) using 400 quality-controlled stations, and with the same interpolation technique and grid used in version v2 of Spain02. This has led the development of two equivalent datasets for a common period (1951-2003) that have been combined in recent studies to obtain the dataset IB02, a precipitation gridded dataset of daily precipitation for the Iberian Peninsula (Ramos et al., 2016; Sousa et al., 2016).

Recently, an extension of the SAFRAN analysis (Durand et al., 1993; 1999) has been applied to the mainland Spain and the Balearic Islands (Quintana-Seguí et al., 2016; 2017), leading to an hourly high-resolution (5 km) gridded dataset based on daily precipitation, and six-hourly temperature, wind speed, relative humidity and cloudiness. SAFRAN also provides modelled incoming visible and infra-red radiation. This way, it provides all the necessary variables to force, for instance, a Land-Surface Model (LSM) or any other physically based distributed

hydrological model. The product covers the period 1979-2014, and is available for research purposes from the Mistrals-HyMex database (Quintana-Seguí, 2015).

On a regional scale, several high spatial resolution (~1 km) daily datasets have been developed in the last years within regional projects or specific analysis/needs of some research groups. Militino et al. (2015) defined and validated a spatio-temporal interpolation method to obtain a precipitation grid with a resolution of 1 km covering the Navarra region. Within the regional project "Escenarios Regionales Probabilísticos de Cambio Climático en Cantabria: Termoplumiometría", Gutierrez et al. (2010) built a similar product for Cantabria, including precipitation and temperature, which was used to project the future climate change scenarios for the region by means of statistical downscaling. Miró Pérez et al. (2015) combined statistical downscaling and spatial interpolation to obtain a very-high resolution grid of daily maximum and minimum temperatures for the Valencia region considering ~300 stations from different institutions (AEMET, CEAM, SIAR and IIG) covering the period 1948-2011. This dataset is referred as SDSITVC in Miró et al. (2016), where it is proposed as a tool for estimating bioclimatic change already occurred in mountainous areas.

Summary and conclusions

In the last few years, a great effort has been carried out to improve the quality of the tools and products used to analyse the climatic system from different points of view. In this sense, several initiatives and projects have emerged aiming to rescue and digitalize existing observational data, with special attention to regions with a poor spatial and/or temporal coverage, and to develop adequate tools and methods to elaborate high-quality datasets for climate analysis.

The Spanish climatological community is involved in most of the current projects and initiatives related to the development of several tools and products (e.g., Climatol package for data analysis and homogenization, or the gridded datasets MOPREDAS, SAFRAN or Spain02), which are being presently used by the international community in many studies, and extending the analysis on data assimilation methods to different models, variables and processes.

In particular, as has been reflected in this article, several regional and national gridded products have been developed in the last years for research purposes covering a wide range of applications, resolutions, variables and time periods. However, the main shortcoming that should be pointed out is that most of the high-quality datasets developed at a local, regional or national scale in Spain are rarely shared with the climatological community, leading to redundant analysis in many cases. On the other hand, there are not adequate intercomparison analyses

between the different datasets developed. Finally, recent changes made by the Spanish Meteorological Agency regarding its open data policy, along with the improvements of new or existing tools, could allow the publication of new and updated gridded climatological products in the near future.

References

- Bedia J., Herrera S., Gutiérrez J.M., 2013: Dangers of using global bioclimatic datasets for ecological niche modeling. Limitations for future climate projections, *Global and Planetary Change*, 107, 1-12, doi: 10.1016/j.gloplacha.2013.04.005
- Belo-Pereira M., Dutra E., and Viterbo P., 2011: Evaluation of global precipitation data sets over the Iberian Peninsula, *J. Geophys. Res.*, 116, D20101, doi:10.1029/2010JD015481.
- Birsan M.V., Dumitrescu A., 2014: ROCADA: Romanian daily gridded climatic dataset (1961-2013) V1.0. Administratia Nationala de Meteorologie, Bucuresti, Romania, doi:10.1594/PANGAEA.833627
- Bojinski S., Verstraete M., Peterson T.C., Richter C., Simmons A., Zemp M., 2014: The concept of essential climate variables in support of climate research, applications, and policy. *Bull. Amer. Meteor. Soc.* 95, 1431-14 43, doi:10.1175/BAMS-D-13-00047.1.
- Dumitrescu A., Birsan, M.V., 2015: ROCADA: a gridded daily climatic dataset over Romania (1961-2013) for nine meteorological variables. *Natural Hazards*, 78(2), 1045-1063, doi:10.1007/s11069-015-1757-z
- Durand Y., Brun E., Merindol L., Guyomarc'h G., Lesaffre B., Martin E., 1993: A meteorological estimation of relevant parameters for snow models, *Ann. Glaciol.*, 18, 65-71.
- Durand Y., Giraud G., Brun E., Merindol L., Martin E., 1999: A computer-based system simulating snowpack structures as a tool for regional avalanche forecasting, *J. Glaciol.*, 45(151), 469-484.
- Garzón-Machado V., Otto R., del Arco Aguilar M. J., 2014: Bioclimatic and vegetation mapping of a topographically complex oceanic island applying different interpolation techniques, *Int J Biometeorol*, 58, 887-899, doi: DOI 10.1007/s00484-013-0670-y
- GCOS, 2010a: Implementation plan for the global observing system for climate in support of the UNFCCC (2010 update). GCOS Rep. 138, 186 pp., Available online at www.wmo.int/pages/prog/gcos/Publications/gcos-138.pdf.
- GCOS, 2010b: Guideline for the generation of datasets and products meeting GCOS requirements. GCOS Rep.

143, 12 pp. Available online at www.wmo.int/pages/prog/gcos/Publications/gcos-143.pdf.

Gonzalo J., 2010: Diagnóstico Fitoclimática de la España Peninsular. Hacia un modelo de clasificación funcional de la vegetación y de los ecosistemas peninsulares españoles. Organismo Autónomo de Parques Nacionales-Botánica General, ISBN: 978-84-8014-787-3. In Spanish.

González-Hidalgo J. C., Brunetti M., de Luis M. (2011), A new tool for monthly precipitation analysis in Spain: MOPREDAS database (monthly precipitation trends December 1945–November 2005). *Int. J. Climatol.*, 31: 715–731. doi:10.1002/joc.2115

Gonzalez-Hidalgo J.C., Peña-Angulo D., Brunetti M., Cortesi N. (2015), MOTEDAS: a new monthly temperature database for mainland Spain and the trend in temperature (1951–2010). *Int. J. Climatol.*, 35: 4444–4463. doi:10.1002/joc.4298

Guijarro J.A., Herrera S., 2017, Climatic observations and instrumental reconstructions I: development of high-quality climatic time series.

Gutiérrez J.M., Herrera S., San-Martín D., Sordo C., Rodríguez J.J., Frochoso M., Ancell R., Fernández J., Cofiño A.S., Pons M.R., Rodríguez M.A., 2010: Escenarios Regionales Probabilísticos de cambio climático en Cantabria: Termoplumiometría, Gobierno de Cantabria-Consejería de Medio Ambiente y Universidad de Cantabria, Santander, Spain (In Spanish)

Haylock M.R., Hofstra N., Klein Tank A.M.G., Klok E.J., Jones P.D., New M., 2008: A European daily high-resolution gridded dataset of surface temperature and precipitation. *J. Geophys. Res (Atmospheres)*, 113, D20119, doi:10.1029/2008JD10201

Herrera S., 2011: Desarrollo, Validación Y Aplicaciones de Spain02: Una Rejilla de Alta Resolución de Observaciones Interpoladas Para Precipitación Y Temperatura en España. PhD thesis (in Spanish), Universidad de Cantabria, Cantabria, Spain. <http://www.meteo.unican.es/tesis/herrera>.

Herrera S., Gutiérrez J.M., Ancell R., Pons M.R., Frías M.D., Fernández J., 2012: Development and Analysis of a 50 year high-resolution daily gridded precipitation dataset over Spain (Spain02), *Int. J. Climatol.*, 32, 74-85, doi: 10.1002/joc.2256

Herrera S., Fernández J., Gutiérrez J.M., 2016: Update of the Spain02 Gridded Observational Dataset for Euro-CORDEX evaluation: Assessing the Effect of the Interpolation Methodology, *Int. J. Climatol.*, 36, 900-908, doi: 10.1002/joc.4391

IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.

Isotta F.A., Frei C., Weigluni V., Perčec Tadić M., Lassègues P., Rudolf B., Pavan V., Cacciamani C., Antolini G., Ratto S.M., Munari M., Micheletti S., Bonati V., Lussana C., Ronchi C., Panettieri E., Marigo G., Vertačnik G., 2014: The climate of daily precipitation in the Alps: development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data. *Int. J. Climatol.*, 34: 1657–1675. doi:10.1002/joc.3794

Militino A.F., Ugarte M.D., Goicoa T., Genton M., 2015: Interpolation of daily rainfall using spatiotemporal models and clustering, *International Journal of Climatology*, 35, 1453–1464, doi:10.1002/joc.4068

Miró Pérez J.J., Estrela Navarro M.J., Olcina-Cantos J., 2015: Statistical downscaling and attribution of air temperature change patterns in the Valencia region (1948–2011), *Atmospheric Research*, 156, 189-212

Miró Pérez J.J., Estrela Navarro M.J., Caselles V., Olcina-Cantos J., 2016. Fine-scale estimations of bioclimatic change in the Valencia region, Spain. *Atmospheric Research*, 180, 150-164. doi:10.1016/j.atmosres.2016.05.020

Ninyerola M., Pons X., Roure J.M., 2007: Monthly precipitation mapping of the Iberian Peninsula using spatial interpolation tools implemented in a Geographic Information System. *Theoretical and Applied Climatology*, 89, 195-209

Pérez F.F., Boscolo R., 2010. Climate in Spain: past, present and future. Regional climate change assessment report. Spanish CLIVAR Committee / Thematic network. http://www.clivar.org/sites/default/files/documents/CLIVAR_Spain_2010_0.pdf

Quintana-Seguí P., 2015: SAFRAN analysis over Spain, doi:10.14768/MISTRALS-HYMEX.1388.

Quintana-Seguí P., Peral C., Turco M., Llasat M.C. and Martín E., 2016: Meteorological Analysis Systems in North-East Spain: Validation of SAFRAN and SPAN, *Journal of Environmental Informatics*, 27 (2), 116-130, doi:10.3808/jei.201600335

Quintana-Seguí P., Turco M., Herrera S., Miguez-Macho G., 2017: Validation of a new SAFRAN-based gridded precipitation product for Spain and comparison to

Spain02 and ERA-Interim. *Hydrology and Earth System Sciences*, 21: 2187-2201, doi: 10.5194/hess-21-2187-2017.

Ramos, A. M., Trigo, R. M. and Liberato, M. L. R., 2017, Ranking of multi-day extreme precipitation events over the Iberian Peninsula. *Int. J. Climatol.*, 37: 607–620. doi:10.1002/joc.4726

Rauhe M., Steiner H., Riediger U., Mazurkiewicz A., Gratzki A., 2013: A Central European precipitation climatology Part I: Generation and validation of a high-resolution gridded daily data set (HYRAS), *Meteorologische Zeitschrift*, 22 (3), 235-256, doi: 10.1127/0941-2948/2013/0436

Ruiz-Arias J.A., Pozo-Vázquez D., Santos-Alamillos F.J., Lara-Fanego V. Tovar-Pescador J., 2011: A topographic geostatistical approach for mapping monthly mean values of daily global solar radiation: A case study in southern Spain, *Agricultural and Forest Meteorology*, 151, 1812-1822

Ruiz-Arias J.A., Quesada-Ruiz S., Fernández E.F., Gueymard A., 2015: Optimal combination of gridded and ground-observed solar radiation data for regional solar resource assessment, *Solar Energy*, 112, 411-424

Ruiz-Arias J.A., Arbizu-Barrena C., Santos-Alamillos F.J., Tovar-Pescador J., Pozo-Vázquez D., 2016: Assessing the surface solar radiation budget in the WRF Model: a spatiotemporal analysis of the bias and its causes, *Monthly Weather Review*, 144, 703-711

Sousa P.M., Barriopedro D., Trigo R.M., Ramos A.M., Nieto R., Gimeno L., Turkman K.F., Liberato M.L.R., Impact of Euro-Atlantic blocking patterns in Iberia precipitation using a novel high resolution dataset, *Climate Dynamics*, 2016, 46, 7-8, 2573

Van den Besselaar E.J.M., Haylock M.R., van der Schrier G., Klein Tank A.M.G., 2011: A European Daily High-resolution Observational Gridded Data set of Sea Level Pressure. *J. Geophys. Res.*, 116, D11110, doi:10.1029/2010JD015468
Vicente-Serrano S.M., Saz-Sánchez M.A., Cuadrat J.M., 2003: Comparative analysis of interpolation methods in the middle Ebro Valley (Spain): application to annual precipitation and temperature, *Climate Research*, 24 (2), 161-180, doi: 10.3354/cr024161

Vicente-Serrano S.M., Lanjeri S., López-Moreno J.I., 2007: Comparison of different procedures to map reference evapotranspiration using geographical information systems and regression-based techniques. *Int. J. Climatol.*, 27: 1103–1118. doi:10.1002/joc.1460

Vicente-Serrano S.M., Beguería S., López-Moreno J.I., García-Vera M.A., Stepanek P., 2010: A complete daily precipitation database for northeast Spain: reconstruction,

quality control, and homogeneity. *Int. J. Climatol.*, 30: 1146–1163. doi:10.1002/joc.1850

WMO, 2011: World Meteorological Organization, 2011: Guide to Climatological Practices, WMO-No. 100, ISBN 978-92-63-10100-6, Geneva.