



Is Alzheimer's Disease at Least Partly a Ciliopathy?

Ubaldo Armato^{1*}, Balu Chakravarthy², Anna Chiarini¹, Ilaria Dal Prà¹ and James F. Whitfield²¹Histology & Embryology Section, Department of Life & Reproduction Sciences, University of Verona Medical School, Verona, Venetia, Italy²Molecular Signaling Group, National Research Council of Canada; Institute for Biological Sciences, Ottawa, Ontario, Canada

Abbreviations: A β : Amyloid- β peptide; AD: Alzheimer's Disease; BBS: Bardet-Biedl Syndrome; DGy: Dentate Gyrus; EC: Entorhinal Cortex; GC: Granule Cell; p75NTR: p75 Neurotrophin Receptor; SGZ: Subgranular Zone; SST: Somatostatin; SSTR3: SSTR3 Receptor 3

In the last few years, two events have happened that could profoundly influence our understanding of Alzheimer's disease (AD) that scourge of elderly brains that begins in young brains where it spreads stealthily along characteristic pathways until years later when it clinically surfaces with a shrinking hippocampus and a frighteningly failing memory. The first is the discovery that the hippocampal dentate gyrus (DGy) granule cells (GCs), which initiate the encoding of novel memories, are continuously generated throughout life; and the second discovery is that this 'adult neurogenesis' and memory formation is driven by signals from the DGy precursor GCs' tiny non-motile primary cilia [1]. It follows from this that silencing the dentate GCs' cilia signaling would stop new memory formation. And if this should happen in the AD brain, then AD would at least partly be a "ciliopathy", a name recently coined for several diseases caused by impaired ciliary functions.

Two important AD targets are the hubs of the memory-encoding machinery, the entorhinal cortex (EC) and the DGy. The latter is the gateway to the memory-encoding hippocampus proper. Data from various sensory regions are collected by the pyramidal neurons of the EC and sent via the perforant pathway from the EC to the DGy, which is a very unusual part of the brain [2]. DGy generates GCs throughout life in its subgranular zone (SGZ), which is one of the two neurogenic sites (the olfactorily directed subventricular zone [SVZ] is the other) in adult rodent and human brains [3-5]. This adult neurogenesis provides newborn "blank-state" (i.e., at first spineless) GCs that can form new sets of distal spines and establish new synapses with EC axons [6]. These new neurons have low-threshold long-term potentiation (LTP) responsiveness and, because they lack the Ca²⁺-binding calbindin, they have an enhanced Ca²⁺-driven release of vesicular glutamate at their synapses with CA3 pyramidal cells [9]. Thus, younger GCs are the principal receivers and processors of new data carried by EC axons that otherwise could not easily access the longer-matured GCs with their heavy synaptic load. These young GCs then send streams of the novel "orthogonalized" data along their mossy axons to be folded into coherent memories by the recurrent collaterals of the CA3 pyramidal neurons, which then send the encoded "memories" on to the pyramidal neurons of CA1 region for postprocessing [2]. But a GC's youthful excitability and taste for novelty soon fade as it acquires calbindin to restrain the Ca²⁺-driven glutamate release and a committed synaptic matrix loaded with "neural cement". So now equipped to receive and more slowly transmit old data, it retires from the youth corps and is replaced by a newly formed GC not yet loaded with committed spines and synapses [1,4,5,8]. Thus, the adult brain is kept ready with newborn GCs to register novelty without interfering with the past, and the DGy has armed itself with a remarkable device, the primary

cilium, to help do this. Each GC, at least in rodent brains, has a non-motile primary cilium consisting of a spine (axoneme) of microtubule doublets wrapped in a plasma membrane loaded with p75NTR (p75 neurotrophin receptor) and SSTR3 (somatostatin receptor 3) connected to signaling machineries packed into the tight confines of the GCs' primary cilia along with the Sonic Hedgehog machinery [9-15]. Signals from these cilia drive neurogenesis, maturation, and memory encoding [11,13]. Neurotrophin (e.g., BDNF)-induced p75NTR signaling from the primary cilia drives the proliferation of GC precursors, as preventing this signaling severely reduces neurogenesis [16,17]. SST (somatostatin)-induced signaling from the ciliary SSTR3 receptors acts later in the DGy GC cell life cycle to drive the newborn GCs' participation in memory formation [15,18].

But how might the DGy GCs' primary cilia be involved in the cognitive decline in the AD brain? A model of possibilities is presented in Figure 1. It appears that the accumulating toxic A β 1-42 oligomers in AD brains at first stimulate the proliferation of GC progenitors until the oligomers are locked away in fibrillar plaques, but the newly formed neuroblasts cannot mature or ultimately survive [5,19,20]. The increased progenitor GC proliferation could be caused by the accumulating A β 1-42 oligomers' neurotrophin-like stimulation of ciliary p75NTR [21-25] (Figure 1). But the failure of the newborn neurons to mature and the resulting GC layer shrinkage and memory failure is likely, at least partly, due to the characteristic decline of SST in AD and with it of the ciliary SSTR3 signaling needed for memory encoding [11,26] (Figure 1). A suggestion of ciliary involvement in AD memory loss is the striking shortening of DGy GC primary cilia that accompanies the severely reduced DGy neurogenesis in AD-transgenic mice accumulating both A β 1-42 and tau (as in human AD brains), but not in mice accumulating only A β 1-42 [27,28]. And similar hippocampal dysgenesis and cognitive deficit in humans and model AD mice are also parts of the Bardet-Biedl syndrome (BBS), a known ciliopathy that is due to ciliary failure resulting from the inability to load key structural parts and signalers such as the SSTR3 into the cilia by mutant BBS protein complexes [29-32].

The possibility of primary cilium damage being responsible of the crippling decline of memory formation in AD demands that we must find out whether human DGy GCs cells are ciliated like their rodent

***Corresponding author:** Prof. Dr. Ubaldo Armato, MD, Histology & Embryology Section, Department of Life & Reproduction Sciences, 8 Strada Le Grazie, I-37134, Italy. Tel: 0039-045-8027159; Fax: 0039-045-8027159; E-mail: ubaldo_armato@yahoo.it

Received June 09, 2011; Accepted July 16, 2011; Published July 18, 2011

Citation: Armato U, Chakravarthy B, Chiarini A, Prà ID, Whitfield JF (2011) Is Alzheimer's Disease at Least Partly a Ciliopathy? J Alzheimers Dis Res 1:101e. doi:10.4172/2161-0460.1000101e

Copyright: © 2011 Armato U, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: A schematic of the role of ciliary p75^{NTR} and SSTR3 receptors in dentate gyrus (DGy) neurogenesis. Both these receptors and their activating ligands drive the proliferation and maturation of murine DGy granule cells (GCs). In fact, BDNF, NGF, and Aβ₁₋₄₂ can all activate p75^{NTR}, the signals from which in progenitor GCs trigger the generation of transit amplifying (TA) precursor GCs. Stimulation of immature postmitotic GCs by signals from non ciliary BDNF-TrkB co-receptor complexes [33] and SSTR3s direct the maturation, migration, and incorporation of these cells into the functional GC layer. While the available evidence suggests that Aβ₁₋₄₂ alone would stimulate precursor GCs to proliferate via p75^{NTR} signaling [5,19,20], in Alzheimer's disease (AD) the accumulating Aβ₁₋₄₂ oligomers-hyperphosphorylated tau combination would disorganize and/or disassemble the microtubule structures (MS) [27,28] and prevent the MS and normal holo-BBS complexes from transporting essential components, such as the SSTR3 receptors, into the primary cilia, thereby inhibiting adult neurogenesis and memory formation [9,11,34-36]. Color codes: pink, GC; light yellow, primary cilium; blue arrows, sequential stages of GC functional maturation; black arrows, receptors signaling effects; red arrow, combination of Aβ₁₋₄₂ and tau in AD.

counterparts, and whether these cilia are indeed damaged in AD. If so, then we will know that in a major part of the cases AD is a ciliopathy.

References

- Whitfield JF, Chakravarthy BR (2009) The neuronal primary cilium: driver of neurogenesis and memory formation in the hippocampal dentate gyrus? Cell Signal 21: 1351-1355.
- Rolls ET (2010) A computational theory of episodic memory formation in the hippocampus. Behav Brain Res 215: 180-196.
- Eriksson PS, Perfilieva E, Björk-Eriksson T, Alborn AM, Nordborg C, et al. (1998) Neurogenesis in the adult human hippocampus. Nat Med 4: 1313-1317.
- Kempermann G (2011) Seven principles in the regulation of adult neurogenesis. Eur J Neurosci 33: 1018-1024.
- Schaeffer EL, Novaes BA, Da Silva ER, Skaf HD, Mendes-Neto AG (2009) Strategies to promote differentiation of newborn neurons into mature functional cells in Alzheimer brain. Prog Neuropsychopharmacol Biol Psychiatry 33: 1087-1102.
- Yuste R (2010) Dendritic Spines. The MIT Press.
- Dumas TC, Powers EC, Tarapore PE, Sapolsky RM (2004) Overexpression of calbindin D (28k) in dentate gyrus granule cells alters mossy fiber presynaptic

function and impairs hippocampal-dependent memory. Hippocampus 14: 701-709.

- Aimone JB, Deng W, Gage FH (2010) Put them out to pasture? What are old granule cells good for, anyway...? Hippocampus 20: 1124-1125.
- Berbari NF, Lewis JS, Bishop GA, Askwith CC, Mykityn K (2008) Bardet-Biedl syndrome proteins are required for the localization of G protein-coupled receptors to primary cilia. Proc Natl Acad Sci USA 105: 4242-4246.
- Chakravarthy B, Gaudet C, Ménard M, Atkinson T, Chiarini A, et al. (2010) The p75 neurotrophin receptor is localized to primary cilia in adult murine hippocampal dentate gyrus granule cells. Biochem Biophys Res Commun 401: 458-462.
- Einstein EB, Patterson CA, Hon BJ, Regan KA, Reddi J, et al. (2010) Somatostatin signaling in neuronal cilia is critical for object recognition memory. J Neurosci. 30: 4306-4314.
- Green JA, Mykityn K (2010) Neuronal ciliary signaling in homeostasis and disease. Cell Mol Life Sci 67: 3287-3297.
- Han YG, Spassky N, Romaguera-Ros M, Garcia-Verdugo JM, Aguilar A, et al. (2008) Hedgehog signaling and primary cilia are required for the formation of adult neural stem cells. Nat Neurosci 11: 277-284.
- Händel M, Schulz S, Stanarius A, Schreff M, Erdtmann-Vourliotis M, et al. (1999) Selective targeting of somatostatin receptor 3 to neuronal cilia. Neuroscience 89: 909-926.
- Stanić D, Malmgren H, He H, Scott L, Aperia A, et al. (2009) Developmental changes in frequency of the ciliary somatostatin receptor 3 protein. Brain Res 1249: 101-112.
- Bernabeu RO, Longo FM (2010) The p75 neurotrophin receptor is expressed by adult mouse dentate progenitor cells and regulates neuronal and non-neuronal cell genesis. BMC Neurosci 11: 136.
- Colditz MJ, Catts VS, Al-menhali N, Osborne GW, Bartlett PF, et al. (2010) p75 neurotrophin receptor regulates basal and fluoxetine-stimulated hippocampal neurogenesis. Exp Brain Res 200: 161-167.
- Feinstein JS, Duff MC, Tranel D (2010) Sustained experience of emotion after loss of memory in patients with amnesia. Proc Natl Acad Sci USA 107: 7674-7679.
- van Tijn P, Kamphuis W, Marlatt MW, Hol EM, Lucassen PJ (2011) Presenilin mouse and zebrafish models for dementia: focus on neurogenesis. Prog Neurobiol. 93: 149-164.
- Waldau B, Shetty AK (2008) Behavior of neural stem cells in the Alzheimer brain. Cell Mol Life Sci 65: 2372-2384.
- Chiarini A, Dal Pra I, Whitfield JF, Armato U (2006) The killing of neurons by beta-amyloid peptides, prions, and pro-inflammatory cytokines. Ital J Anat Embryol 111: 221-246.
- Kuner P, Schubeneil R, Hertel C (1998) Beta-amyloid binds to p75NTR and activates NFkappaB in human neuroblastoma cells. J Neurosci Res 54: 798-804.
- Perini G, Della-Bianca V, Politi V, Della Valle G, Dal-Pra I, et al. (2002) Role of p75 neurotrophin receptor in the neurotoxicity by beta-amyloid peptides and synergistic effect of inflammatory cytokines. J Exp Med 195: 907-918.
- Sothibundhu A, Sykes AM, Fox B, Underwood CK, Thangnipon W, et al. (2008) Beta-amyloid(1-42) induces neuronal death through the p75 neurotrophin receptor. J Neurosci 28: 3941-3946.
- Yaar M, Zhai S, Pilch PF, Doyle SM, Eisenhauer PB, et al. (1997) Binding of beta-amyloid to the p75 neurotrophin receptor induces apoptosis. A possible mechanism for Alzheimer's disease. J Clin Invest 100: 2333-2340.
- Burgos-Ramos E, Hervás-Aguilar A, Aguado-Llera D, Puebla-Jiménez L, Hernández-Pinto AM, et al. (2008) Somatostatin and Alzheimer's disease. Mol Cell Endocrinol 286: 104-111.
- Chakravarthy B, Gaudet C, Ménard M, Brown L, Atkinson T, et al. (2011) Changes in the expression of somatostatin 3 receptor (SSTR3) and p75NTR in the hippocampal primary cilia of AD transgenic mice. ICAD Meeting, Paris, July 16-21, Abstract Book.
- Rodríguez JJ, Jones VC, Tabuchi M, Allan SM, Knight EM, et al. (2008) Impaired adult neurogenesis in the dentate gyrus of a triple transgenic mouse model of Alzheimer's disease. PLoS One 3: e2935.
- Baker K, Northam GB, Chong WK, Banks T, Beales P, et al. (2011) Neocortical

- and hippocampal volume loss in a human ciliopathy: A quantitative MRI study in Bardet-Biedl syndrome. *Am J Med Genet A* 155A: 1-8.
30. Berbari NF, Lewis JS, Bishop GA, Askwith CC, Mykytyn K (2008) Bardet-Biedl syndrome proteins are required for the localization of G protein-coupled receptors to primary cilia. *Proc Natl Acad Sci USA* 105: 4242-4246.
31. Davis RE, Swiderski RE, Rahmouni K, Nishimura DY, Mullins RF, et al. (2007) A knockin mouse model of the Bardet-Biedl syndrome 1 M390R mutation has cilia defects, ventriculomegaly, retinopathy, and obesity. *Proc Natl Acad Sci USA* 104: 19422-19427.
32. Bennouna-Greene V, Kremer S, Stoetzel C, Christmann D, Schuster C, et al. (2011) Hippocampal dysgenesis and variable neuropsychiatric phenotypes in patients with Bardet-Biedl syndrome underline complex CNS impact of primary cilia. *Clin Genet*.
33. Young KM, Merson TD, Sotthibundhu A, Coulson EJ, Bartlett PF (2007) p75 neurotrophin receptor expression defines a population of BDNF-responsive neurogenic precursor cells. *J Neurosci* 27: 5146-5155.
34. Jin H, White SR, Shida T, Schulz S, Aguiar M, et al. (2010) The conserved Bardet-Biedl syndrome proteins assemble a coat that traffics membrane proteins to cilia. *C ell* 141: 1208-1218.
35. Zaghoul NA, Katsanis N (2009) Mechanistic insights into Bardet-Biedl syndrome, a model ciliopathy. *J Clin Invest* 119: 428-437.
36. King ME, Kan HM, Baas PW, Erisir A, Glabe CG, et al. (2006) Tau-dependent microtubule disassembly initiated by prefibrillar β -amyloid. *J. Cell Biol.* 175: 541-546.

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 100 Open Access Journals
- 10,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/acrgroup/>