

1 **Genetic diversity of Koala retrovirus (KoRV) *env* gene subtypes: Insights**
2 **into northern and southern koala populations**

3 **Nishat Sarker**^{1,6*}, Jessica Fabijan², Jennifer Seddon¹, Rachael Tarlinton³, Helen Owen¹, Greg
4 Simmons¹, Joshua Thia⁴, Adam Mark Blanchard⁷, Natasha Speight², Jasmeet Kaler³, Richard David
5 Emes^{3,5}, Lucy Woolford², Darren Trott², Farhid Hemmatzadeh², and Joanne Meers^{1*}

6 ¹School of Veterinary Science, The University of Queensland, Australia

7 ²School of Animal and Veterinary Sciences, The University of Adelaide, Australia

8 ³School of Veterinary Medicine and Science, University of Nottingham, United Kingdom

9 ⁴School of Biological Sciences, The University of Queensland, Australia

10 ⁵Advanced Data Analysis Centre (ADAC), University of Nottingham, United Kingdom

11 ⁶Laboratory Sciences & Services Division, International Centre for Diarrhoeal Disease Research,
12 Bangladesh

13 ⁷School of Animal, Rural and Environmental Sciences, Nottingham Trent University, United Kingdom

14

15 *Corresponding authors:

16 (a) Professor Joanne Meers

17 Email: j.meers@uq.edu.au

18

19 (b) Nishat Sarker

20 Email: nishat.sarker@icddr.org; n.sarker@uq.edu.au

21

22

23 **Key words: KoRV, *env* subtype, koala, KoRV-B, genetic diversity**

24

25

26

27 **Repositories:** Sequence reads are uploaded in NCBI database, Sequence Read Archive under accession

28 number SRR8375764

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

34 **Abstract**

35 Koala retrovirus (KoRV) is a recently endogenised retrovirus associated with neoplasia and
36 immunosuppression in koala populations. The virus is known to display sequence variability and to be
37 present at varying prevalence in different populations, with animals in southern Australia displaying
38 lower prevalence and viral loads than northern animals. This study used a PCR and next generation
39 sequencing strategy to examine the diversity of the KoRV *env* gene in both proviral DNA and viral
40 RNA forms in two distinct populations representative of the “northern” and “southern” koala genotypes.
41 The current study demonstrated that the full range of KoRV subtypes is present across both populations,
42 and in both healthy and sick animals. KoRV-A was the predominant proviral subtype in both
43 populations, but there was marked diversity of DNA and RNA subtypes within individuals. Many of
44 the northern animals displayed a higher RNA viral diversity than evident in their proviral DNA,
45 indicating relatively higher replication efficiency of non-KoRV-A subtypes. The southern animals
46 displayed a lower absolute copy number of KoRV than the northern animals as reported previously and
47 a higher preponderance of KoRV-A in individual animals. These discrepancies in viral replication and
48 diversity remain unexplained but may indicate relative protection of the southern population from
49 KoRV replication due to either viral or host factors and may represent an important protective effect
50 for the host in KoRV’s ongoing entry into the koala genome.

51 **1. Introduction**

52 Koala retrovirus (KoRV) is reportedly the youngest endogenized retrovirus (virus integrated in the
53 host’s germline and inherited as part of its host’s genome), having been integrated in the koala genome
54 only about 22,200–49,900 years ago (Ishida et al., 2015). The low genetic diversity of the long terminal
55 repeat (LTR) regions of KoRV provirus sequences suggests that endogenous KoRV probably arose as
56 part of a single outbreak (Ishida et al., 2015). KoRV is also evident in an apparently exogenous
57 (horizontally infectious) form (Hobbs et al., 2017), with many koalas possessing high levels of KoRV
58 RNA in plasma reflecting active viral replication. KoRV is found at a proviral prevalence of 100% in
59 koala populations in northern regions of Australia (Tarlinton et al., 2005; Simmons et al., 2012) and at
60 much lower prevalence (15-25%) in southern populations (Simmons et al., 2012; Legione et al., 2017).
61 This range of prevalence has led to suggestions that KoRV is currently spreading throughout the
62 Australian koala population following a ‘northern to southern’ transmission wave (Fiebig et al., 2006;
63 Tarlinton et al., 2008). Other recent work has indicated that, in southern populations that have
64 apparently low KoRV prevalence, KoRV-negative animals may in fact have defective versions of the
65 virus, missing the *pol* and *env* gene portions most commonly used in virus detection studies (Tarlinton
66 et al., 2017).

67 The outcome of natural KoRV infection is variable, and little is known about the mechanisms of
68 pathogenesis. KoRV is associated with neoplasia and immunosuppression leading to clinical chlamydial
69 disease (Canfield et al., 1988; Hanger et al., 2000; Xu et al., 2013; Fabijan et al., 2017; Gonzalez-
70 Astudillo et al., 2017; Nyari et al., 2017; Burnard et al., 2018). For retroviruses in other species,
71 mutation or recombination events in *env* genes play a significant role in pathogenicity, such as
72 immunosuppression induced by subtypes of feline leukaemia virus (FeLV) (Overbaugh et al., 1988;
73 Anderson et al., 2000; Chandhasin et al., 2005a; Chandhasin et al., 2005b).

74 Classification of KoRV subtypes (in line with similar naming schemes used for better studied
75 gammaretroviruses such as FeLV) is based around the nucleotide sequence of the *env* gene, which
76 encodes the surface protein (SU) and transmembrane protein (TM) of the virus. The 5' end of this gene,
77 known as the hypervariable region, is of particular importance in subtype classification. This region of
78 the *env* gene encodes the protein most exposed to the host's immune response as it is external to the
79 virus membrane and is therefore typically the most variable portion of a retrovirus. This region of the
80 virus also determines host receptor specificity (and therefore cellular tropism) and is one of the major
81 determinants of pathogenicity in FeLV (Bolin and Levy, 2011). The classification system of KoRV is
82 based around phylogenetic groupings of *env* gene nucleotide sequences. For some subtypes (though not
83 all) receptor binding differences have also been determined. The generally accepted classifications are:
84 KoRV-A (Hanger et al., 2000), which binds to the sodium-dependent phosphate transporter Pit-1,
85 KoRV-B and J which bind to the thiamine transporter encoded by THTR1 (Xu et al., 2013), KoRV-C,
86 KoRV-D (Shojima et al., 2013), KoRV-E, KoRV-F (Xu et al., 2015), KoRV-G, KoRV-H, and KoRV-
87 I (Chappell et al., 2017). The receptor usage of subtypes C, D, E, F, G, H and I have not been determined.
88 KoRV-A is found in every KoRV-positive koala and is considered the endogenous version of KoRV
89 from which other subtypes have arisen (Chappell et al., 2017; Hobbs et al., 2017). The other subtypes
90 of KoRV are possibly not germ line transmitted, as they were present in only low copy number in the
91 koala reference genome animal, and as such were considered putative somatic insertions (Hobbs et al
92 2017). The same study also reported that KoRV-D and KoRV-E were present only as defective viruses
93 and the authors hypothesised that these subtypes may be transmitted with a replication competent
94 "helper" virus as has been documented for other retroviruses (Hobbs et al., 2017). KoRV-B is thought
95 to be more pathogenic than KoRV-A, having been reported at an increased prevalence in animals with
96 chlamydiosis or neoplasia than in healthy animals (Waugh et al., 2017). KoRV B and J isolates have
97 also been reported to have variable numbers of copies of repeat regions (these are present as single
98 copies in all KoRV A variants) in their LTRs. These types of repeats are known to enhance replication
99 efficiency in other retroviruses such as FeLV (Xu et al., 2013; Chaban et al., 2017; Waugh et al., 2017).

100 This study explored the evolutionary patterns of KoRV *env* gene subtypes in two koala populations.
101 Patterns of KoRV genetic diversity were investigated in one "northern" genotype in South-East

102 Queensland (QLD) with a KoRV prevalence of 100% and one “southern” genotype in the Mount Lofty
 103 Ranges, South Australia (SA), with an unknown prevalence (Figure 1). Patterns of KoRV *env* subtype
 104 diversity were compared in paired DNA and RNA samples in a subset of koalas from both populations
 105 to understand *env* gene variation in integrated proviral genome (DNA) and in circulating virus (RNA).
 106 Further, this study examined the possible relationship of *env* gene subtypes with a diverse range of
 107 clinical diseases.

108 2. Results

109 This study assessed *env* gene diversity of both integrated KoRV provirus DNA and expressed plasma
 110 viral RNA. Thirty-three “northern” (Queensland, QLD) koalas were assessed, comprising 28 with
 111 paired DNA and RNA samples and five with only DNA as plasma was not available. Twenty-eight
 112 “southern” (South Australia, SA) koalas were included, comprising five with paired DNA and RNA
 113 samples, and 23 with only DNA samples because plasma was not available (10 animals) or the RNA
 114 sample was negative in the *env* gene RT-PCR (13 animals). The overall summary of the demographic
 115 details and clinical status of the animals is shown in Table 1 and details are in supplementary file 2.

116 After quality evaluation and filtration, an average of 22719 total reads were generated for each provirus
 117 DNA sample, ranging between 6169 to 59558 total reads. However, the number of total reads was
 118 higher from amplified RNA, averaging 104894 total reads with a range of 12717 to 245827.

119 **Table 1:** Overall details of study samples (percentages of the total number of animals are given in
 120 parentheses). Not all information was available for all koalas.

Variables	Characters	Queensland	South Australia
Sex	Male	n= 20 (60.6%)	n= 14 (50%)
	Female	n= 13 (39.4%)	n= 14 (50%)
	Total	n= 33	n= 28
Age group	Juvenile	n = 2 (6.3%)	n = 3 (11.5%)
	Young adult	n = 8 (25%)	n = 14 (53.8%)
	Adult	n = 22 (68.8%)	n = 9 (34.6%)
	Total	n = 32	n = 26
Body condition score	0 to <3	n = 15 (46.9%)	n = 4 (15.4%)
	≥ 3 to 5	n = 17 (53.1%)	n = 22 (84.6%)
	Total	n = 32	n = 26
Clinical status	Healthy	n = 11 (33.3%)	n = 5 (17.9%)
	Neoplasia	n = 8 (24.2%)	n = 5 (17.9%)
	Oxalate nephrosis	negative	n = 5 (17.9%)
	Chlamydiosis	n = 16 (48.5%)	n = 11 (39.3%)

121

122 2.1 *Env* subtypes

123 From deep sequencing, a total of 169 unique sequences were generated after sequence validation from
124 all samples of both populations. Sequence reads are available under Sequence Read Archive accession
125 number SRR8375764. The sequence alignment is shown in supplementary file 3. The Bayesian
126 phylogenetic tree (Figure 2) showed high genetic diversity in the KoRV *env* gene at the population
127 level. The identified sequences were grouped with previously recognized subtypes A, B, D, and I.
128 Subtypes B and I were monophyletic in the tree with posterior probability support of 1.0. Subtype A
129 formed a well-supported monophyletic clade. In contrast, subtype D exhibited multiple sub-clades with
130 relatively long branches, high posterior probability values and sequences that were divergent in the
131 hyper-variable region of the *env* receptor binding domain (RBD) (Supplementary Files 3 and 4). This
132 subtype D grouping, with posterior probability support of 0.93, contained reference sequences that had
133 been previously identified as subtypes F, G and H in addition to previously assigned subtype D
134 sequences (Chappell et al., 2017). We identified 13 distinct sub-clades within the subtype D grouping;
135 each of these sub-clades was strongly supported with posterior probabilities of 0.99-1.0. Two of these
136 sub-clades contained only KoRV-G or KoRV-H sequences, and for consistency with the literature, we
137 retained these names despite their phylogenetic placement as sub-clades within subtype D. The
138 remaining 11 sub-clades within subtype D were designated D1 to D11. Sequences that had previously
139 been designated KoRV-F by different authors (Xu et al., 2015; Chappell et al., 2017) belonged to two
140 different subtype D sub-clades in our analysis, with the Chappell et al. (2017) sequence strongly
141 clustering with sub-clade D3 and the Xu et al. (2015) sequence clustering with moderate support
142 (posterior probability 0.83) with D9. Previously assigned KoRV-D sequences clustered either with sub-
143 clade D1 (Shojima et al., 2013) or D4 (Chappell et al., 2017). Sequences clustering in sub-clades D2,
144 D5, D6, D7, D8, 10 and D11 were not matched or clustered with any reference sequences. Subtypes C
145 and E, which have been previously described (Miyazawa et al., 2011; Shojima et al., 2013; Xu et al.,
146 2015), clustered together close to but distinct from group D.

147 Sequences in this study were assigned to all of the afore-mentioned subtypes except for subtypes C, E
148 and H, which were not found in any of our samples. A total of 63 sequences were found from the KoRV-
149 A subtype, 22 from subtype B, 16 from subtype I, and 68 from subtype D. Within the subtype D
150 sequences, most (22) sequences were within the D1 sub-clade, with one sequence in each of D4 and
151 D10, two in each of D6, D8 and KoRV-G, four sequences in each of D2 and D7, five in D3, six in D5,
152 eight sequences in D9, and 11 sequences in the D11 sub-clade.

153 The average read count of each unique sequence was highly diverse between animals, between RNA
154 and DNA forms in the same animal and within the two koala populations (QLD and SA). The number
155 of different KoRV DNA sequences within an individual, indicative of the number of provirus insertions,

156 was significantly higher (Mann Whitney U, $p < 0.0001$) in QLD individuals (median 77, range 63-100)
157 in comparison to SA koalas median 59, range 43 – 74).

158 The read count details of unique sequences in individual animals are available in Supplementary File 5
159 and the relative percentage levels of each subtype (group of sequences) are available in Supplementary
160 File 6. There was variation among individuals in overall read count and so read counts of subtypes were
161 converted to the relative percentage of each subtype of the total reads for that individual with following
162 equation:

163 Relative percentage of each specific subtype (for each individual) = (total number of unique
164 sequence reads of the subtype / total read count of all subtypes) x 100

165 **2.2 Env subtype abundance in QLD and SA koala population**

166 As shown in Figure 3, the absolute read count values of *env* gene subtypes in the DNA of QLD koalas
167 were significantly higher (p value < 0.0001) in comparison to SA koalas. Each koala had multiple *env*
168 subtypes in their genome (Figure 4). In the QLD animals, KoRV-A, KoRV-B, and KoRV-D (sub-clades
169 10 and 11) were present in proviral DNA form at some level in all individuals (supplementary file 5
170 and 6) KoRV-A was the dominant subtype (present at $>40\%$ of reads) in proviral DNA in 30 of the 33
171 QLD animals (Figure 4A). The exceptions were two animals where the D1 subclade was in higher
172 abundance and one koala with the D6 subclade as the highest abundance. For the SA koalas, subtypes
173 A, B and sub-clade D10 were present in the DNA of all koalas, while sub-clades D1 (27/28) and D11
174 (27/28) were also represented in the majority of koalas. KoRV-A dominated the proviral DNA subtypes
175 in all SA koalas with a much higher relative percentage than in the QLD animals. The median Shannon
176 diversity index was also significantly lower in the SA than the QLD animals (Mann Whitney U
177 < 0.0001).

178 The subtypes present in the RNA of individual animals differed from those present in the DNA. The
179 median Shannon diversity index was lower for DNA than RNA samples for both populations but did
180 not reach significance (Mann Whitney U test, QLD $p=0.4$ and SA $p=0.5$). For the QLD animals subtypes
181 A, B, D2, D3 and D5 were more abundant in the DNA samples and D10 and G more abundant in the
182 RNA samples (FDR p values > 0.005). All QLD ($n=28$) and SA koalas ($n=5$) had subtypes or subclades
183 A, B, D1, D10 and D11 at some level in their viral RNA; additionally all SA koalas also had subclades
184 D2 and D3.

185 All five SA koalas showed a very high relative percentage of KoRV-A (92.5-99.9%) in viral RNA. In
186 contrast, none of the RNA samples for QLD koalas showed KoRV-A to be the most abundant subtype,
187 with KoRV-B ($n=8$), D11 ($n=7$), D1 ($n=3$), D3 ($n=2$), KoRV-I ($n=1$), D2 ($n=1$), D5 ($n=2$), D6 ($n=2$),

188 D7 (n=1) or D8 (n=1) the most abundant subtypes or subclades within individual koalas (supplementary
189 file 6)

190 **2.3 Distribution differences between viral DNA and RNA of env subtypes**

191 The difference in the distribution of subtypes between viral DNA and RNA of individual koalas was
192 striking, in particular amongst the QLD koalas (Figure 4A and 5 A,C). In some koalas, the predominant
193 viral RNA subtype formed only a very minor proportion of the proviral DNA subtype distribution. As
194 examples, koalas Q2 and Q27 had an overwhelming predominance of subtype/subclades D2 and D8 in
195 their viral RNA, comprising 85% and 88% of their RNA subtype distribution, respectively, whereas
196 these two subtypes comprised only 13% and 8%, respectively, of the proviral DNA subtype distribution
197 in these koalas. Within individual koalas, it is clear that some KoRV proviral subtypes have very high
198 rates of expression while others are poorly expressed.

199 These results probably reflect a greater replication rate (and overall viral load) in the QLD animals with
200 viral diversity increasing in the RNA form of the virus (a greater number and range of non KoRV-A
201 subtypes being produced). In the SA animals where the viral load (and presumably the replication rate)
202 is lower this difference in viral diversity is not seen, with these animals continuing to display a higher
203 preponderance of the ancestral A subtype (both when compared to the QLD animals and when RNA
204 and DNA forms within the SA animals are compared).

205 **2.4 KoRV-A and KoRV-B status based on conventional PCR**

206 Conventional PCR of DNA using KoRV-A and KoRV-B specific primers demonstrated a 100%
207 prevalence of KoRV-A in QLD koalas and 96.4% in SA koalas, while the KoRV-B prevalence was
208 48.5% in QLD and 0% in SA koalas. This is in contrast to the MiSeq deep sequencing results where all
209 animals in both populations were positive for both subtypes. There was a significant differences
210 ($p < 0.0001$) between a positive test for KoRV B with conventional PCR and a Miseq read count of
211 > 2700 for KoRV B. With one exception, koalas with raw read counts below 2700 were negative by
212 KoRV-B specific conventional PCR.

213 **2.5 Subtype correspondence with clinical status of respective koalas**

214 Amongst the 33 QLD koalas, 11 were clinically healthy, 13 had chlamydiosis, four had neoplasia, four
215 had both chlamydiosis and neoplasia and one had a non-neoplastic hepatic mass (Table 1 and
216 Supplementary File 2). All DNA and RNA subtypes, including the putative pathogenic KoRV-B, were
217 found in both healthy and diseased animals. Of the 28 SA koalas, 12 had chlamydiosis, five had
218 neoplasia, five had oxalate nephrosis (a genetic kidney disease not commonly found in QLD animals),

219 and six were healthy. As with the QLD animals, all SA koalas had both KoRV-A, KoRV-B and
220 subclades of KoRV-D.

221 There were too few animals (particularly in the SA population) with RNA for a sensible analysis of
222 disease status vs viral subtypes. For proviral DNA there was no clear association between the abundance
223 of any particular subtype and any particular disease syndrome. Graphs of subtypes A, B, combined D
224 and I versus disease categories of healthy, neoplasia, oxalate nephrosis and chlamydiosis for each
225 population are presented in supplementary file 7. There was a trend towards healthy animals (in both
226 the QLD and SA populations) and the oxalate nephrosis animals in the SA population (this disease is
227 thought to have a genetic basis) having a lower viral diversity (with a greater preponderance of KoRV
228 A) than those with neoplasia or chlamydiosis (the diseases thought to be associated with KoRV
229 infection) though a major confounding factor for more robust analysis here was the number of animals
230 with multiple disease syndromes and the small number of animals in some disease categories in each
231 population.

232 **3. Discussion**

233 Despite the high prevalence of KoRV and its potential impact on the health of koalas, there are few
234 reports available about KoRV genetic diversity in the Australian koala population. Most of the
235 information about KoRV diversity comes from studies in overseas captive koalas (Miyazawa et al.,
236 2011; Shojima et al., 2013; Xu et al., 2013; Xu et al., 2015) or wild South-East Queensland (SE QLD)
237 koalas (Chappell et al., 2017), all of which are of the “northern” or mixed genotypes. Here, we made a
238 substantial contribution to knowledge in this field by investigating KoRV *env* gene diversity in diseased
239 and healthy koalas from both “northern” and “southern” genotype populations (SE QLD and Mt Lofty
240 Ranges, SA) highlighting the differences in abundance of KoRV subtypes at both DNA and RNA level
241 between these populations, with the southern animals demonstrating both a lower viral load, a reduced
242 viral diversity and a greater preponderance of KoRV A abundance. The paired DNA-RNA samples in
243 individual koalas also demonstrated that the abundance of different DNA and RNA subtypes within
244 individual koalas do not correspond to each other, with a trend (though not significant) towards a higher
245 diversity in the RNA samples, indicating variable expression of proviral DNA subtypes.

246 Our study demonstrated that the full range of KoRV subtypes was present in both northern and southern
247 koala populations, and in both DNA and RNA forms of the virus. The finding of KoRV-B in all southern
248 animals studied was unexpected and is in contrast to recent PCR-based studies of KoRV-B prevalence
249 which have reported varying (Waugh et al., 2017) or absent (Legione et al., 2017) prevalence rates of
250 KoRV-B in “southern” animals.

251 Phylogenetic analysis of the KoRV *env* genes in this study found four major subtypes; three were
252 strongly supported monophyletic clades clustering with previously designated as A, B and I. The fourth
253 subtype was the large paraphyletic group D, which this study classified into 13 sub-clades comprising
254 previously designated subtypes G and H and newly designated subtypes D1 to D11. Two reference
255 sequences that had previously been designated KoRV-F clustered with two distinct group D subtypes
256 (KoRV-D3 and KoRV-D9). The paraphyletic nature of the subtype D grouping highlights the
257 difficulties of assigning KoRV subtypes. Rather than following convention and designating our newly
258 identified sequences as further alphabetical subtypes (KoRV-K, L, M, N, etc), we recognised that these
259 sequences belong to a large phylogenetic grouping and should not be classified as distinct lettered
260 subtypes, but rather as sub-clades of subtype D. We cannot entirely rule out PCR related recombination
261 of differing loci or PCR based errors in the sequences (particularly for the KoRV-D group) though the
262 parameters set for including sequences in subsequent phylogenetic analysis (sequences present in at
263 least two animals, a minimum read count of four and a 99% clustering threshold) will have removed
264 sequences that appeared only once in the data.

265 KoRV-A, KoRV-B, and KoRV-D sub-clades 1, 10 and 11 were highly prevalent in individuals in this
266 study, while KoRV-C, E and H were not identified. KoRV-C was identified at a Japanese zoo from
267 captive koalas (Shojima et al., 2013) and to date has not been found in any wild koala (Chappell et al.,
268 2017; Hobbs et al., 2017). KoRV-E was identified from a zoo in USA (Xu et al., 2015) and was also
269 not found by Chappell et al. (2017), although a defective form is present in the reference genome animal
270 (Hobbs et al., 2017). KoRV-H is rare, having been found in only one animal in viral RNA form
271 (Chappell et al., 2017); in our study, KoRV-H sequences clustered within a larger KoRV-D clade, so
272 it is possible that this subtype may exist in other geographic ranges. Overall these data highlight the
273 extreme intra-animal variability of KoRV with many subtypes being reported in only a small number
274 of animals.

275 This study is not able to distinguish between endogenous (incorporated into the genome and vertically
276 transmitted) and exogenous (horizontally transmitted) virus. Indeed, in a newly integrated virus like
277 KoRV, this distinction may not be very helpful as there is no reason why the virus cannot be both
278 vertically and horizontally transmitted. The original demonstration of KoRV as an endogenous virus
279 (Tarlinton et al 2006) did not use methods that would distinguish the different KoRV subtypes, though
280 subsequent analysis of the variants present in the reference genome animal (of the northern genotype)
281 (Hobbs et al., 2017) indicated that KoRV-A is endogenous in this animal (present at high copy number)
282 while variants B-I are likely present only as low copy number somatic cell insertions (and so likely not
283 vertically transmitted) though to date this has only been examined in this one individual. A number of
284 sequencing efforts from museum specimens have only demonstrated KoRV-A and not the other variants
285 in historical specimens (though DNA quality is an issue in these specimens). There has been limited

286 sequencing of KoRV strains outside of the variable region of *env*; analysis of the koala reference
287 genome indicated that variants D and E were defective in the source animal (Hobbs et al., 2017). This
288 does not mean that these variants are not horizontally transmitted as there are multiple examples of
289 retroviruses in other species (notably cats and chickens) where defective viruses are transmitted
290 alongside replication competent “helper viruses”. KoRV-A might represent remnants of ancestral germ-
291 line infections by exogenous retroviruses with other forms of the virus representing those still active
292 due to continual reinfection or retro-transposition in cis within germ-line cells as reported in other
293 retroviral systems (Boeke and Stoye, 1997; Belshaw et al., 2005). This theory is potentially supported
294 by the phylogenetic pattern evident in the KoRV-A isolates in this study, with the long branch lengths
295 obvious within the KoRV-A cluster (supplementary figure 4) consistent with very closely related
296 endogenous proviruses that have diverged post integration.

297 KoRV-A is at any rate present in all KoRV positive koalas and is consistent with being endogenous
298 (Xu et al., 2015; Legione et al., 2017; Waugh et al., 2017). This study confirms KoRV-A as being
299 present in all KoRV positive koalas as previously reported by many groups. It also highlights the
300 previously described lower viral load in southern animals (Legione et al., 2017, Simmons et al., 2012).
301 This lower viral load corresponds with a reduced viral diversity and a higher relative percentage
302 abundance of KoRV-A in southern koalas. In addition, in the northern animals, the relative percentage
303 of KoRV-A was much higher in the integrated proviral DNA than in the viral RNA, with other subtypes
304 variable among all samples. This may reflect a relatively greater viral diversity in animals with higher
305 viral loads due to the greater rate of mutation in actively replicating virus. This phenomenon is well
306 described in other retroviruses such as HIV where viral diversity increases with viral replication (Theys
307 et al., 2018). Alternatively, some variants of KoRV-A (particularly endogenous loci) may not be very
308 effectively transcribed, either directly or as a result of competition with high copy number of other
309 transcribed subtypes. Another possibility is that a mutation of the provirus may disrupt DNA sequence
310 elements from the promoter which are essential for transcription. Indeed, the original KoRV-A isolate
311 does not replicate efficiently in cell culture, probably due to sequence changes in its LTR when
312 compared with more replication competent clones (Shimode et al., 2014). It is also possible that
313 transcription from individual KoRV-A loci is uneven with some highly transcribed loci responsible for
314 the RNA detected. These particular loci may be less prevalent in QLD animals. Another potential
315 confounding factor in blood samples is that the levels may not directly reflect viral transcription in other
316 tissues (there is likely differential transcription in different tissues as has been demonstrated for many
317 retroviruses, both endogenous and exogenous). However, replicating virus in any tissue likely produces
318 virions spilling over in the blood.

319 Several other studies have reported a linkage between detection of KoRV-B provirus and neoplasia or
320 chlamydial disease occurrence (Chaban et al., 2017; Waugh et al., 2017) (Xu et al., 2013). However,

321 the current study does not support an association between the presence of particular virus subtypes in
322 either DNA or RNA forms and the occurrence of disease. Previous studies on the association between
323 KoRV-B subtype and disease were based on conventional PCR. The NGS approach used here is not
324 reliant on sequence specific primers for each KoRV subtype and is therefore able to detect a more
325 comprehensive range of subtypes in individual animals. The NGS approach was also more sensitive in
326 detection of KoRV-B, with only 48.5% of QLD animals and no SA animals testing positive for KoRV-
327 B with conventional PCR, in contrast to 100% of the same animals testing positive with the PCR and
328 NGS approach. There was a significant association between higher read counts of KoRV-B in the NGS
329 data and the likelihood of testing positive for KoRV-B on conventional PCR. These findings indicate
330 that previously reported results for an association of KoRV-B with disease in animals might reflect an
331 association between higher viral load and disease rather than the presence of the KoRV-B subtype per
332 se. In terms of koala population management decisions, the findings also indicate that testing for KoRV-
333 B via endpoint PCR, as has been adopted by some zoological collections, is probably not a useful screen
334 for future neoplasia risk.

335 Indeed, this study does not provide convincing evidence for an association of any particular virus
336 subtype with a particular disease syndrome, although a trend towards a reduced viral diversity and an
337 increased preponderance of KoRV-A is evident in healthy animals and in the SA animals the oxalate
338 nephrosis animals (this is a genetic disease seen predominantly in the southern population). As with the
339 differences in viral diversity between the populations this probably reflects the previously reported
340 relationship between higher viral load (and therefore sequence diversity), neoplasia and clinical
341 chlamydiosis in KoRV affected animals (Tarlinton et al., 2005). Important caveats where are the small
342 numbers of animals in some disease categories, the differences in disease patterns between the two
343 populations and the numbers of animals with multiple diseases which will have confounded this
344 analysis. These confounding factors also made more appropriate statistical analysis techniques for this
345 type of data (like multivariate modelling) inaccurate.

346 It still remains unexplained why the South Australian animals have a lower level of KoRV replication
347 and a reduced level of abundance of non-KoRV-A subtypes. This study only looked at *env* diversity
348 and there are other factors that can affect viral replication efficiency. In particular the LTR sequences
349 of retroviruses are known to be major determinants of replication efficiency (Pantginis et al., 1997;
350 Chandhasin et al., 2004) and variations in KoRV-B/J isolate LTRs have been reported previously
351 (Shimode et al., 2014) Hobbs et al 2017) that appear to affect replication efficiency. It is also possible
352 that the South Australian population has defects in the receptor for one or more variants of KoRV
353 (KoRV-A and B are known to use different receptors) affecting the efficiency of viral re-infection in
354 these animals, although preliminary analysis of unpublished transcriptome sequences from the two
355 populations would indicate that this is not the case. Other unpublished data indicate that the SA animals

356 may have a defective form of the virus, which is missing most of the *gag*, *pol* and *env* genes (Tarlinton
357 et al., 2017) and it is possible that this defective virus inhibits replication of the full length virus as has
358 been reported for some other retroviruses (Boeke and Stoye, 1997).

359 Overall, this study analysed KoRV *env* gene diversity in paired samples of provirus DNA and viral
360 RNA within individual koalas from two different zones of koala habitat representing northern and
361 southern koala populations. The identified sequences significantly enhance the number of *env* gene
362 sequences known for KoRV and highlighted significant variation between the abundance of transcribed
363 variants of KoRV present in the RNA of individuals when compared with the provirus complement in
364 the DNA. This probably reflects differential transcription efficiency of different loci and subtypes.
365 KoRV-A is the likely ancestral version of KoRV, with other variants likely generated via mutations,
366 deletions, or recombination events. These other subtypes have now become the predominant transcribed
367 form of KoRV in the Queensland population. It remains unexplained why the southern animals display
368 such lower viral loads and reduced viral diversity than the northern population, along with such a
369 different disease pattern, however this study highlights that this is not as simple as the presence or
370 absence of particular virus subtypes as has been previously hypothesised.

371 **4. Methods**

372 **4.1 Sample collection and preparation**

373 In South-East QLD, animals were sourced from Moggill Koala Hospital, Australia Zoo Wildlife
374 Hospital, RSPCA Wacol and Sea World Paradise Country. South Australian (Mount Lofty Ranges)
375 samples were collected from Fauna Rescue of South Australia (Figure 1, Table 1). Blood (2-3 ml) was
376 collected from live and clinically healthy captive koalas using a sterile 22-gauge butterfly catheter and
377 5 ml syringe. Wild koalas hospitalised due to disease or serious injury following trauma or animal attack
378 were euthanased and necropsied. Koalas were anaesthetised with 0.25 ml Zoletil (Virbac)
379 intramuscularly. Euthanasia was performed with an intravenous injection of pentobarbitone.
380 Immediately following euthanasia, 10 – 15 ml of blood was withdrawn from a femoral vein or by
381 cardiac puncture into EDTA tubes.

382 DNA was extracted from 100 µl EDTA whole blood using Qiagen DNeasy Blood & Tissue Kit
383 according to manufacturer's (Qiagen) instructions. A 1-2 ml aliquot of blood was centrifuged at 3000
384 g for 5 mins and 200 µl of plasma was removed and added to 300 µl of RNAlater stabilisation agent
385 (Qiagen) within 15 min of blood collection. RNA was extracted using Qiagen QIAmp Viral RNA mini
386 kit with on-column Qiagen RNase free DNase steps. Briefly, 140 µl of RNAlater diluted plasma was
387 suspended in 560 µl viral lysis buffer containing carrier RNA and extraction continued following the

388 extraction kit procedures and finally eluted in 30 µl water. The extracted RNA samples were stored at
389 -80°C until required.

390 **4.2 KoRV-A and KoRV-B real-time and conventional PCR**

391 To test for the presence of KoRV-A and B, we used real time and conventional PCR. Initially, KoRV
392 positivity of the extracted DNA and RNA was initially assessed with a real-time PCR of the KoRV *pol*
393 gene using published primers and probe (Tarlinton et al., 2005). DNA and RNA samples were amplified
394 using TaqMan gene expression master mix (Applied Biosystem) and SuperScript® III One-Step RT-
395 PCR System with Platinum® Taq DNA Polymerase (Invitrogen) respectively, following
396 manufacturers' instructions, in a BioRad CFX 96. Samples were considered KoRV-positive if the CT
397 value was ≤ 35 .

398 Conventional PCR of the *env* gene was performed using published primers to specifically amplify each
399 of the KoRV-A and KoRV-B *env* subtypes (Waugh et al., 2017) as a preliminary assessment of KoRV
400 subtype prevalence. Primers used in this study shown in Table 2. The Qiagen HotStartTaq Plus Master
401 Mix kit was used for PCR of DNA samples following the manufacturer's instructions with 35 cycles of
402 amplification and an annealing temperature of 51°C. KoRV-A and KoRV-B positive samples were
403 directly purified with ExoSAP-IT (Thermo Fisher Scientific), following manufacturer's directions and
404 Sanger sequenced to validate the amplification of this subtype. Sequencing was undertaken using Big
405 Dye Terminators (ThermoFisher Scientific) at the Animal Genetics Laboratory, University of
406 Queensland. Sequences were subjected to BLAST analysis through the NCBI database to determine the
407 percentage of homology to known subtypes.

408 **Table 2:** Primers used in this study for PCR

Region	Forward	Reverse	Reference
Pol	TTGGAGGAGGAATACCGATTACAC	GCCAGTCCCATACCTGCCTT	(Tarlinton et al., 2005)
Env KoRV-A specific	TCCTGGGAACTGGAAAAGAC	GGGTTCCCAAGTGATCTG	(Waugh et al., 2017)
Env KoRV-B specific	TCCTGGGAACTGGAAAAGAC	GGCGCAGACTGTTGAGATTC	(Waugh et al., 2017)

409 **4.3 Sample preparation for Illumina sequencing**

410 Previously published oligonucleotide primers flanking the hypervariable region of the *env* gene
411 (Chappell et al., 2017) were used to amplify a 500 bp fragment of target sequence by PCR. The primers
412 contained the Illumina adaptor sequences (*italics*) ligated to *env* gene complementary regions.

413 The Qiagen HotStartTaq Plus Master Mix kit was used to amplify from DNA and the Qiagen OneStep
414 RT-PCR kit was used to amplify from RNA, both following the manufacturer's instructions with an
415 annealing temperature of 58°C and 35 rounds of amplification. We adopted recently established deep
416 sequencing methodology for analysis of the *env* gene hypervariable region, such that consistency was
417 retained between current and previous findings (Chappell et al., 2017). Samples were prepared
418 following the Illumina 16S Metagenomic Sequencing Library Preparation guidelines. The purification
419 and sequencing of PCR amplicons was performed at the Ramaciotti Centre for Genomics (University
420 of New South Wales, Sydney, Australia). Purification was performed using Agencourt AMPure XP
421 beads (Beckman Coulter, USA) and purified DNA was indexed with unique 8 bp barcodes using the
422 Illumina Nextera XT 384 sample Index Kit A-D (Illumina FC-131-1002) following standard PCR
423 conditions. Indexed amplicons were pooled together in equimolar concentrations and sequenced on the
424 MiSeq Sequencing System (Illumina, USA) using paired end sequencing with V3 300bp following
425 manufacturer's protocols.

426 **4.4 Sequence assembly**

427 The overlapped forward and reverse reads of the Illumina next generation sequencing (NGS) were
428 assembled using the OL assembly method in the *dDocent* pipeline (Puritz et al., 2014). The reads were
429 trimmed using Trimmomatic (Bolger et al., 2014) and assembled using *Rainbow* (Chong et al., 2012)
430 and *CD-HIT* (Fu et al., 2012). A series of optimization assemblies were run to assess how the number
431 of contigs would be affected by parameter choice in *dDocent* and the effect of clustering threshold: (a)
432 the minimum number of samples in which a sequence had to be represented (1–10 samples); and (b)
433 the clustering threshold (80–98%). These preliminary optimization runs indicated that the level of
434 clustering did not have a substantial effect on the total number of unique contigs assembled. What did
435 have a major impact was the number of individuals required to represent a sequence: when this was 1,
436 the number of contigs that could be assembled from a single individual was substantially larger than
437 when assemblies required a sequence to be found in ≥ 2 samples. This result implicates considerable
438 subtype sequence variation, but it is hard to resolve this from technical or sequencing error that may
439 generate false variation. Therefore, the final assembly included the parameter selection of: (i) sequences
440 present in at least two samples, (ii) a minimum read count of four and (iii) a 99% clustering threshold.
441 Graphical view of optimisation assemblies are shown in supplementary file 1.

442 The representative sequences were aligned with KoRV-A (AF151794) and KoRV-B (KC779547.1)
443 using the *ClustalW* alignment in the program *BioEdit* (Hall, 1999) to identify the presence of any
444 anomalous contigs. Sequences that failed to show homology against reference sequences were removed
445 from further analysis. The putative sequences were mapped by *BWA* (Li and Durbin, 2009) with the
446 following parameters: match score = 1, mismatch penalty = 4, gap open penalty = 15. Finally, *SAMTools*

447 (Li et al., 2009) was used to filter the alignment bam files (for a MapQ score of 30) and to extract the
448 counts of reads mapped to each contig.

449 **4.5 Phylogenetic analysis**

450 The representative unique sequences were imported into the Geneious v11.0.4 software package
451 (<https://www.geneious.com/>) and combined with previously published KoRV env sequences; KoRV-A
452 (AF151794, KX587957.1 and KP792565.1), KoRV-B (KX588002.1, KX588011.1, KX588027.1,
453 KC779547.1, AB822553.1, KX588031.1, KX588053.1), KoRV-C (AB828005.1, KP792564.1),
454 KoRV-D (KX587952.1, KX587991.1, KX588043.1, KX587993.1, KX587972.1, KX587972.1,
455 AB828004.1, KX587997.1), KoRV-E (KU533853.1), KoRV-F (KX588025.1, KX588028.1,
456 KX587994.1, KU533852.1), KoRV-G (KX587961.1 and KX587998.1), KoRV-H (KX588036.1 and
457 KX587979.1), and KoRV-I (KX587976.1 and KX588021.1) were used. Moreover, four other
458 sequences from the viruses previously determined to be the most closely related to KoRV (Simmons et
459 al., 2014) in other were used as outgroups to root the KoRV phylogeny: two Gibbon ape leukaemia
460 virus (GALV) sequences, (KT724047.1, KT724048.1) and two *Melomys burtoni* retrovirus (MbRV)
461 sequences (KF572486.1, KF572485.1). Sequences were aligned using *ClustalW* alignment with a gap
462 opening cost of 15 and a gap extension cost of 7 as implemented in Geneious 11.0.4. The alignments
463 were further edited by hand to fill the blanks at the beginning and end.

464 A Bayesian phylogenetic tree was determined from the aligned reads using the Geneious plugin of
465 MrBayes 3.2.6 (Huelsenbeck and Ronquist, 2001) with a chain length of 10,100,000, a subsampling
466 frequency of 2000 and a burn-in-length of 1,100,100, with all others parameters set at defaults.
467 Sequences were manually allocated to a KoRV subtype based on clustering with previously identified
468 reference subtypes and phylogenetic topology.

469 **4.6 Statistical analysis**

470 The comparison of the number of unique proviral sequences and read counts between QLD and SA
471 populations was statistically evaluated through non-parametric Mann Whitney U test. The concordance
472 of the NGS results with the conventional PCR testing for KoRV-B was assessed through Pearson's chi-
473 squared tests. The taxonomic count data was analysed for statistically significant differences, between
474 QLD vs SA provirus, and DNA vs RNA for both SA and QLD samples, in R (Ihaka and Gentleman,
475 1996) using the EdgeR wrapper (Robinson et al., 2010) as part of the phyloseq package (McMurdie and
476 Holmes, 2013). Diversity statistics were calculated using vegan (Dixon, 2003) and differences were
477 assessed for significance using Mann-Whitney U tests in Prism 8.01 (GraphPad Software Inc. USA).

478

479 **FUNDING INFORMATION**

480 This project and scholarship for NS were funded by the Queensland Department of the Environment
481 and Heritage Koala Research Grant Programme 2012. NS was also supported by a Keith Mackie Lucas
482 Travel Scholarship from the University of Queensland.

483

484 **CONFLICTS OF INTEREST**

485 The authors declare that there is no conflict of interest.

486

487 **ETHICAL STATEMENTS**

488 Ethical approval for this study was granted by the University of Queensland Animal Ethics Committee,
489 permit number ANFRA/SVS/461/12 and ANRFA/SVS/445/15, the Queensland Government
490 Department of Environment and Heritage Protection permit number WISP11989112, University of
491 Adelaide Animal Ethics Committee permit number S-2013-198 and South Australian Government
492 Department of Environment, Water and Natural Resources Scientific Research Permit Y26054.

493

494 **AUTHOR CONTRIBUTIONS**

495 N.S. performed DNA and RNA extraction, laboratory experiments, data analysis and drafted
496 manuscript. J. Meers, J.M.S., G.S. and H.O. helped in laboratory experiment set up, data
497 interpretation and manuscript preparation. J. T. helped in bioinformatics analysis. R.D.E and R.T
498 edited the manuscript. J.F. and N. Speight helped in sample collection and reviewing manuscript.
499 J.K. and A.B.M reviewed the statistical analysis and edited the manuscript. F.H, D.T. and L.W.
500 reviewed the manuscript. All authors read and approved the final manuscript.

501

502 **REFERENCES**

- 503 Anderson, M.M., Lauring, A.S., Burns, C.C., Overbaugh, J., 2000. Identification of a cellular
504 cofactor required for infection by feline leukemia virus. *Science* 287, 1828-1830.
- 505 Belshaw, R., Katzourakis, A., Pačes, J., Burt, A., Tristem, M., 2005. High Copy Number in
506 Human Endogenous Retrovirus Families is Associated with Copying Mechanisms in
507 Addition to Reinfection. *Molecular Biology and Evolution* 22, 814-817.
- 508 Boeke, J.D., Stoye, J.P. 1997. Retrotransposons, Endogenous Retroviruses, and the Evolution
509 of Retroelements, In: Coffin, J.M., Hughes, S.H., Varmus, H.E. (Eds.) *Retroviruses*.
510 Cold Spring Harbor Laboratory Press, Cold Spring Harbor (NY).
- 511 Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
512 sequence data. *Bioinformatics* 30, 2114-2120.
- 513 Bolin, L.L., Levy, L.S., 2011. Viral determinants of FeLV infection and pathogenesis: lessons
514 learned from analysis of a natural cohort. *Viruses* 3, 1681-1698.
- 515 Burnard, D., Gillett, A., Polkinghorne, A., 2018. Chlamydia pecorum in Joint Tissue and
516 Synovial Fluid of a Koala (*Phascolarctos cinereus*) with Arthritis. *Journal of wildlife*
517 *diseases*.

518 Canfield, P.J., Sabine, J.M., Love, D.N., 1988. Virus particles associated with leukaemia in a
519 koala. *Australian veterinary journal* 65, 327-328.

520 Chaban, B., Ong, V.A., Hanger, J., Timms, P., 2017. Molecular dynamics and mode of
521 transmission of Koala Retrovirus (KoRV) as it invades and spreads through a wild
522 Queensland koala population. *Journal of virology*.

523 Chandhasin, C., Coan, P.N., Levy, L.S., 2005a. Subtle mutational changes in the SU protein of
524 a natural feline leukemia virus subgroup A isolate alter disease spectrum. *J Virol* 79,
525 1351-1360.

526 Chandhasin, C., Coan, P.N., Pandrea, I., Grant, C.K., Lobelle-Rich, P.A., Puetter, A., Levy,
527 L.S., 2005b. Unique long terminal repeat and surface glycoprotein gene sequences of
528 feline leukemia virus as determinants of disease outcome. *J Virol* 79, 5278-5287.

529 Chandhasin, C., Lobelle-Rich, P., Levy, L.S., 2004. Feline leukaemia virus LTR variation and
530 disease association in a geographical and temporal cluster. *J Gen Virol* 85, 2937-2942.

531 Chappell, K.J., Brealey, J.C., Amarilla, A.A., Watterson, D., Hulse, L., Palmieri, C., Johnston,
532 S.D., Holmes, E.C., Meers, J., Young, P.R., 2017. Phylogenetic Diversity of Koala
533 Retrovirus within a Wild Koala Population. *Journal of virology* 91.

534 Chong, Z., Ruan, J., Wu, C.I., 2012. Rainbow: an integrated tool for efficient clustering and
535 assembling RAD-seq reads. *Bioinformatics* 28, 2732-2737.

536 Dixon, P., 2003. VEGAN, A Package of R Functions for Community Ecology. *Journal of*
537 *Vegetation Science* 14, 927-930.

538 Fabijan, J., Woolford, L., Lathe, S., Simmons, G., Hemmatzadeh, F., Trott, D.J., Speight, N.,
539 2017. Lymphoma, Koala Retrovirus Infection and Reproductive Chlamydiosis in a
540 Koala (*Phascolarctos cinereus*). *Journal of Comparative Pathology* 157, 188-192.

541 Fiebig, U., Hartmann, M.G., Bannert, N., Kurth, R., Denner, J., 2006. Transspecies
542 transmission of the endogenous koala retrovirus. *Journal of virology* 80, 5651-5654.

543 Fu, L., Niu, B., Zhu, Z., Wu, S., Li, W., 2012. CD-HIT: accelerated for clustering the next-
544 generation sequencing data. *Bioinformatics* 28, 3150-3152.

545 Gonzalez-Astudillo, V., Allavena, R., McKinnon, A., Larkin, R., Henning, J., 2017. Decline
546 causes of Koalas in South East Queensland, Australia: a 17-year retrospective study of
547 mortality and morbidity. *Sci Rep* 7, 42587.

548 Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis
549 program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41, 95-98.

550 Hanger, J.J., Bromham, L.D., McKee, J.J., O'Brien, T.M., Robinson, W.F., 2000. The
551 nucleotide sequence of koala (*Phascolarctos cinereus*) retrovirus: a novel type C
552 endogenous virus related to Gibbon ape leukemia virus. *Journal of virology* 74, 4264-
553 4272.

554 Hobbs, M., King, A., Salinas, R., Chen, Z., Tsangaras, K., Greenwood, A.D., Johnson, R.N.,
555 Belov, K., Wilkins, M.R., Timms, P., 2017. Long-read genome sequence assembly
556 provides insight into ongoing retroviral invasion of the koala germline. *Sci Rep* 7,
557 15838.

558 Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogenetic trees.
559 *Bioinformatics* 17, 754-755.

560 Ihaka, R., Gentleman, R., 1996. R: A Language for Data Analysis and Graphics. *Journal of*
561 *Computational and Graphical Statistics* 5, 299-314.

562 Ishida, Y., Zhao, K., Greenwood, A.D., Roca, A.L., 2015. Proliferation of endogenous
563 retroviruses in the early stages of a host germ line invasion. *Molecular biology and*
564 *evolution* 32, 109-120.

565 Legione, A.R., Patterson, J.L., Whiteley, P., Firestone, S.M., Curnick, M., Bodley, K., Lynch,
566 M., Gilkerson, J.R., Sansom, F.M., Devlin, J.M., 2017. Koala retrovirus genotyping

567 analyses reveal a low prevalence of KoRV-A in Victorian koalas and an association
568 with clinical disease. *J Med Microbiol* 66, 236-244.

569 Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler
570 transform. *Bioinformatics* 25, 1754-1760.

571 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
572 Durbin, R., 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
573 25, 2078-2079.

574 McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis
575 and graphics of microbiome census data. *PLoS One* 8, e61217.

576 Miyazawa, T., Shojima, T., Yoshikawa, R., Ohata, T., 2011. Isolation of koala retroviruses
577 from koalas in Japan. *The Journal of veterinary medical science / the Japanese Society*
578 *of Veterinary Science* 73, 65-70.

579 Nyari, S., Waugh, C.A., Dong, J., Quigley, B.L., Hanger, J., Loader, J., Polkinghorne, A.,
580 Timms, P., 2017. Epidemiology of chlamydial infection and disease in a free-ranging
581 koala (*Phascolarctos cinereus*) population. *PLoS one* 12, e0190114.

582 Overbaugh, J., Donahue, P.R., Quackenbush, S.L., Hoover, E.A., Mullins, J.I., 1988.
583 Molecular cloning of a feline leukemia virus that induces fatal immunodeficiency
584 disease in cats. *Science* 239, 906-910.

585 Pantginis, J., Beaty, R.M., Levy, L.S., Lenz, J., 1997. The feline leukemia virus long terminal
586 repeat contains a potent genetic determinant of T-cell lymphomagenicity. *J Virol* 71,
587 9786-9791.

588 Puritz, J.B., Hollenbeck, C.M., Gold, J.R., 2014. dDocent: a RADseq, variant-calling pipeline
589 designed for population genomics of non-model organisms. *PeerJ* 2, e431.

590 Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2010. edgeR: a Bioconductor package for
591 differential expression analysis of digital gene expression data. *Bioinformatics (Oxford,*
592 *England)* 26, 139-140.

593 Shimode, S., Nakagawa, S., Yoshikawa, R., Shojima, T., Miyazawa, T., 2014. Heterogeneity
594 of koala retrovirus isolates. *FEBS Lett* 588, 41-46.

595 Shojima, T., Yoshikawa, R., Hoshino, S., Shimode, S., Nakagawa, S., Ohata, T., Nakaoka, R.,
596 Miyazawa, T., 2013. Identification of a novel subgroup of Koala retrovirus from Koalas
597 in Japanese zoos. *Journal of virology* 87, 9943-9948.

598 Simmons, G., Clarke, D., McKee, J., Young, P., Meers, J., 2014. Discovery of a novel
599 retrovirus sequence in an Australian native rodent (*Melomys burtoni*): a putative link
600 between gibbon ape leukemia virus and koala retrovirus. *PLoS One* 9, e106954.

601 Simmons, G.S., Young, P.R., Hanger, J.J., Jones, K., Clarke, D., McKee, J.J., Meers, J., 2012.
602 Prevalence of koala retrovirus in geographically diverse populations in Australia.
603 *Australian veterinary journal* 90, 404-409.

604 Tarlinton, R., Meers, J., Hanger, J., Young, P., 2005. Real-time reverse transcriptase PCR for
605 the endogenous koala retrovirus reveals an association between plasma viral load and
606 neoplastic disease in koalas. *The Journal of general virology* 86, 783-787.

607 Tarlinton, R., Meers, J., Young, P., 2008. Biology and evolution of the endogenous koala
608 retrovirus. *Cellular and molecular life sciences : CMLS* 65, 3413-3421.

609 Tarlinton, R.E., Sarker, N., Fabijan, J., Dottorini, T., Woolford, L., Meers, J., Simmons, G.,
610 Owen, H., Seddon, J., Hemmatzedah, F., Trott, D., Speight, N., Emes, R., 2017.
611 Differential and defective expression of Koala Retrovirus reveal complexity of host and
612 virus evolution. *bioRxiv*.

613 Theys, K., Libin, P., Pineda-Peña, A.-C., Nowé, A., Vandamme, A.-M., Abecasis, A.B., 2018.
614 The impact of HIV-1 within-host evolution on transmission dynamics. *Current Opinion*
615 *in Virology* 28, 92-101.

616 Waugh, C.A., Hanger, J., Loader, J., King, A., Hobbs, M., Johnson, R., Timms, P., 2017.
617 Infection with koala retrovirus subgroup B (KoRV-B), but not KoRV-A, is associated
618 with chlamydial disease in free-ranging koalas (*Phascolarctos cinereus*). *Sci Rep* 7, 134.
619 Xu, W., Gorman, K., Santiago, J.C., Kluska, K., Eiden, M.V., 2015. Genetic diversity of koala
620 retroviral envelopes. *Viruses* 7, 1258-1270.
621 Xu, W., Stadler, C.K., Gorman, K., Jensen, N., Kim, D., Zheng, H., Tang, S., Switzer, W.M.,
622 Pye, G.W., Eiden, M.V., 2013. An exogenous retrovirus isolated from koalas with
623 malignant neoplasias in a US zoo. *Proceedings of the National Academy of Sciences*
624 *of the United States of America* 110, 11547-11552.

625

626

627 **Figures:**

628 **Fig. 1:** Location of sample collection site. Red dot showing the sample collection cities. From
629 Queensland, samples were collected from Gold Coast, Brisbane, and Sunshine Coast and from South
630 Australia, koalas were collected from Mount Lofty region. Map was adapted from Australian Koala
631 Foundation (AKF) website.

632 **Fig. 2:** Phylogenetic tree from aligned KoRV sequences (including 169 newly identified in this study
633 and 24 previously published sequences with two sequences from GALV and MbRV (used as outgroups
634 to root the tree) generated through Geneious implemented Bayesian approach. Previously published
635 KoRV env sequences; KoRV-A (AF151794, KX587957.1 and KP792565.1), KoRV-B (KX588002.1,
636 KX588011.1, KX588027.1, KC779547.1, AB822553.1, KX588031.1, KX588053.1), KoRV-C
637 (AB828005.1, KP792564.1), KoRV-D (KX587952.1, KX587991.1, KX588043.1, KX587993.1,
638 KX587972.1, KX587972.1, AB828004.1, KX587997.1), KoRV-E (KU533853.1), KoRV-F
639 (KX588025.1, KX588028.1, KX587994.1, KU533852.1), KoRV-G (KX587961.1 and KX587998.1),
640 KoRV-H (KX588036.1 and KX587979.1), and KoRV-I (KX587976.1 and KX588021.1) were used.
641 Outgroup sequences: Gibbon ape leukaemia virus (GALV) sequences, (KT724047.1, KT724048.1) and
642 *Melomys burtoni* retrovirus (MbRV) sequences (KF572486.1, KF572485.1). Bayesian value are shown.
643 Paraphyletic KoRV-D has multiple subclades numbered D1 to D11 and also includes previously
644 designated KoRV-G and KoRV-H. Clades color and weight are marked as gradient following posterior
645 probabilities values

646 **Fig. 3:** Comparison of the read counts of KoRV env subtypes A, B, I and D (including subclades) in
647 the proviral DNA form within QLD (n = 33) and SA (n = 28) koala populations. Mean read counts with
648 one standard deviation error bars are shown. Although all SA animals had KoRV-B and D10, their
649 lower level read counts are not observable at this scale.

650 **Fig. 4:** Genetic diversity of KoRV env subtypes among paired DNA and RNA samples was illustrated
651 through the relative percentage of total reads of (A) QLD and (B) SA koala populations. Colors indicate

652 the different subtypes. (A) Among 33 QLD koalas, 28 were present in both DNA and RNA forms and
653 (B) among 28 SA koalas, 5 had both DNA and RNA forms.

654 **Fig. 5:** Percentage relative abundance of viral subtypes A (green), B (orange), I (red) and combined D
655 (purple). Compared between QLD and SA animals for A) DNA, B) RNA, C) RNA and DNA for paired
656 QLD samples, and D) RNA and DNA for paired SA samples. Median and interquartile ranges
657 displayed.

658

659

660 **Supplementary Files:**

661 **File 1 (figure):** Preliminary optimisation assemblies using *dDocent*. Two parameters were explored:
662 (1) the minimum number of samples required to represent a sequence in the assembly (colours, see
663 legend), and (2) the clustering thresholds used to group reads into similar sequences. The clear effect
664 of singletons (only requiring a sequence to be represented in one individual) on the number of contigs
665 probably arises from two competing (non-mutually exclusive) hypotheses: firstly, large variation exists
666 within and between individuals; and secondly, technical and sequencing error introduces sequence
667 variation. This problem appears to disappear when the number of samples was ≥ 2 .

668 **File 2:** Details of koala samples used for analysis of *env* gene diversity using 16S Metagenomics
669 sequencing.

670 **File 3 (figure):** Alignment of unique Koala Retrovirus (KoRV) *env* sequences from koalas sampled in
671 Queensland and South Australia.

672 **File 4 (figure):** Simple view of Bayesian phylogenetic tree. Unique sequences were aligned with
673 previously published KoRV *env* sequences; KoRV-A (AF151794, KX587957.1 and KP792565.1),
674 KoRV-B (KX588002.1, KX588011.1, KX588027.1, KC779547.1, AB822553.1, KX588031.1,
675 KX588053.1), KoRV-C (AB828005.1, KP792564.1), KoRV-D (KX587952.1, KX587991.1,
676 KX588043.1, KX587993.1, KX587972.1, KX587972.1, AB828004.1, KX587997.1), KoRV-E
677 (KU533853.1), KoRV-F (KX588025.1, KX588028.1, KX587994.1, KU533852.1), KoRV-G
678 (KX587961.1 and KX587998.1), KoRV-H (KX588036.1 and KX587979.1), and KoRV-I
679 (KX587976.1 and KX588021.1) using ClustalW alignment programme. Moreover, four other
680 sequences were used as outgroups to root the KoRV phylogeny: two Gibbon ape leukaemia virus
681 (GALV) sequences, (KT724047.1, KT724048.1) and two *Melomys burtoni* retrovirus (MbRV)
682 sequences (KF572486.1, KF572485.1). Alignments were further edited by hand to fill the blanks at the

683 beginning and end. Phylogenetic tree was determined from the aligned reads using the Geneious plugin
684 of MrBayes 3.2.6.

685 **File 5:** The read count of unique KoRV *env* sequences from PCR and next-generation
686 sequencing of individual koala DNA and RNA samples.

687 **File 6:** The relative percentage of each subtype of KoRV *env* gene in individual DNA and RNA
688 samples of koalas.

689 **File 7 (figure) 7:** Percentage abundance of each major subtype of KoRV (A, B, Combined D, I) in
690 the DNA of individuals compared with disease status (Healthy, Neoplasia, Oxalate Nephrosis,
691 Chlamydiosis). Median and interquartile range displayed.