



Water-based training enhances both physical capacities and body composition in healthy young adult women

Camilla Raffaelli¹ · Chiara Milanese¹ · Massimo Lanza¹ · Paola Zamparo¹

Received: 12 January 2016 / Accepted: 21 March 2016 / Published online: 8 April 2016
© Springer-Verlag Italia 2016

Abstract

Purpose The purpose of this study was to determine the effectiveness of a 9 weeks aquatic training program on aerobic capacity, muscle strength, flexibility, balance and body composition in 34 healthy young adult women.

Methods Five typical water based exercises (WE) of known intensity were utilized during the classes; intensity ranged from “moderate” to “hard” according to ACSM criteria (RPE range 12–14).

Results The group physical activity level and food intake were not significantly different before and after training. A significant decrease for the skin folds sum (−4.6 %) and for %fat mass (−3.8 %) calculated according to skin folds technique was found after training. DXA regional data showed a significant increase in the fat-free mass of arms (2.4 %) and trunk (0.9 %). According to the Astrand step test, the training program led to an increase of estimated maximal oxygen uptake (14.9 %) and a decrease in sub-maximal Heart Rate (−6.9 %). A significant change in the majority of the physical capacities tested was found: abdominal and upper body muscular endurance (21 and 36 %, respectively), leg flexors and extensors maximal strength (12 and 8 %, respectively) as well as balance (34 %) all improved after training ($p < 0.05$)

Conclusions A training program based on WE of known intensity and tailored to the ACSM recommendations can significantly improve cardio respiratory fitness, muscular

endurance, strength, balance and some aspects of body composition in active young adult women.

Keywords Water fitness · Aquatic exercise · Cardio respiratory fitness · Training effects

Introduction

The selection of an appropriate physical activity (in terms of exercise mode, intensity, frequency and duration) is essential to obtain actual training effects, to avoid injuries and to ensure exercise adherence. The American College of Sport Medicine guidelines [1, 2] place a strong emphasis on the different components of physical fitness and underline how, given the specific nature of exercise adaptation, a well rounded training program should consist of aerobic, resistance and flexibility exercises.

Water-based activities (WA), in recent years, have gained popularity and are considered as one of the possible alternatives among the traditional physical activities for well-being and health [3–14]. The specific properties of water have probably encouraged this success. In particular: buoyancy reduces the effect of weight bearing on skeletal joints while the larger density of water (compared to air) provides loading during all movements. In addition, the hydrostatic pressure and the water temperature improve blood flow and favourably alter the hemodynamic responses at rest and during exercise [15–19]. Thanks to these features many people chose to perform physical activity in water rather than on land, especially people with low levels of physical fitness. These activities are often prescribed for elderly people [7, 9–11, 20–40], post-menopausal women [41, 42], subjects with orthopaedic or neurological disabilities [43–53], subjects with pulmonary

✉ Paola Zamparo
paola.zamparo@univr.it

¹ Department of Neurological, Biomedical and Movement Sciences, University of Verona, via F. Casorati 43, 37131 Verona, Italy

disease [54–57], patients with coronary artery disease [58–60], athletes with surgically, or otherwise, treated injuries [61, 62] and obese persons [63–68]. WA are widely used also to improve the physical condition of young adult healthy individuals who regularly take part in recreational training [3, 69–71] and by athletes, as complement of training [72, 73].

The physiological and training effects of WA have indeed been investigated by several authors in different populations (especially in older adults, elderly, obese person and in subject with orthopaedic and neurological disease): some studies have examined mainly cardio respiratory factors, body composition and lipoproteins-lipids patterns [3–5, 9–11, 14, 20, 22, 28, 32, 36, 37, 42, 63, 64, 66, 68], whereas others studied mainly the resistance component of static and dynamic strength [3, 7, 25, 27, 28, 30, 32, 33, 36, 41, 43, 69, 72, 74–76] and only few studies considered other physical capacities such as flexibility [7, 10, 27, 31, 46], agility and balance [7, 28, 31, 38]. In some cases cognitive function, activity of daily living, quality of life and adherence to exercise in the aquatic environment were also considered [22, 24, 31, 32, 34, 35, 43, 49, 54, 56, 64].

However, the quantification and the control of work intensity in water remain still difficult, particularly so for aquagym, aquaerobics or callisthenics exercises (generally utilized in water fitness programs) as compared to walk, run or aqua bike, which are more easy to standardize.

Two methods to monitor exercise intensity in water are available. Colado et al. [77] showed that the exercise intensity during aquatic resistance movements can be controlled by setting the rhythm of execution and by checking the perceived effort. Raffaelli et al. [78] showed that it is possible to standardize the most common water based exercises (WE) in terms of amplitude and frequency of movement and that the intensity of exercise could be controlled by changing the type of exercise and/or the frequency of the music track. Thus, the intensity level of an aquatic fitness lesson can be predicted and the intensity of a training program could be determined and set to correspond to ACSM criteria [1, 2, 79] to maintain and improve cardiovascular fitness in a given population (e.g. young adult healthy women).

The purpose of the present study was to determine the effectiveness of a 9-week aquatic training program on aerobic capacity and body composition, in healthy young adult women. Training duration was determined based on ACSM indications (9 weeks is the minimum duration to obtain these adaptations in these subjects) [1, 2]. The intensity of the WE utilized during training ranged from moderate to hard, according to ACSM criteria [1, 2, 79]. A complementary analysis of other health related physical capacities (muscle strength, flexibility and balance) was

also carried out to better understand whether these WA can be considered a well rounded training program as well.

Materials and methods

Subjects

Thirty-four physically active women (26.4 ± 3.8 years; 1.63 ± 0.05 m of stature; 56.9 ± 8.3 kg of body mass; 19.7 ± 1.5 kg m⁻² of BMI) were recruited from the Faculty of Human Movement Sciences of the University of Verona and voluntarily participated to the study (WA are mainly performed by females). The inclusion criteria were an age range between 20 and 30 years, a stature range between 1.6 and 1.7 m and a body mass index (BMI) lower than 25 kg m⁻². The stature was selected as an inclusion criteria to standardize, as much as possible, the water level for all subjects.

The participants were instructed not to modify their physical activity level and diet for the duration of the study; all completed the study.

All participants received written and oral instructions before the study and gave their informed consent to the experimental procedure. The experimental protocol was approved by the Institutional Review Board.

Procedures

The International Physical Activity Questionnaire (IPAQ) was compiled before and after training to monitor the physical activity level of the participants; an alimentary diary was also compiled to monitor the changes in food intake before and after training.

Body composition and regional fat distribution was investigated based on measures of body stature, body mass, skin folds and selected anatomical circumferences as well as by means of total body of dual-energy X-ray absorptiometry (DXA). These measurements will be described in detail below.

Physical capacities were assessed by means of the following tests: astrand step test for aerobic capacity; sit-up and push-up tests for muscular endurance; leg-curl, leg-extension and pectoral machine tests for maximal strength; flamingo balance test for balance; sit and reach test and back stretch test for flexibility.

The participants were familiarized with each test and their performance was continuously checked, and eventually corrected, by two experienced trainers. The tests were performed at the same time of day, in the same environment and in the same order during the pre-training and post-training session with 20 min recovery in between. During each test, the subjects were verbally encouraged to

perform maximally. These tests will be described in detail below.

Each subject performed the tests 1 week before training and 1 week after training. Training was carried out twice a week in a swimming pool (water depth: 1.2 m; water temperature: 28 °C) for 9 weeks. At the end of each training session the subjects were asked to report their rate of perceived exertion by means of the Borg Scale [80].

Anthropometric measurements

Body height to the nearest 1 mm was measured using a wall-mounted stadiometer (Holtain Limited, UK), body mass was measured to the nearest 0.1 kg using an electronic scale (Tanita electronic scale BWB-800 MA, Wunder SA.BI. Srl) and the body mass index was calculated from these data ($BMI = \text{body mass (kg)}/\text{height (m}^2\text{)}$).

The selected anatomical circumferences (arm, wrist, waist, hip, calf and thigh) were measured to the nearest 0.1 cm using anthropometry tapes. In total, eight skin folds (triceps, sub-scapular, chest, abdominal, front thigh, medial calf, supra iliac and mid-axillary) were measured using a skinfold caliper (type Harpenden—GIMA), on the right side of the body, while the participant was standing with arms by her sides. All measurements were carried out by the same trained examiner under identical environmental conditions. The average of two trials was used for further analysis and, when the measurement differed by more than 1.0 mm, a third measurement was obtained and the mean value was utilized.

Total body and regional composition (lean mass, fat mass, and mineral mass) was evaluated by means of dual-energy X-ray absorptiometry (DXA) using a total body scanner (QDR Explorer W, Hologic, MA, USA; fan beam technology, software for Windows XP version 12.6.1) according to the manufacture's procedures. The percentage of fat-free mass from DXA was calculated as the sum of lean tissue mass and bone mineral content; the percentage of body fat was calculated as the ratio of fat tissue to body weight.

In addition, percent fat mass was assessed based on measurements of skin folds: the sum of seven skin folds (triceps, sub-scapular, chest, abdominal, front thigh, supra iliac and mid-axillary) was used to assess body density according to Jackson and Pollock [81], while percent fat mass was calculated on the basis of body density according to Siri [82].

Physical capacities tests

The Step Test was used to estimate aerobic capacity. The subjects were asked to step up and down a bench (bench height: 33 cm) with a cadence of 22.5 beats per minute for

3 min: during this period the heart rate (HR) was recorded. Oxygen consumption was estimated based on the ACSM stepping equations [79]. Finally, maximal oxygen consumption ($\dot{V}O_{2\max}$) was calculated on the basis of the Astrand-Rhyming Nomogram equation [83].

The sit and reach test was used to test trunk flexion (the flexibility of the lower body) and was performed according to ACSM instructions [79]. The distance between the fingers and the vertical surface of the box at full stretch was measured (if the subject cannot reach the toes, a negative score was recorded) and the best of three trials (distance, cm) was recorded.

The Back Stretch test was used to assess upper-body (shoulder) flexibility and was performed according to Morrow et al. [84]. The test's score was the distance (cm) of overlap or the distance between the tips of the middle fingers. The best of three trials was recorded.

The Sit-Up test was used to assess abdominal endurance and the Push-Up test was used to assess upper-body endurance. Both tests were performed according to Morrow et al. [84]: the test's score was the total number of correct repetitions.

The exercises chosen to assess maximum dynamic strength were: leg-extension, leg-flexion and adduction of arms (leg-curl, leg-extension and pectoral machine—Personal Selection TUV, Technogym, Italy); the method utilized was the One Repetition Maximum (1-RM): the subjects started from a workload that they could move up to 5/6 times; the workload was gradually increased to the load that could be moved one-time only: the test's score was this final load.

The Flamingo Balance test was used to assess balance capacity. In our case the test was modified as proposed by Zurc et al. [85]. The test's score was the time passed from the start to the moment the bent leg touched the floor; maximal duration of the test was set to 1 min.

Training Program (in the swimming pool)

Each subject underwent a training program two times a week for 9 weeks. Each lesson lasted for 45 min: the program consisted of a 10 min of warm-up, 30 min of workout and 5 min of cool-down; each session was led by trained fitness instructors. The frequency of training (twice a week) was the most convenient for our participants and was also chosen to avoid dropouts.

Five typical water based exercises (WE), commonly utilized during water fitness activities (without specific equipment, such as aquagym and aquaerobics), were utilized during the classes [78]:

1. "Running on the spot raising the knees high" (S)
2. "Jumping on the spot moving the legs sideways (in the frontal plane)" (SJ)

3. “Jumping on the spot moving the legs backward and forward (in the sagittal plane)” (FJ)
4. “Alternate forward kicks (in the sagittal plane)” (FK)
5. “Alternate sideways kicks (in the frontal plane)” (SK).

These exercises were performed at different frequencies (corresponding to different music tracks): $F_1 = 110\text{--}120$ bpm; $F_2 = 120\text{--}130$ bpm; $F_3 = 130\text{--}140$ bpm (corresponding to 1.8–2.0, 2.0–2.17 and 2.17–2.33 Hz, respectively). Every leg movement (e.g. raising the leg or lowering the leg, in the skip) corresponded to a sound beat. These frequencies were selected because the most frequently utilized in the music tracks that accompany WA. Participants were verbally encouraged to maintain the correct movement amplitude and frequency.

During training, the warm up and cool down phases were carried out asking the subjects to perform a specific combination of S, SJ and FJ at F_1 and SJ and FJ at F_2 . As indicated by Raffaelli et al. [78] this combination of exercises and frequencies correspond, for subjects of this age and fitness level, to a “light” intensity level (20–39 % of $\dot{V}O_{2\max}$) according to ACSM classification [2]. The 30 min of workout were carried out using mainly FK and SK. These exercises can indeed be considered as “moderate” (40–59 of $\dot{V}O_{2\max}$) when performed at F_1 and F_2 and as “hard” (60–84 of $\dot{V}O_{2\max}$) at F_3 . In the workout phase we also used S, SJ and FJ exercises but at higher frequencies (F_2 and F_3) than during the warm up and cool down phases since, at these frequencies their intensity becomes “moderate” (40–59 of $\dot{V}O_{2\max}$) as indicated by Raffaelli et al. [78].

As indicated by Table 1, were the training schedule is schematically represented, the warm up and cool down phases were always the same. On the contrary the workout was progressively increased: during the first 2 weeks the intensity was always “moderate”, from the third to the fifth week the intensity ranged from “moderate” to “hard” while during the last 4 weeks the intensity was always “hard”. As indicated by Table 1, during training the participants worked at about 20–39 % of their $\dot{V}O_{2\max}$ during warm up and cool down and at 40–84 % of $\dot{V}O_{2\max}$ during the workout. This level of intensity should, according to ACSM guidelines [2], maintain and improve cardiovascular fitness in subjects of this age and physical capacity level.

Statistical analysis

Data are presented as mean ± 1 standard deviation (SD). Percent changes (before to after) were calculated as well. Normality of distribution was assessed for each variable with the Shapiro–Wilk test. Normally distributed variables

were compared (pre-post training) by means of a parametric test (two-tailed paired Student’s test), otherwise a non-parametric test (Wilcoxon Signed Rank Test) was applied; the level of significance was set at $p < 0.05$. Statistical analysis was performed by using SigmaStat 3.5.

Results

The group attendance to training was about 93 %. The study was completed without any case of injury. The group’s physical activity level (see Fig. 1) and food intake were not significantly different ($p < 0.05$) before and after training.

The IPAQ questionnaire results showed that 83 % of participants in the study was moderately active (from 700 to 2500 METs min week⁻¹) or active (over 2500 METs min week⁻¹) whereas 17 % was not sufficiently active (less than 700 METs min week⁻¹). Also these subjects, however, performed moderate physical activity at least twice a week for at least 30 min.

Anthropometric measurements

A significant decrease was found for the majority of the selected skin folds and circumferences (see Table 2) after training: even when the changes were not significant, the trend was towards a decrease (except for hip circumference). Indeed, a significant decrease in skin fold sum and % fat mass (calculated according to these measurements) was found after training (see Table 3).

The total body DXA data indicated no significant changes after training even if the trend was for a decrease, in agreement with what found for the anthropometric measures (see Table 3). On the other hand, the DXA regional data showed a significant improvement in the fat-free mass of arms and trunk (see Table 3).

Physical capacities tests

RPE was measured every training session in all subjects: the average values (± 1 SD) are reported in Fig. 2 which shows that perceived exertion was in the 12–14 range for the entire duration of training (from moderate to hard) in line with ACSM recommendations.

Submaximal HR during the step test and estimated maximal oxygen consumption changed significantly after training: HR decreased from 146.7 ± 15.3 to 136.5 ± 12 bpm ($\Delta \% = -6.9$, $p < 0.001$) and $\dot{V}O_{2\max}$ improved from 47.3 ± 9.1 to 54.3 ± 8.8 mL min⁻¹ kg⁻¹ ($\Delta \% = 14.9$, $p < 0.001$) from the first to the ninth week of training (see Fig. 3).

Table 1 A general overview of the training program

Week	Training program			Workout (30 min)		
	Exercise type and movement frequency	Relative intensity (% $\dot{V}O_{2max}$)	Target ACSM	Exercise type and movement frequency	Relative intensity (% $\dot{V}O_{2max}$)	Target ACSM
I	SJ,F1-FJ,F1	31	Light	SJ,F3-FJ,F3	43	Moderate
	S,F1-SJ,F2-FJ,F2	37		S,F2	45	
II	SJ,F1-FJ,F1	31	Light	FK,F1-SK,F1	47	Moderate
	S,F1- SJ,F2-FJ,F2	37		S,F3	54	
III	SJ,F1-FJ,F1	31	Light	FK,F2-SK,F2	56	Moderate
	S,F1-SJ,F2- FJ,F2	37		SJ,F3-FJ,F3	43	
IV	SJ,F1-FJ,F1	31	Light	S,F2	45	Hard
	S,F1-SJ,F2- FJ,F2	37		FK,F1-SK,F1	47	
V	SJ,F1- FJ,F1	31	Light	S,F3	54	Moderate
	S,F1-SJ,F2- FJ,F2	37		FK,F2-SK,F2	56	
VI	SJ,F1- FJ,F1	31	Light	FK,F3-SK,F3	63	Hard
	S,F1- SJ,F2 -FJ,F2	37		S,F3	54	
VII	SJ,F1-FJ,F1	31	Light	FK,F2-SK,F2	56	Hard
	S,F1-SJ,F2-FJ,F2	37		FK,F3-SK,F3	63	
VIII	SJ,F1-FJ,F1	31	Light	S,F3	54	Hard
	S,F1-SJ,F2- FJ,F2	37		FK,F3-SK,F3	63	
IX	SJ,F1-FJ,F1	31	Light	FK,F3-SK,F3	63	Hard
	S,F1-SJ,F2- FJ,F2	37		FK,F3-SK,F3	63	

S: “running on the spot raising the knees high”; SJ: “jumping on the spot moving the legs sideways (in the frontal plane)”; FJ: “jumping on the spot moving the legs backward and forward (in the sagittal plane)”; FK: “alternate forward kicks (in the sagittal plane)”; SK: “alternate sideways kicks (in the frontal plane)”; $F_1 = 110\text{--}120$ bpm; $F_2 = 120\text{--}130$ bpm; $F_3 = 130\text{--}140$ bpm

Before and after training there was also a significant change in the majority of the physical capacities tested: muscular endurance, maximal strength and balance all improved after the training program. On the contrary, flexibility wasn't improved after the training program (see Table 4).

Discussion

The purpose of the present study was to determine the effectiveness of a 9 weeks aquatic training program on aerobic capacity, muscle strength, flexibility, balance and

body composition in healthy young adult women with the aim to understand if WA, standardized in terms of exercises intensity, can be considered a well rounded training program. To this purpose we choose to utilize field tests that can be easily reproduced by water fitness teachers.

The high adherence to the study suggests that aquatic exercise training is well tolerated by these subjects; this type of physical activity seems to facilitate adherence to training and to limit dropouts. Indeed our RPE data (range 12–14) are in line with the target value associated with high adherence to training as indicated by ACSM [2]. High adherence to a water-based training program was found




Fig. 1 Physical Activity Levels (MET min wk⁻¹) before (pre) and after (post) training; bars represent 1 SD

also by Mc Namara et al. [54] in people with chronic obstructive pulmonary disease. These authors have shown that six factors were highly rated to exercise adherence: staff support, enjoyment, sense of achievement, noticeable improvements, personal motivation and participant support.

The physical activity level of the subjects was over the range of 450–750 METs min week⁻¹ which is indicated as the minimal value for this population [1]; according to the replies received in the questionnaires, the participants' lifestyle patterns (which include physical activity levels and dietary intakes) did not seem to have changed throughout the training period. It is therefore fair to assume that the pre to post differences can indeed be attributed to training itself. This is in line with previous studies showing that, in healthy young active women, physical capacity and body composition do not change in such a short period of

time (9 weeks) unless there are important changes in food intake and/or physical activity level [65, 86].

Anthropometric characteristics

In this study no significant changes in body mass were recorded (with both measurement techniques). This result is in agreement with Rica et al. [64], Takeshima et al. [10], Bocalini et al. [36] and Broman et al. [37] who did not find changes in body mass after a water-based training program. On the contrary, Volaklis et al. [87] and Gappmaier et al. [68] did find significant decreases in body mass (1.7–2.0 and about 6 %, respectively) as well as Greene et al. [66]. The regional DXA data, however, showed a significant increase of fat-free mass for trunk and arms in line with other studies that found a significant increase in fat free mass after water resistance training [20, 66, 69, 74].

Our training program revealed adaptations on body composition as assessed with anthropometric measures (skin folds and circumferences) and DXA regional data. Anthropometric techniques do not estimate body composition per se but are an excellent indicator of subcutaneous adiposity. For these parameters (skin folds and circumferences, see Table 2) our data are in line with the results of Takeshima et al. [10], Volaklis et al. [87] and Gappmaier et al. [68] that found a significant decrease in skin fold sum after aquatic training.

While % fat mass assessed by skin folds was decreased after training in agreement with Gappmaier et al. [68],

Table 2 Effects of training on the anthropometric characteristics of the subjects

	Pre	Post	% Change	<i>p</i>	95 % IC
Skin folds (mm)					
Triceps	17.6 ± 5	15.9 ± 4.8	-9.2 ± 11.9	0.000	0.863 to 2.381
Sub-scapular	10.4 ± 4.3	10.1 ± 4.2	-2.5 ± 7.7	0.108	-
Chest	8.8 ± 3.6	8.4 ± 3.2	-4.6 ± 18.8	0.175	-
Abdominal	18.1 ± 7.9	17.1 ± 6.7	-5.9 ± 12.5	0.048	n.p.
Front thigh	26.4 ± 6.6	25.2 ± 6.5	-4.5 ± 9.5	0.011	0.292 to 2.096
Medial calf	15.3 ± 3.8	14.7 ± 3.7	-3.4 ± 8	0.040	n.p.
Supra-iliac	14.5 ± 6.6	14.3 ± 6.4	-1.2 ± 10.5	0.517	-
Mid-axillary	8.5 ± 4.2	8.4 ± 4.2	-1.3 ± 10.6	0.475	-
Circumferences (cm)					
Arm	25.6 ± 2.7	25.1 ± 2.4	-1.7 ± 3.3	0.014	n.p.
Wrist	14.9 ± 0.6	14.8 ± 0.6	-0.4 ± 1.7	0.241	-
Waist	68.3 ± 6.1	67.8 ± 6	-0.8 ± 1.6	0.009	0.150 to 0.956
Hip	93.7 ± 17.9	96.3 ± 5.5	2.7 ± 20.2	0.696	-
Calf	34.4 ± 2.2	34.3 ± 1.9	-0.4 ± 2.5	0.344	-
Thigh	50.6 ± 3.5	49.6 ± 3.1	-1.9 ± 2.6	0.000	0.521 to 1.467

Data are mean ± 1 SD. In the last column are reported the 95 % IC of the paired Student's *t* test (for *p* < 0.05) and it is indicated whether a Wilcoxon Signed Rank Test was utilized instead (non-parametric test: n.p.)

Table 3 Effects of training on body composition

	Pre	Post	% Change	<i>p</i>	95 % IC
Skin fold technique					
Sum of skin folds (mm)	104.3 ± 34.7	99.5 ± 32.1	-4.6 ± 7.7	0.002	0.945 to 7.736
Fat mass (%)	19.5 ± 5.2	18.7 ± 4.8	-3.8 ± 6.5	0.002	0.283 to 1.198
Body mass (kg)	56.9 ± 8.3	56.6 ± 8.1	-0.5 ± 2	0.176	-
DXA					
Total mass (kg)	56.4 ± 8.2	56.2 ± 8	-0.2 ± 0.5	0.903	-
Total fat mass (kg)	16.3 ± 5.6	16 ± 5.5	-1.6 ± 6.9	0.495	-
Total fat-free mass (kg)	38 ± 3.9	38.1 ± 3.8	0.3 ± 2.2	0.384	-
Fat mass (%)	28.4 ± 5.5	28.0 ± 5.5	-1.4 ± 5.7	0.166	-
Trunk fat-free mass (kg)	19.2 ± 2	19.4 ± 2	0.9 ± 2	0.037	-33.04 to -10.92
Arms fat-free mass (kg)	1.74 ± 0.2	1.79 ± 2	2.4 ± 0.5	0.005	-70.39 to -13.64
Legs fat-free mass (kg)	6.3 ± 0.7	6.2 ± 0.7	-1 ± 2.2	0.138	-

Data are mean ± 1 SD. In the last column are reported the 95 % IC of the paired Student's *t* test (for *p* < 0.05)

DXA: total body densitometry (DXA Hologic QDR Explorer W, Fan Beam technology)




Fig. 2 RPE average values (bars represent 1 SD) during the eighteen training sessions (9 weeks, two sessions per week)




Fig. 3 Percentage changes (pre-post training) of sub-maximal Heart Rate (*black column*) and estimated Maximal Oxygen Consumption (*white column*) as assessed by means of the Astrand step test; *bars* represent 1 SD

Colado et al. [69], Greene et al. [66], Bergamini et al. [28], Kasprzak et al. [63] and Irandoust et al. [20], % fat mass assessed by DXA showed only a decreasing trend. Moreover, the values of % fat mass assessed by means of skin folds measurements were found to be substantially lower than those assessed by means of DXA.

This difference was previously reported by Peterson et al. [88], Silva et al. [89] and España-Romero et al. [90] who examined the accuracy of the existing equations to estimate percent body fat (such as the Jackson and Pollock's equation utilized in this study); these authors have indeed shown that % fat mass assessed by means of anthropometric measurements is underestimated when compared to the three or four-compartment model, as found in our study. In general terms, the studies that compared body composition measurements using DXA,

Table 4 Effects of training on physical capacities

Test	Pre	Post	% Change	<i>p</i>	95 % IC
Muscular endurance					
Sit-up (N of repetitions)	37.6 ± 26.6	45.4 ± 29.9	20.8 ± 44.6	0.004	n.p.
Push-up (N of repetitions)	23.5 ± 13.1	31.9 ± 15.4	35.9 ± 36	0.000	−11.39 to −5.49
Maximal strength					
Leg-curl (1 RM)	32.6 ± 7.5	36.5 ± 7.6	11.9 ± 15.9	0.000	n.p.
Leg-extension (1 RM)	43.8 ± 14.9	47.5 ± 16.9	8.5 ± 25.2	0.036	n.p.
Pectoral machine (1RM)	15.4 ± 4.3	17.4 ± 6.2	12.9 ± 33.8	0.051	n.p.
Balance					
Flamingo balance (s)	28.4 ± 22.7	42.8 ± 22.7	33.9 ± 91	0.049	n.p.
Flexibility					
Sit and reach (cm)	6.5 ± 7.6	5.9 ± 9.4	−8.6 ± 104.9	0.613	–
Back stretch (cm)	8 ± 5.9	7.7 ± 4.2	−4.36 ± 67.2	0.318	–

Data are mean ± 1 SD. In the last column are reported the 95 % IC of the paired Student's *t* test (for $p < 0.05$) and it is indicated whether a Wilcoxon Signed Rank Test was utilized instead (non-parametric test: n.p.)

BIA (Bioelectrical Impedance) and skinfold measurement techniques in children and in adults [91–93], have consistently shown that body composition estimates are highly dependent on the method utilized: these results therefore underline the importance of using the same method when comparing data before and after training.

Our data suggest that 9 weeks of aquatic training are not sufficient to modify total body mass in our population. Despite this, small changes in body composition (especially for subcutaneous adiposity and for the lean tissue of trunk and arms) and a decreasing trend for all investigated body composition parameters (as assessed by both methods) were observed after training.

Physical capacities test

The duration of our training program was of 9 weeks. The ACSM guidelines [2] indicate that a 15–20 weeks training may be an adequate minimum standard to evaluate the efficacy of exercises of various intensities, frequencies, and durations on fitness variables. However, these guidelines also indicate that to evaluate the time course of adaptations to training, shorter training programs such as the one of our study, may suffice.

The physical capacity tests we utilized are field tests that can be easily reproduced by any water fitness teacher to evaluate the training effects of their program.

Aerobic capacity

The majority of the data published in the literature about water fitness training (from 8 to 12 weeks) show increases

in $\dot{V}O_{2\max}$ similar to the ones reported in this study (about 15 %). Our results are in line with data reported by Pinto et al. [3], Greene et al. [66], Bocalini et al. [32], Ruoti et al. [40], Tauton et al. [94], Takeshima et al. [10] and Broman et al. [37] who found an improvement in $\dot{V}O_{2\max}$ of 10–15 %. In comparison with our study, Chu et al. [60] and Bocalini et al. [36] found larger (22 and 42 % respectively) and Cider et al. [95] lower (6 %) gains in $\dot{V}O_{2\max}$ after water fitness training. Other studies [55, 57, 64] investigated the changes in “aerobic capacity” after water-fitness training finding improvements in this parameter; a direct comparison with our results is, however, difficult because, in those papers, the change in aerobic capacity (e.g. $\dot{V}O_{2\max}$) was not directly measured, but estimated based on the outcome of a 6-min walking test.

As far as data of HR are concerned, the majority of the studies reported in the literature analyse changes in resting heart rate, rather than in sub maximal heart rate but for Broman et al. [37] who found a decrease in sub maximal heart rate (3 %) after training comparable to the one reported in our study (6.7 %).

It is worth noting that all the above cited papers refer to elderly subjects [10, 32, 36, 37, 40, 60, 64, 94], to people with chronic obstructive pulmonary disease [55–57], to obese older women [64] or to physically inactive overweight and obese adults [66], and/or to subjects with chronic heart failure [95] or chronic stroke [60]. Only Pinto et al. [3] investigated the change in aerobic capacity in young women. Our data therefore extend this body of knowledge to young adult female subjects and indicate that this kind of activity is suitable to improve cardiovascular fitness in physically active subjects too.

Muscular endurance and strength

The muscular endurance and strength test utilized in this study to assess the effects of training were similar to the movements that were practiced during the training itself and the effectiveness of these tests to assess functional mobility has been well established [79, 96]. Moreover, these tests were selected because suitable for different populations, because they can be performed outside the laboratory without expensive and sophisticated equipment and because they allow to assess (although indirectly) health related physical capacity.

Generally, when the training's aim is to improve strength, aquatic resistance devices or elastic bands are used to modulate drag force [7, 10, 69, 74, 76]. In our study we utilized the frequency of movement as a "resistance device" (hydrodynamic resistance indeed increases with the square of the speed of movement) and hence our training was standardized both in terms of exercise intensity ($\%V'O_{2max}$) and in terms of hydrodynamic resistance.

On this line of reasoning it seems possible to compare data reported in the literature about water resistance training with the results obtained in this study. Our results indicated that 9 weeks of aquatic training increased abdominal and upper body muscular endurance by 21 and 36 %, respectively. These improvements are in line with the results reported by Ruoti et al. [40] whereas Colado et al. [74] found higher improvements of abdominal and upper body muscle endurance (28 and 50 %, respectively); the training period of that study, however, was about three times longer than ours (24 vs. 9 weeks). On the contrary, Tauton et al. [94] did not found any significant change before and after training for muscular endurance data.

In our study, maximum dynamic strength was assessed by means of the method of One Repetition Maximum (1-RM) and improved by 12 % for leg flexors ($p = 0.000$), 8 % for leg extensors ($p = 0.036$) and 13 % for pectoral muscles ($p = 0.051$) after training. These results are in line with Tsourlou et al. [7], who found an improvement of dynamic strength in leg-extension (29.4 %), leg press (29.55 %) and chest press (25.7 %) and with Pinto et al. who found an improvement in lower body 1RM both in postmenopausal women [41] and in young women [71]. Our results are also comparable with data reported by Colado et al. [69] who found significant improvements in maximal strength of the upper limbs assessed by vertical row, horizontal bench press, horizontal bench row, arm lateral raise (3.7 %, on the average), with data reported by Volaklis et al. [87] who found an improvement of total strength assessed by bench press, pull down, seated row, leg extension and hamstring curl (12.8 %, on the average) and with data reported by Oh et al. [27] who found a significant improvement for strength of hip abduction and

adduction. In line with these results, Bento and Rodacki [25] and Rica et al. [64] reported an improvement in maximal dynamic strength while Bergamini et al. [28] reported that water-based training was beneficial only in maintaining knee-extension strength. Finally, Graef et al. [33] showed that a significant increase in maximal 1RM can be obtained only when the exercise load can be accurately determined and administered.

In the literature, maximum strength is also evaluated by means of different methods, e.g. by measuring the peak isometric torque; peak torque of knee extension and flexion was shown to improve significantly after aquatic training [7, 10, 25, 30, 41, 60, 71, 76]. These data are therefore in agreement with our findings and indicate that aquatic training produces significant effects on muscular endurance and strength.

It is worth remarking that most data published in the literature about the effects of water exercise on endurance and muscle strength refer to older subjects [7, 10, 25, 27, 28, 30, 33, 40, 64] or postmenopausal women [41, 74], to subjects with chronic heart failure [87] or chronic stroke [60] and that only few papers [3, 69, 71, 76] refer to fit young subjects. Our data therefore extend this body of knowledge to young adult female subjects and indicate that this kind of activity is suitable to improve neuromuscular performance in physically active subjects too.

Flexibility

Data reported in this study did not show any significant improvement in lower and upper body flexibility in line with the results of Oh et al. [27] and of Taunton et al. [94]. Other studies have shown a significant improvements of this physical capacity after aquatic exercise training: Sanders et al. [26] found a significant improvement in flexibility (8 %) as Sato et al. [31] who presented similar data. Takeshima et al. [10] found a significant change in trunk extension (11 %) whereas Bergamini et al. [28], Tsourlou et al. [7], Bocalini et al. [36] and Colado et al. [74] found an improvement of lower body flexibility (from 11 to 44 %) and upper body flexibility after training. In other papers [e.g. 46, 47] statistically significant differences after water training are reported but, in these cases, the range of motion and not flexibility is considered. The lack of differences observed in our study could be attributed to the fact that, unlike the above cited studies, we did not use specific flexibility exercises during training.

Balance

Our results showed significant improvements in balance (33 %, $p = 0.049$). It is quite difficult to compare this finding with those from other studies [e.g. 7, 21, 26, 28, 36,

38, 60] where functional mobility, agility and fear of falling were assessed instead of balance (balance is, however, a strong component of these parameters) but our findings are in line with those of Sanders et al. [26], Devereux et al. [38], Fiskén et al. [43], Tsourlou et al. [7] and Bocalini et al. [36]. Bento et al. [21] and Bergamini et al. [28] founded improvements after training only for dynamic balance. On the contrary, in the paper of Chu et al. [60], no improvement was reported in this parameter; in their study balance was assessed by the 14-item Berg Balance Scale (that measures balance while a person is sitting, standing or stepping). However, the balance assessment is highly dependent on the method utilized so, it seems difficult to compare data obtained from different tests.

Limitations of this study

This is a pre-test, post-test single group design; no control group was recruited. As discussed in the literature (e.g. Micallef [97]) there are some advantages in using a single group instead of a double group design since non-active participants (control group) could lose interest and drop out or develop a competitive attitude against the intervention group (“compensatory rivalry”). Moreover, a within subjects analysis has the advantage that subjects act as their own controls. At this regard, we carefully checked that no changes in physical activity of daily living (physical activity level) and in food intake occurred during training. These parameters have been measured/estimated by means of the IPAQ questionnaire and the alimentary diary and, since no pre to post differences were observed in these parameters the effects of training could indeed be attributed to training itself.

Conclusions

Our results indicate that a training program based on water exercises of known intensity and tailored to the ACSM recommendations, can significantly improve cardio respiratory fitness, muscular endurance and strength as well as balance and some aspects of body composition in active young adult women. In addition, the high adherence to our protocol seems to indicate that this type of physical activity is well tolerated by this population of subjects even if the specific properties of medium make it particularly suitable for elderly people and person with orthopaedic and neurological disease. On the basis of our findings, WA of known intensity (from moderate to hard) can indeed be considered a well rounded training program to improve health related aspects of physical fitness in healthy subjects as well.

Acknowledgments No funding was received for this work.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

- Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA et al (2007) Physical Activity and Public Health: update recommendation for adults from the American College of Sport Medicine and the American Heart Association. *Med Sci Sports Exerc* 39:1423–1434
- American College of Sport Medicine (1998) The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc* 30:975–991
- Pinto SS, Alberton CL, Cadore EL, Zaffari P, Baroni BM, Lanferdini FJ, Radaelli R, Pantoja PD, Peyrè-Tartaruga LA, Wolf Schoenell MC, Vaz MA, Krueger LF (2015) Water-based concurrent training improves peak oxygen uptake, rate of force development, jump height and neuromuscular economy in young women. *J Strength Cond Res* 29(7):1846–1854. doi:10.1519/JSC.0000000000000820
- Nagle EF, Sanders ME, Shafer A, Barone Gibbs B, Nagle JA, Deldin AR, Franklin BA, Robertson RJ (2013) Energy expenditure, cardiorespiratory, and perceptual responses to shallow-water aquatic exercise in young adult women. *Phys Sportsmed* 41(3):67–76. doi:10.3810/psm.2013.09.2018
- Guimaraes GV, Cruz LG, Tavares AC, Dorea EL, Fernandes-Silva MM, Bocchi EA (2013) Effects of short-term heated water-based exercise training on systemic blood pressure in patients with resistant hypertension: a pilot study. *Blood Press Monit* 18(6):342–345. doi:10.1097/MBP.0000000000000000
- Meredith-Jones K, Waters D, Legge M, Jones L (2011) Upright water-based exercise to improve cardiovascular and metabolic health: a qualitative review. *Complement Ther Med* 19(2):93–103. doi:10.1016/j.ctim.2011.02.002
- Tsourlou T, Benik A, Diplakou K, Zafeiridis A, Kellis S (2006) The Effects of a Twenty-Four-Week Aquatic training Program on Muscular Strength Performance in Healthy Elderly Women. *J Strength Cond Res* 20:811–818
- Benelli P, Ditroilo M, De Vito G (2004) Physiological Responses to fitness Activities: a comparison between Land-Based and water-Based Aerobics Exercise. *J Strength Cond Res* 18:719–722
- Campbell JA, D’Acquisto LJ, D’Acquisto DM, Cline MG (2003) Metabolic and Cardiovascular Response to Shallow Water Exercise in Young and Older Women. *Med Sci Sports Exerc* 35:675–681
- Takekuma N, Rogers ME, Watanabe E, Brechue WF, Okada A, Yamada T et al (2002) Water-based exercise improves health-related aspects of fitness in older women. *Med Sci Sports Exerc* 34:544–551

11. D'Acquisto LJ, D'Acquisto DM, Renne D (2001) Metabolic and Cardiovascular Responses in Older Women During Shallow-Water Exercise. *J Strength Cond Res* 15:12–19
12. Darby LA, Yaekle BC (2000) Physiological responses during two types of exercise performed on land and in water. *J Sports Med Phys Fitness* 40:303–311
13. Evans EM, Cureton KJ (1998) Metabolic, Circulatory and Perceptual Responses to Bench Stepping in Water. *J Strength Cond Res* 12:95–100
14. Cassady SL, Nielsen DH (1992) Cardiorespiratory responses of healthy subjects to calisthenics performed on Land versus in Water. *Phys Ther* 72:538–552
15. Christie JL, Sheldahl LM, Tristani FE (1990) Cardiovascular regulation during head-out water immersion exercise. *J Appl Physiol* 69:657–664
16. Sheldahl LM, Tristani FE, Clifford PS, Hughes CV, Sobicinsky KA, Morris RD (1987) Effect of head-out water immersion on cardiorespiratory response to dynamic exercise. *J Am Coll Cardiol* 10:1254–1258
17. Epstein M (1976) Cardiovascular and renal effects of head-out water immersion in man. *Circ Res* 39:619–629
18. Arborelius MU, Balldin I, Lilja B, Lundgren CEG (1972) Hemodynamic changes in man during immersion with the head above water. *Aerospace Med* 43:592–598
19. Rennie DW, Di Prampero P, Cerretelli P (1971) Effects of water immersion on cardiac output, heart-rate and stroke volume of man at rest and during exercise. *Med Sport* 24:223–228
20. Irandoust K, Taheri M (2015) The effects of aquatic exercise on body composition and nonspecific low back pain in elderly males. *J Phys Ther Sci* 27(2):433–435. doi:10.1589/jpts.27.433
21. Bento PC, Lopes MF, Cebolla EC, Wolf R, Rodacki A (2015) The Effects of a Water-based Training on Static and Dynamic Balance of Older Women. *Rejuvenation Res* 24 [Epub ahead of print]
22. Fedor A, Garcia S, Gunstad J (2015) The effect of a brief, water-based exercise intervention on cognitive function in older adults. *Arch Clin Neuropsychol* 30(2):139–147. doi:10.1093/arclin/acv001
23. Bergamin M, Ermolao A, Matten S, Sieverdes JC, Zaccaria M (2015) Metabolic and cardiovascular responses during aquatic exercise in water at different temperatures in older adults. *Res Q Exerc Sports* 86(2):163–171. doi:10.1080/02701367.2014.981629
24. Sato D, Seko C, Hashitomi T, Sengoku Y, Nomura T (2015) Differential effects of water-based exercise on the cognitive function in independent elderly adults. *Aging Clin Exp Res* 27(2):49–149. doi:10.1007/s40520-014-0252-9
25. Bento PC and Rodacki AL (2014) Muscle function in aged women in response to a water-based exercises program and progressive resistance training. *Geriatr Gerontol* 11. doi:10.1111/ggi.12418 [Epub ahead of print]
26. Sanders ME, Takeshima N, Rogers ME, Colado JC, Borreani S (2013) Impact of the s.w.e.a.T.TM water-exercise method on activities of daily living for older women. *J Sports Sci Med* 12(4):707–715
27. Oh S, Lim JM, Kim Y, Song W, Yoon B (2015) Comparison of the effects of water and land-based exercises on the physical function and quality of life in community-dwelling elderly people with history of falling: a single-blind, randomized controlled trial. *Arch Gerontol Geriatr* 60(2):288–293. doi:10.1016/j.archger.2014.11.001
28. Bergamin M, Ermolao A, Tolomio S, Berton L, Sergi G, Zaccaria M (2013) Water- versus land-based exercise in elderly subjects: effects on physical performance and body composition. *Clin Interv Aging* 8:1109–1117. doi:10.2147/CIA.S44198
29. Elbar O, Tzedek I, Vered E, Shvarth G, Friger M, Melzer I (2013) A water-based training program that includes perturbation exercises improves speed of voluntary stepping in older adults: a randomized controlled cross-over trial. *Arch Gerontol Geriatr* 56(1):134–140. doi:10.1016/j.archger.2012.08.003
30. Bento PCB, Pereira G, Ugrinowitsch C, Rodacki ALF (2012) The effects of a water-based exercise program on strength and functionality of older adults. *J Aging Phys Act* 20(4):469–483
31. Sato D, Kaneda K, Wakabayashi H, Shimoyama Y, Baba Y, Nomura T (2011) Comparison of once and twice weekly water exercise on various bodily functions in community-dwelling frail elderly requiring nursing care. *Arch Gerontol Geriatr* 52(3):331–335. doi:10.1016/j.archger.2010.05.002
32. Bocalini DS, Serra AJ, Rica RL, Dos Santos L (2010) Repercussions of training and detraining by water-based exercise on functional fitness and quality of life: a short-term follow-up in healthy older women. *Clinics* 65(12):1305–1309
33. Graef FI, Pinto RS, Alberton CL, de Lima WC, Krueel LF (2010) The effects of resistance training performed in water on muscle strength in the elderly. *J Strength Cond Res* 24(11):3150–3156. doi:10.1519/JSC.0b013e3181e2720d
34. Sato D, Kaneda K, Wakabayashi H, Nomura T (2009) Comparison of 2-year effects of once and twice weekly water exercise on activities of daily living ability of community-dwelling frail elderly. *Arch Gerontol Geriatr* 49(1):123–128. doi:10.1016/j.archger.2008.05.011
35. Sato D, Kaneda K, Wakabayashi H, Nomura T (2009) Comparison two-year effects of once-weekly and twice-weekly water exercise on health-related quality of life of community-dwelling frail elderly people at a day-service facility. *Disabil Rehabil* 31(2):84–93. doi:10.1080/09638280701817552
36. Bocalini DS, Serra AJ, Murad N, Levy RF (2008) Water- versus land – based exercise effects on physical fitness in older women. *Geriatr Gerontol Int* 8:265–271
37. Broman G, Quintana M, Lindberg T, Jansson E, Kaijser L (2006) High intensity deep water training can improve aerobic power in elderly women. *Eur J Appl Physiol* 98:117–123
38. Devereux K, Robertson D, Briffa NK (2005) Effects of a water-based program on women 65 years and over: a randomised controlled trial. *Aust J Physiother* 51(2):102–108
39. Takeshima N, Nakata M, Kobayashi F, Tanaka K, Pollok ML (1997) Oxygen uptake and heart rate differences between walking on land and in water in the elderly. *JAPA* 5:126–134
40. Ruoti RG, Troup JT, Berger RA (1994) The effects of non-swimming water exercises on older adults. *J Orthop Spors Phys Ther* 19:140–145
41. Pinto SS, Alberton CL, Bagatini NC, Zaffari P, Cadore EL, Radaelli R, Baroni BM, Lanferdini FJ, Ferrari R, Kanitz AC, Pinto RS, Vaz MA, Krueel LF (2015) Neuromuscular adaptations to water-based concurrent training in postmenopausal women: effects of intrasession exercise sequence. *Age (Dordr)* 37(1):9751. doi:10.1007/s11357-015-9751-7
42. Arca EA, Martinelli B, Martin LC, Waisberg CB, Franco RJ (2014) Aquatic exercise is as effective as dry land training to blood pressure reduction in postmenopausal hypertensive women. *Physiother Res Int* 19(2):93–98. doi:10.1002/pri.1565
43. Fisker AL, Waters DL, Hing WA, Steele M, Keogh JW (2015) Comparative effects of 2 aqua exercise programs on physical function, balance and perceived quality of life in older adults with osteoarthritis. *J Geriatr Phys Ther* 38(1):17–27. doi:10.1519/JPT.000000000000019
44. Ayan C, Cancela JM (2012) Feasibility of 2 different water-based exercise training programs in patients with Parkinson's disease: a pilot study. *Arch Phys Med Rehabil* 93(10):1709–1714. doi:10.1016/j.apmr.2012.03.029
45. Bidonde J, Busch AJ, Webber SC, Schachter CL, Danyliw A, Overend TJ, Richards RS, Rader T (2014) Aquatic exercise training for fibromyalgia. *Cochrane Database Syst Rev* 28;10:CD011336. doi:10.1002/14651858.CD011336

46. Nemicic T, Budisin V, Vrabec-Matkovic D, Grazio S (2013) Comparison of the effects of land-based and water-based therapeutic exercises on the range of motion and physical disability in patients with chronic low-back pain: single-blinded randomized study. *Acta Clin Croat* 52(3):321–327
47. Johansson K, Hayes S, Speck RM, Schmitz KH (2013) Water-based exercise for patients with chronic arm lymphedema: a randomized controlled pilot trial. *Am J Phys Med Rehabil* 92(4):312–319. doi:[10.1097/PHM.0b13e318278b0e8](https://doi.org/10.1097/PHM.0b13e318278b0e8)
48. Hale LA, Waters D, Herbison P (2012) A randomized controlled trial to investigate the effects of water-based exercise to improve falls risk and physical function in older adults with lower-extremity osteoarthritis. *Arch Phys Med Rehabil* 93(1):27–34. doi:[10.1016/j.apmr.2011.08.004](https://doi.org/10.1016/j.apmr.2011.08.004)
49. Ayan C, Cancela JM, Gutierrez-Santiago A, Prieto I (2014) Effects of two different exercise programs on gait parameters in individuals with Parkinson's disease: a pilot study. *Gait Posture* 39(1):648–651. doi:[10.1016/j.gaitpost.2013.08.019](https://doi.org/10.1016/j.gaitpost.2013.08.019)
50. Kim IS, Chung SH, Park YJ, Kang HY (2012) The effectiveness of an aquaerobic exercise program for patients with osteoarthritis. *Appl Nurs Res* 25(3):181–189. doi:[10.1016/j.apnr.2010.10.001](https://doi.org/10.1016/j.apnr.2010.10.001)
51. Hall J, Swinkels A, Briddon J, McCabe CS (2008) Does aquatic exercise relieve pain in adults with neurologic or musculoskeletal disease? A systematic review and meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil* 89(5):873–883. doi:[10.1016/j.apmr.2007.09.054](https://doi.org/10.1016/j.apmr.2007.09.054)
52. Robert JJ, Jones L, Bobo M (1996) The Physiologic response of exercising in the water and on land with and without the X100 Walk'N Tone Exercise Belt. *Res Q Exerc Sport* 67:310–315
53. Danneskiold-Samsøe B, Lyngberg K, Risum T, Telling M (1987) The effect of water exercise therapy given to patients with rheumatoid arthritis. *Scand J Rehabil Med* 19:31–35
54. McNamara RJ, McKeough ZJ, McKenzie DK, Alison JA (2015) Acceptability of the aquatic environment for exercise training by people with chronic obstructive pulmonary disease with physical comorbidities: additional results from a randomized controlled trial. *Physiotherapy* 101(2):187–192. doi:[10.1016/j.physio.2014.09.002](https://doi.org/10.1016/j.physio.2014.09.002)
55. McNamara RJ, McKeough ZJ, McKenzie DK, Alison JA (2013a) Water-based exercise training for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 18;12:CD008290. doi:[10.1002/14651858.CD008290](https://doi.org/10.1002/14651858.CD008290)
56. McNamara RJ, McKeough ZJ, McKenzie DK, Alison JA (2013) Water-based exercise in COPD with physical comorbidities: a randomized controlled trial. *Eur Respir J* 41(6):1284–1291. doi:[10.1183/09031936.00034312](https://doi.org/10.1183/09031936.00034312)
57. Wadell K, Sundelin G, Henriksson-Larsen K, Lundgren R (2004) High intensity physical group training in water – an effective training modality for patients with COPD. *Respir Med* 98(5):428–438
58. Choi JH, Kim BR, Joo SJ, Han EY, Kim SM, Lee SY, Yoon HM (2015) Comparison of cardiorespiratory responses during aquatic and land treadmill exercise in patients with coronary artery disease. *J Cardiopulm Rehabil Prev* 35(2):140–146. doi:[10.1097/HCR.0000000000000094](https://doi.org/10.1097/HCR.0000000000000094)
59. Teffaha D, Mourot L, Vernochet P, Ounissi F, Regnard J, Monpere C, Dugue B (2011) Relevance of water gymnastics in rehabilitation programs in patients with chronic heart failure or coronary artery disease with normal left ventricular function. *J Card Fail* 17(8):676–683. doi:[10.1016/j.cardfail.2011.04.008](https://doi.org/10.1016/j.cardfail.2011.04.008)
60. Chu KS, Eng JJ, Dawson AS, Harris JE, Ozcaplan A, Gylfadottir S (2004) Water-Based Exercise for Cardiovascular Fitness in People with Chronic Stroke: a Randomized Controlled Trial. *Arch Phys Med Rehabil* 85:870–874
61. Svendenhag J, Seger J (1992) Running on land and in water: comparative exercise physiology. *Med Sci Sports Exerc* 24:1155–1160
62. McKenzie DC, McLuckie SL (1991) Running in water as an alternative method for injured runners. *Clin J Sport Med* 1:243–346
63. Kasprzak Z and Pilaczynska-Szczesniak L (2014) Effects of regular physical exercises in the water on the metabolic profile of women with abdominal obesity. *J Hum Kinet* 8(41):71–79. doi:[10.2478/hukin-2014-0034](https://doi.org/10.2478/hukin-2014-0034). eCollection 2014
64. Rica RL, Carneiro RM, Serra AJ, Pontes Rodriguez D, Junior FL, Bocalini DS (2013) Effects of water-based exercise in obese older women: impact of short-term follow-up study on anthropometric, functional fitness and quality of life parameters. *Geriatr Gerontol* 13(1):209–214. doi:[10.1111/j.1447-0594.2012.00889.x](https://doi.org/10.1111/j.1447-0594.2012.00889.x)
65. American College of Sport Medicine (2001) Appropriate intervention strategies for weight loss and prevention of weight gain for adults. *Med Sci Sports Exerc* 33:2145–2156
66. Greene NP, Lambert BS, Greene ES, Carbuhn AF, Green JS, Crouse SF (2009) Comparative efficacy of water and land treadmill training for overweight or obese adults. *Med Sci Sports Exerc* 41(9):1808–1815. doi:[10.1249/MSS.0b013e3181a237ff](https://doi.org/10.1249/MSS.0b013e3181a237ff)
67. Jones LM, Meredith-Jones K, Legge M (2009) The effect of water-based exercise on glucose and insulin response in overweight women: a pilot study. *J Womens Health* 18(10):1653–1659. doi:[10.1089/jwh.2008.1147](https://doi.org/10.1089/jwh.2008.1147)
68. Gappmaier E, Lake W, Nelson AG, Fisher AG (2006) Aerobic exercise in water versus walking on land: effect on indices of fat reduction and weight loss of obese women. *J Sports Med Phys Fitness* 46:564–569
69. Colado JC, Tella V, Triplett NT, Gonzalez LM (2009) Effects of a short-term aquatic resistance program on strength and body composition in fit young men. *J Strength Cond Res* 23:549–559
70. Colado JC (2004) Physical conditioning in the aquatic way. Paidotribo, Barcelona
71. Pinto SS, Cadore EL, Alberton CL, Zaffari P, Bagatini NC, Baroni BM, Radaelli R, Lanferdini FJ, Colado JC, Pinto RS, Vaz MA, Bottaro M, Krueger LF (2014) Effects of intra-session exercise sequence during water-based concurrent training. *Int J Sports Med* 35(1):41–48. doi:[10.1055/s-0033-1345129](https://doi.org/10.1055/s-0033-1345129)
72. Martel GF, Harmer ML, Logan JM, Parker CB (2005) Aquatic plyometric training increases vertical jump in female volleyball players. *Med Sci Sports Exerc* 37:1814–1819
73. McWaters JG (1988) Deep water exercise for health and fitness. Public Editions, California
74. Colado JC, Triplett NT, Tella V, Saucedo P, Abellán J (2009) Effects of aquatic resistance training on health and fitness in postmenopausal women. *Eur J Appl Physiol* 106:113–122
75. Robinson LE, Devor ST, Merrick MA, Buckworth J (2004) The Effects of Land vs Aquatic Plyometrics on Power, Torque, Velocity and Muscle Soreness in women. *J Strength Cond Res* 18:84–91
76. Poyhonen T, Sipilä S, Keskinen KL, Hautala A, Savolainen J, Malkia E (2002) Effects of aquatic resistance training on neuromuscular performance in healthy women. *Med Sci Sports Exerc* 34:2103–2109
77. Colado JC, Tella V, Triplett NT (2008) A Method for Monitoring Intensity during Aquatic Resistance Exercises. *J Strength Cond Res* 22:2045–2049
78. Raffaelli C, Lanza M, Zanolla L, Zamparo P (2010) Exercise intensity of head-out water based activities (water fitness). *Eur J Appl Physiol* 109(5):829–838. doi:[10.1007/s00421-010-1419-5](https://doi.org/10.1007/s00421-010-1419-5)
79. American College of Sport Medicine (2005) ACSM's Guidelines for Exercise Testing and Prescription, 7th edn. Lippincott, Williams & Wilkins, Philadelphia

80. Borg G (1998) Borg's Perceived Exertion and Pain Scales. Human Kinetics, Champaign
81. Jackson AS, Pollock ML (1978) Generalized equations for predicting body density for men. *Br J Nutr* 40:497–504
82. Siri WE (1961) Body composition from fluid spaces and density: analysis of methods. In: Brozek J, Henschel A (eds) *Techniques for measuring body composition*. National Academy of Sciences, National Research Council, Washington, pp 223–244
83. Astrand PO, Ryhming I (1954) A Nomogram for calculation of Aerobic Capacity (Physical fitness) from Pulse Rate during Submaximal Work. *J Appl Physiol* 7:218–221
84. Morrow JR, Jackson AW, Disch JG, Mood DP (2000) Measurement and evaluation in human performance. Human Kinetics, Champaign
85. Zurc J, Pisot R, Strojnik V (2005) Gender differences in motor performance in 6.5-year-old children. *Kinesiology Slovenica* 11:90–104
86. Hawkins SA, Wiswell RA (2003) Rate and mechanism of maximal oxygen consumption decline with ageing. Implications for exercise training. *Sports Med* 33:877–888
87. Volaklis KA, Spassis AT, Tokmakidis SP (2007) Land versus water exercise in patients with coronary artery disease: effects on body composition, blood lipids and physical fitness. *Am Heart Journal* 154:559–566
88. Peterson MJ, Czerwinski SA, Siervogel RM (2003) Development and validation of skinfold-thickness prediction equations with a 4-compartment model. *Am J Clin Nutr* 77:1186–1191
89. Silva AM, Fields DA, Quiterio AL, Sardina LB (2009) Are skinfold-based models accurate and suitable for assessing changes in body composition in highly trained athletes? *J Strength Cond Res* 23:1688–1696
90. España-Romero V, Ruiz JR, Ortega FB, Artero EG, Rodriguez GV, Moreno LA et al (2009) Body fat measurement in elite sport climbers: comparison of skinfold thickness equations with dual energy X-ray absorptiometry. *Journal of Sports Sciences* iFirst article:1–9
91. Ellis KJ (1996) Measuring body fatness in children and young adults: comparison of bioelectric impedance analysis, total body electrical conductivity, and dual energy X-ray absorptiometry. *Int J Obes* 20:866–873
92. Gutin B, Litaker M, Islam S, Manos T, Smith C, Treiber F (1996) Body-composition measurement in 9–11-y-old children by dual-energy X-ray absorptiometry, skinfold-thickness measurements, and bioimpedance analysis. *Am J Clin Nutr* 63:287–292
93. Duz S, Kocak M, Korkusuz F (2009) Evaluation of body composition using three different methods compared to dual energy X-ray absorptiometry. *Euro J Sport Sci* 9:181–190
94. Taunton JE, Rhodes EC, Wolski LA, Donnelly M, Warren J, Elliot J et al (1996) Effects of Land-Based and Water-Based Fitness Programs on the Cardiovascular Fitness, Strength and Flexibility of Women Aged 65–75 Years. *Gerontology* 42:204–210
95. Cider A, Schaufelberger M, Sunnerhagen KS, Andersson B (2003) Hydrotherapy—a new approach to improve function in the older patient with chronic heart failure. *Eur J of Heart Failure* 5:527–535
96. American College of Sport Medicine (2005) *Health-related physical fitness assessment manual*. Lippincott, Williams & Wilkins, Philadelphia
97. Micallef C (2014) The effectiveness of an 8-week Zumba programme for weight reduction in a group of Maltese overweight and obese women. *Sport Sciences for Health* 10:211–217. doi:10.1007/s11332-014-0195-8