

1. Sasco JA, Fontaniere B. A population based series of 10 male breast cancer cases. *Eur J Cancer Clin Oncol* 1991, 12, 1713.
2. Isquierdo MA, Alonso C, Andres LDE, Ojeda B. Male breast cancer: report of 50 cases. *Acta Oncologica* 1994, 33(7), 767-771.
3. Ciatto S, Iossa A, Bonardi R, Pacini P. Male breast carcinoma: review of a multicentre series of 150 cases. *Tumori* 1990, 76, 555-558.
4. Wagner JI, Thomas CR, Koh W-J, Rudolph RH. Carcinoma of the male breast: update 1994. *Med Pediatr Oncol* 1995, 24(2), 123-32.
5. Hecht JR, Winchester DJ. Male breast cancer. *Am J Clin Pathol* 1994, 102 (Suppl. 1), S25-S30.
6. Thomas DB. Breast cancer in men. *Epidemiol Rev* 1993, 15, 220-231.
7. Sasco AJ, Lowenfels AB, Pasker-De Jong P. Epidemiology of male breast cancer. A meta-analysis of published case control studies and discussion of selected aetiological factors. *Int J Cancer* 1993, 53, 538-549.
8. Schon M, Zaiac M, Schilag OM. Male breast cancer. *Onkologie* 1995, 18(1), 16-21.
9. Anelli A, Anelli TFM, Youngson B, Rosen PP, Borgen PI. Mutations of the P53 gene in male breast cancer. *Cancer* 1995, 75(9), 2233-2238.
10. Thorlacius S, Tryggvadottir L, Olafsdottir GH, et al. Linkage to BRCA2 region in hereditary male breast cancer. *Lancet* 1995, 346, 544-545.

European Journal of Cancer Vol. 32A, No. 8, pp. 1436-1437, 1996.
 Copyright © 1996 Elsevier Science Ltd. All rights reserved.
 Printed in Great Britain
 0959-8049/96 \$15.00 + 0.00

PII: S0959-8049(96)00126-8

High-dose Recombinant Interleukin-2/Verapamil Combination in Advanced Cancer

P. Tagliaferri,¹ P. Correale,¹ M. Mottola,²
 G. de Simone,³ V. Montesarchio,¹
 E. Matano,¹ A. Rea,¹ A. Morabito,¹
 M. Famiani,¹ F. Ciardiello,¹ G. Tortora,¹
 M. Caraglia,¹ C. Barile,¹ G. Palmieri¹ and
 A.R. Bianco¹

¹Cattedra di Oncologia Medica, Dipartimento di Endocrinologia ed Oncologia Molecolare e Clinica;

²Cattedra di Cardiocirurgia; and ³Istituto di Medicina Interna e Malattie Dismetaboliche, Università "Federico II", Via S. Pansini 5, 80131 Naples, Italy

rIL-2-BASED IMMUNOTHERAPY is a promising approach for patients with advanced melanoma and renal cell carcinoma [1]. IL-2 generates lymphokine-activated killing activity (LAK) by peripheral blood lymphocytes (PBL) which acquire the capacity of MHC-unrestricted killing of tumour cells [2-4]. We have demonstrated that human tumour cell lines, resistant to LAK-cytotoxicity *in vitro*, become sensitive to cytotoxic lymphocytes after 24-48 h exposure to the calcium channel blocker verapamil [5]. Verapamil is active at concentrations

(10-100 nM) which can be reached *in vivo* [6]. Therefore, we designed a Phase I clinical study on rIL-2/verapamil combination. After committee approval, written informed consent was obtained from each patient before study entry.

All patients enrolled in the study were given the same verapamil dose, while the dose of rIL-2 given by continuous infusion for 72 h starting day 2 was escalated in the different cohorts of patients (Table 1). Dose escalation was allowed if a minimum of 3 patients had been treated at each dose level of infusional rIL-2 without grade 3-4 toxicities in 2 of 3 patients. Dose escalation over 18 IMU/m²/day, which can be considered a standard dose [7] and close to the maximum tolerated dose (MTD) [8] of rIL-2 by continuous infusion, was not planned. 10 IMU/m²/day rIL-2 were given by subcutaneous administration on days 6-10 to all patients who entered the study. Verapamil was administered at the dose of 0.2 mg/kg/h for 3 consecutive days by continuous infusion after a loading bolus dose of 0.15 mg/kg. 18 patients—8 males, 10 females, 10 with malignant melanoma, 7 with renal cell carcinoma, 1 with colon cancer—were enrolled in the study and completed the two planned courses of rIL-2/verapamil. Median age was 49 years (range 22-72). 3 additional patients were treated with 9 IMU/m² rIL-2 to further explore the rIL-2 dose which we considered, on the basis of biological activity (see below), as the recommended one for Phase II trials (Table 1).

Grade 1 elevation of BUN and creatinine occurred in 5 patients (28%), respectively, while grade 2 increase of creatinine occurred in 1 patient (6%). Oliguria occurred in 11 patients (61%) and inversion of urine electrolyte in 13 patients (72%). 2 patients experienced grade 2-3 hypotension which resulted in only temporary suspension of therapy. Grade 1 hypotension occurred in 9 patients (50%), but did not result in discontinuation of drug infusion. Synus tachycardia was observed in 7 patients (39%), first degree atrioventricular (AV) heart block in 2 patients (11%), AV dissociation in 1 (6%) and premature ventricular beats in 1. 1 patient had synus bradycardia immediately after bolus verapamil injection which resolved spontaneously with temporary discontinuation of verapamil infusion. Fever occurred in 14 (78%) patients, dyspnoea in 1 (6%), abdominal pain in 13 (72%), anaemia in 10 (56%), nausea and vomiting in 4 (22%), constipation in 4 (22%), increase of serum bilirubin in 5 (28%), diarrhoea in 1 (6%).

We also evaluated the immunomodulatory properties of the rIL-2/verapamil combination by determining the number of circulating eosinophils and induction of cytotoxic activity against the human leukaemia K562 cells. Both peripheral blood lymphocytes cytotoxic activity and circulating eosinophils were maximally increased at the rIL-2 dose of 9 IMU/m²/day. We observed 1 complete remission in a patient with renal cell carcinoma, lasting 30+ months, and 2 partial remissions (renal cell carcinoma and melanoma), lasting 6 and 9 months. Minor objective responses were recorded in 4 patients (1 with melanoma and 3 with renal cell carcinoma). Disease progressed in 5 patients (Table 1). In conclusion, we demonstrated that a rIL-2 dose of 18 IMU/m² could be administered in combination with high-dose verapamil in absence of cumulative toxicity. Furthermore, this combination has biomodulatory properties and antitumour activity and deserves evaluation in Phase II clinical studies.

Correspondence to P. Tagliaferri.

Received 31 Aug. 1995; revised 21 Feb. 1996; accepted 25 Mar. 1996.

1. Rosenberg SA, Lotze MT, Yang J, et al. Experience with the use of high dose interleukin-2 in the treatment of 652 cancer patients. *Ann Surg* 1989, 210, 474-483.

Table 1. Tumour types and responses to rIL-2/verapamil combination therapy

rIL-2 (IMU/m ² /day)	Tumour type	Sites of disease	Clinical response
1.2	Renal cell carcinoma	Liver, nodes	PD
1.2	Melanoma	Liver, bone	NC
1.2	Melanoma	Skin, nodes	NC
2.4	Colon carcinoma	Liver, lung	NC
2.4	Melanoma	Liver, breast	NC
2.4	Melanoma	Liver, spleen, bone, lung	NC
4.8	Melanoma	Nodes	PD
4.8	Melanoma	Pleura, lung, liver, nodes Uterus	PD
4.8	Renal cell carcinoma	Mediastinal nodes	CR
9	Melanoma	Adrenal gland, nodes	MR
9	Renal cell carcinoma	Lung	MR
9	Melanoma	Liver	PR
9	Renal cell carcinoma	Lung	MR
9	Melanoma	Nodes	NC
9	Renal cell carcinoma	Lung, nodes	PD
18	Melanoma	Lung, skin, nodes	PD
18	Renal cell carcinoma	Lung, bone	MR
18	Renal cell carcinoma	Lung	PR

CR, complete remission; PR, partial remission; MR, minimal response; NC, no change; PD, progression of disease.

- Lotze MT, Matory SE, Ettinghausen SE, *et al.* *In vivo* administration of purified recombinant interleukin-2. II. Half life, immunological effects and expansion of peripheral lymphoid cells with recombinant IL-2. *J Immunol* 1985, **135**, 2865-2875.
- Hank JA, Kohler PC, Weill-Hillman G. *In vivo* induction of the lymphokine activated killer phenomenon: interleukin-2 dependent human non-major histocompatibility complex restricted cytotoxicity generated *in vivo* during administration of human recombinant interleukin-2. *Cancer Res* 1988, **48**, 1965-1971.
- Grimm EA, Mazumber A, Zhang HZ, *et al.* Lymphocyte activated killing phenomenon: lysis of natural killer resistant fresh solid tumor cells by interleukin-2 activated autologous human peripheral blood lymphocytes. *J Exp Med* 1982, **155**, 1823-1841.
- Correale P, Tagliaferri P, Celio L, *et al.* Verapamil upregulates sensitivity of human colon and breast cancer cells to LAK cytotoxicity *in vitro*. *Eur J Cancer* 1991, **27A**, 1393-1395.
- Pennock GD, Dalton WS, Roeske WR. Systemic toxic effects associated to high dose verapamil infusion and chemotherapy administration. *J Natl Cancer Inst* 1991, **83**, 105-110.
- Negrier S, Philip T, Stoter G. Interleukin-2 with or without LAK cells in metastatic renal cell carcinoma: a report of a European multicentre study. *Eur J Cancer Clin Oncol* 1989, **25** (Suppl. 3), 21-28.
- Gaynor ER, Weiss GR, Margolin KA, *et al.* Phase I study of high-dose continuous-infusion recombinant interleukin-2 and autologous lymphokine-activated killer cells in patients with metastatic or unresectable malignant melanoma and renal cell carcinoma. *J Natl Cancer Inst* 1990, **82**, 1397-1402.

Acknowledgements—Supported by the Associazione Italiana per la Ricerca sul Cancro (A.I.R.C.) and the Consiglio Nazionale Ricerche (C.N.R., A.C.R.O. project). Michele Caraglia is recipient of a fellowship from A.I.R.C. We thank Dr Michelino De Laurentiis, University of Naples, for useful discussions and critical reading of the manuscript.

European Journal of Cancer Vol. 32A, No. 8, pp. 1437-1438, 1996.
Copyright © 1996 Elsevier Science Ltd. All rights reserved
Printed in Great Britain
0959-8049/96 \$15.00 + 0.00

PII: S0959-8049(96)00128-1

Activity of Gemcitabine in Platinum-resistant Ovarian Germ Cell Cancer

K. Maas, T. Daikeler, L. Kanz and C. Bokemeyer

Eberhard Karls University, Department of Internal Medicine II, Otfried-Müller-Str. 10, 72076 Tübingen, Germany

OVARIAN GERM cell cancer (OGCT) is a rare disease accounting for less than 5% of all ovarian carcinomas. The importance of the disease is due to its relatively high incidence among children and young adults and the potential for cure by the use of platinum-containing chemotherapy [1]. The current recommendation of the Gynecologic Oncology Group (U.S.A.) for the treatment of advanced OGCT consists of cytoreductive surgery followed by three to six cycles of chemotherapy with bleomycin, etoposide and cisplatin (BEP) resulting in a 75% cure rate for advanced stage disease [1]. However, treatment of platinum-refractory OGCT remains difficult [2]. Paclitaxel is among the few drugs that have shown activity in platinum-resistant ovarian and testicular germ cell cancer [3, 4]. We report here the case of a patient with

Correspondence to K. Maas.
Received 4 Mar. 1996; accepted 7 Mar. 1996.