

World Journal of *Stem Cells*

World J Stem Cells 2018 July 26; 10(7): 78-105





EDITORIAL

- 78 Ectopic expression of the osteogenic master gene *RUNX2* in melanoma
Valenti MT, Dalle Carbonare L, Mottes M

REVIEW

- 82 Stem cell therapy for faecal incontinence: Current state and future perspectives
Trébol J, Carabias-Orgaz A, García-Arranz M, García-Olmo D

ABOUT COVER

Editorial Board Member of *World Journal of Stem Cells*, Manabu Akahane, MD, PhD, Associate Professor, Public Health, Health Management and Policy, Nara Medical University School of Medicine, Nara 634-8521, Japan

AIM AND SCOPE

World Journal of Stem Cells (*World J Stem Cells*, *WJSC*, online ISSN 1948-0210, DOI: 10.4252), is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJSC covers topics concerning all aspects of stem cells: embryonic, neural, hematopoietic, mesenchymal, tissue-specific, and cancer stem cells; the stem cell niche, stem cell genomics and proteomics, and stem cell techniques and their application in clinical trials.

We encourage authors to submit their manuscripts to *WJSC*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Stem Cells (*WJSC*) is now indexed in PubMed, PubMed Central, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports/Science Edition, Biological Abstracts, and BIOSIS Previews. The 2018 Edition of Journal Citation Reports cites the 2017 impact factor for *WJSC* as 4.376 (5-year impact factor: N/A), ranking *WJSC* as 7 among 24 journals in Cell and Tissue Engineering (quartile in category Q2), and 65 among 190 journals in Cell Biology (quartile in category Q2).

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Wen-Wen Tan*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Stem Cells

ISSN
 ISSN 1948-0210 (online)

LAUNCH DATE
 December 31, 2009

FREQUENCY
 Monthly

EDITORS-IN-CHIEF
Tong Cao, BM BCh, DDS, PhD, Associate Professor, Doctor, Department of Oral Sciences, National University of Singapore, Singapore 119083, Singapore

EDITORIAL BOARD MEMBERS
 All editorial board members resources online at <http://www.wjgnet.com/1948-0210/editorialboard.htm>

EDITORIAL OFFICE
 Jin-Lei Wang, Director
World Journal of Stem Cells

Baishideng Publishing Group Inc
 7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: editorialoffice@wjgnet.com
 Help Desk: <http://www.wjgnet.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
 Baishideng Publishing Group Inc
 7901 Stoneridge Drive, Suite 501,
 Pleasanton, CA 94588, USA
 Telephone: +1-925-2238242
 Fax: +1-925-2238243
 E-mail: bpgoffice@wjgnet.com
 Help Desk: <http://www.wjgnet.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
 July 26, 2018

COPYRIGHT
 © 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non-commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
 All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.wjgnet.com>

Ectopic expression of the osteogenic master gene *RUNX2* in melanoma

Maria Teresa Valenti, Luca Dalle Carbonare, Monica Mottes

Maria Teresa Valenti, Luca Dalle Carbonare, Department of Medicine, University of Verona, Verona 37100, Italy

Monica Mottes, Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona 37100, Italy

ORCID number: Maria Teresa Valenti (0000-0003-1166-8033); Luca Dalle Carbonare (0000-0003-3263-6671); Monica Mottes (0000-0002-7390-2246).

Author contributions: Valenti MT and Mottes M conceived the study; Valenti MT, Dalle Carbonare L and Mottes M drafted the manuscript; all authors approved the final version of the paper.

Conflict-of-interest statement: The authors have no conflict of interest to declare.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Maria Teresa Valenti, BSc, PhD, Chief Technician, Research Fellow, Department of Medicine, University of Verona, Ple Scuro 10, Verona 37100, Italy. mariateresa.valenti@univr.it
Telephone: +39-45-8128450

Received: May 23, 2018

Peer-review started: May 25, 2018

First decision: June 5, 2018

Revised: June 11, 2018

Accepted: June 28, 2018

Article in press: June 28, 2018

Published online: July 26, 2018

Abstract

The transcription factor *RUNX2* is the osteogenic master gene expressed in mesenchymal stem cells during osteogenic commitment as well as in pre-osteoblasts and early osteoblasts. However, *RUNX2* is also ectopically expressed in melanoma and other cancers. Malignant melanoma (MM) is a highly metastatic skin cancer. The incidence of MM has increased considerably in the past half-century. The expression levels and mutation rates of genes such as *BRAF*, *KIT*, *NRAS*, *PTEN*, *P53*, *TERT* and *MITF* are higher in melanoma than in other solid malignancies. Additionally, transcription factors can affect cellular processes and induce cellular transformation since they control gene expression. Recently, several studies have identified alterations in *RUNX2* expression. In particular, the regulation of *KIT* by *RUNX2* and the increased expression of *RUNX2* in melanoma specimens have been shown. Melanocytes, whose transformation results in melanoma, arise from the neural crest and therefore show "stemness" features. *RUNX2* plays an important role in the re-activation of the MAPK and PI-3K/AKT pathways, thus endowing melanoma cells with a high metastatic potential. In melanoma, the most frequent metastatic sites are the lung, liver, brain and lymph nodes. In addition, bone metastatic melanoma has been described. Notably, studies focusing on *RUNX2* may contribute to the identification of an appropriate oncotarget in melanoma.

Key words: *RUNX2*; Mesenchymal stem cells; Epithelial mesenchymal transition; Bone; Melanoma

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: In addition to its physiological expression in osteogenic cells, *RUNX2* ectopic expression is also reported in several cancers. Melanoma cells, which arise from the neural crest, are not typical epithelial cells and

exhibit “stemness” features. For this reason the epithelial mesenchymal transition process may be enhanced in melanoma cells. *RUNX2* expression in such context increases the migration and invasiveness of melanoma cells; it can therefore be considered a new oncotarget in melanoma.

Valenti MT, Dalle Carbonare L, Mottes M. Ectopic expression of the osteogenic master gene *RUNX2* in melanoma. *World J Stem Cells* 2018; 10(7): 78-81 Available from: URL: <http://www.wjgnet.com/1948-0210/full/v10/i7/78.htm> DOI: <http://dx.doi.org/10.4252/wjsc.v10.i7.78>

INTRODUCTION

In recent times, the incidence of malignant melanoma (MM) has increased. The increased number of MM patients may be ascribed to lifestyle and environmental changes. Unfortunately, the mortality rate of MM is very high, as MM is highly metastatic and genetically resistant to apoptosis. The surgical treatment of skin lesions before cell spreading is considered a successful therapeutic approach. However, after metastasis and dissemination of cells has occurred, the therapeutic tools are limited. Despite an initial positive response to treatment, melanoma cells become resistant to chemotherapy^[1]. Many studies performed in animal models and in cell culture have identified the complex mechanism involved in the metastatic progression of melanoma. Mutations in master transcription regulators such as *BRAF*, *KIT*, *NRAS*, *PTEN*, *P53*, *TERT*, and *MITF* have been found frequently^[2]. In particular, the microphthalmia-associated transcription factor (*MITF*) is an important oncogene, and it plays a critical role in melanoma transformation due to its role in melanocyte proliferation and differentiation^[3]. However, transcription factors in general may affect cellular processes and thereby induce cellular transformation. Among them, *RUNX2*, the master gene of osteogenic differentiation, has been shown to be involved in the transformation and metastatic progression of melanoma^[4-6]. *RUNX2*, together with *RUNX1* and *RUNX3*, belongs to the *RUNT*-related gene family^[7]. These heterodimeric transcription factors have a DNA-binding “A” subunit and a non-DNA binding “B” subunit. The genomic and cDNA structures of the *RUNT* family members are evolutionarily conserved^[8]. *RUNX2* is located on human chromosome 6; its coding sequence is organized in eight exons and controlled by two promoters. Isoforms may originate by the alternative use of promoters or by the alternative splicing of exons; however, the DNA-binding domain remains invariant^[7]. *RUNX2* induces the commitment of mesenchymal stem cells to the osteogenic lineage by acting either as a monomer or as a heterodimeric complex, which shows a higher DNA binding activity^[7]. Interestingly, differences in *RUNX2* expression are associated with the skeletal differ-

ences between modern humans and Neanderthals^[9]. Otto *et al.*^[10] using an in vivo model, showed that the ossification process is lacking in *RUNX2* KO mice. *RUNX2* is expressed not only in mesenchymal stem cells to commit these cells to osteogenic differentiation but also in pre-osteoblasts and early osteoblasts. It is then downregulated, as its continuous expression would prevent osteoblast maturation. This aspect is important for bone remodelling. In acromegaly patients, who are characterized by an excessive production of GH and IGF-1, a high fracture risk occurs; we have demonstrated that *RUNX2* gene overexpression induces bone loss and impairs bone remodelling in these patients^[11].

RUNX2 OVEREXPRESSION IN CANCER

The *RUNX* family members are frequently deregulated in human cancer. Runx proteins are involved in critical cellular processes such as DNA repair, apoptosis, hypoxia, the inflammatory response, EMT, stem cell functions, and oncogene-induced senescence, and they interact with different signalling pathways such as the RAS-ERK, WNT, TGF β , BMP and Notch pathways^[12]. In addition, *RUNX2* is involved in osteolytic diseases, neoangiogenic processes, and invasion and metastasis of solid tumours^[4]. *RUNX2* can be associated with HDA6 and can thereby prevent the pro-apoptotic activity of P53 through its deacetylation^[13]. *RUNX2* is ectopically expressed in several solid tumours such as breast cancer, pancreatic cancer, prostate cancer, lung cancer, and ovarian epithelial cancer; therefore, we have proposed *RUNX2* as a mesenchymal stem marker for cancer^[14,15]. Interestingly, we observed higher levels of *RUNX2* expression in cancer patients with bone metastases compared to patients without bone metastases^[15]. In thyroid cancer patients, we observed higher expression of *RUNX2* in cancer tissue as well as in circulating mRNA compared to those levels of *RUNX2* expression in controls^[16]. In addition, *RUNX2* expression was higher in cancer patients with microcalcifications compared to patients without microcalcifications^[16]. Finally, it has been reported that *RUNX2* induces its target genes SDF-1, CXCR7 and BSP, promoting bone homing^[17].

RUNX2 IN MELANOMA

Ectopic expression of *RUNX2* in melanoma tissue and in melanoma cell lines has been shown to be associated with bone sialoprotein (BSP)^[18]. BSP has been associated with tumour invasiveness^[18]. *RUNX2* also regulates several matrix metalloproteinases that are involved in melanoma progression; therefore, *RUNX2* may be regarded as a common regulator of both BSP and metalloproteinases^[19]. This finding could also explain the expression of other bone proteins, such as osteopontin and osteocalcin, in melanoma^[20,21]. TGF β , which induces metastasis formation in advanced melanoma stages^[22], can upregulate the expression of *RUNX2* in melanoma

cell lines^[23]. In contrast, the tumour suppressor p14ARF reduces *RUNX2* expression^[24]. Loss-of-function *p14ARF* mutations lead to increased expression of *RUNX2* and melanoma progression^[24]. Ectopic expression of *RUNX2* in melanoma has been associated with invasiveness; it has been demonstrated that *RUNX2* overexpression mediates the migration ability of melanoma cells^[6]. Notably *RUNX2* stimulates *VEGF* gene expression^[25] and the upregulation of angiogenesis, consequent to *RUNX2* overexpression, may represent a critical step for tumor metastasis.

Furthermore, *RUNX2* overexpression upregulates *SOX9*, *SNAI2* and *SMAD3*. These transcription factors are involved in EMT as well as in cytoskeletal remodelling and thus enhance the motility and invasive potential of cancer cells^[17]. In melanoma, the overexpression of *EGF-R*, *PDGFR β* , *AXL* and *IGF-1R* induces the reactivation of the MAPK and PI3K/AKT pathways, which are involved in migration and invasion processes^[26]. The reactivation of these pathways is due to tyrosine kinase receptor-based autocrine loops, in which *RUNX2* plays a pivotal role^[5]. The MAPK and AKT pathways regulate most physiological processes, such as proliferation, differentiation and cell survival. It has been suggested that the MAPK and AKT pathways are strongly associated with each other in melanoma; the link between the PI3K/AKT (AKT pathway) and MAPK/ERK1/2 (MAPK pathway) cascades is well known^[27]. AKT enhances *RUNX2* expression either directly or by inactivating *RUNX2* regulators. Similarly, *RUNX2* activates the PI3K/AKT pathway by regulating different components^[4]. This reciprocal activation induces tumour progression and aggressiveness. In addition, *RUNX2* promotes the crosstalk between MAPK and AKT through EGFR, as has been shown in human mammary epithelial cells^[28] through the reduction of ERK-mediated inhibition of EGFR and the AKT pathway. The lung, liver, brain and lymph nodes are the most frequent metastatic sites for melanoma cells^[29]. Since melanocytes arise from the neural crest, they are not typical epithelial cells; they show "stemness" features. This last finding could partially explain the high metastatic potential of melanoma^[29]. Solid malignancies such as breast, prostate and lung cancers form metastases in bone. However, metastatic bone diseases from MM are under-investigated. The data in the literature report an incidence of skeletal metastases in MM patients that ranges from 0.85% to 18.6%^[30]. Recently, Zekri *et al.*^[31], managing MM patients at two referral cancer centres in England, showed that bone metastases occur in 4.1% of patients at all stages of MM. In addition, another study showed that 17.2% of metastatic melanoma patients had metastases in bone tissue.

CONCLUSION

MM accounts for less than 5% of cutaneous malignancies. However, its incidence has been increasing con-

siderably in the past half-century. Therefore, extensive research focusing on the genes involved in melanoma transformation and on the modulation of these genes is needed. Several studies have highlighted the involvement of *RUNX2*, the master gene of osteogenic differentiation, in melanoma. The invasiveness and metastatic features conferred by *RUNX2* seem to be related to its ability to promote EMT. In addition, melanocytes are not typical epithelial cells, and their stemness features could explain melanoma invasiveness. Since metastatic melanoma has a poor prognosis, new and more effective therapeutic tools should be developed in order to implement current therapies. In this context, the exploitation of molecules actually involved in melanoma, such as *RUNX2*, represents an important frontier for the identification of new oncotargets. Transcription factors have been ignored thus far by the pharmaceutical industry; nonetheless, many studies have identified their central role in cellular transformation. We believe that future efforts directed towards unravelling the complex roles of *RUNX2* may contribute to the identification of new therapeutic tools that improve the quality of life of melanoma patients.

REFERENCES

- 1 **Whiteman DC**, Pavan WJ, Bastian BC. The melanomas: a synthesis of epidemiological, clinical, histopathological, genetic, and biological aspects, supporting distinct subtypes, causal pathways, and cells of origin. *Pigment Cell Melanoma Res* 2011; **24**: 879-897 [PMID: 21707960 DOI: 10.1111/j.1755-148X.2011.00880.x]
- 2 **Damsky WE**, Bosenberg M. Melanocytic nevi and melanoma: unraveling a complex relationship. *Oncogene* 2017; **36**: 5771-5792 [PMID: 28604751 DOI: 10.1038/ncr.2017.189]
- 3 **Levy C**, Khaled M, Fisher DE. MITF: master regulator of melanocyte development and melanoma oncogene. *Trends Mol Med* 2006; **12**: 406-414 [PMID: 16899407 DOI: 10.1016/j.molmed.2006.07.008]
- 4 **Cohen-Solal KA**, Boregowda RK, Lasfar A. *RUNX2* and the PI3K/AKT axis reciprocal activation as a driving force for tumor progression. *Mol Cancer* 2015; **14**: 137 [PMID: 26204939 DOI: 10.1186/s12943-015-0404-3]
- 5 **Boregowda RK**, Medina DJ, Markert E, Bryan MA, Chen W, Chen S, Rabkin A, Vido MJ, Gunderson SI, Chekmareva M, Foran DJ, Lasfar A, Goydos JS, Cohen-Solal KA. The transcription factor *RUNX2* regulates receptor tyrosine kinase expression in melanoma. *Oncotarget* 2016; **7**: 29689-29707 [PMID: 27102439 DOI: 10.18632/oncotarget.8822]
- 6 **Boregowda RK**, Olabisi OO, Abushahba W, Jeong BS, Haenssen KK, Chen W, Chekmareva M, Lasfar A, Foran DJ, Goydos JS, Cohen-Solal KA. *RUNX2* is overexpressed in melanoma cells and mediates their migration and invasion. *Cancer Lett* 2014; **348**: 61-70 [PMID: 24657655 DOI: 10.1016/j.canlet.2014.03.011]
- 7 **Dalle Carbonare L**, Innamorati G, Valenti MT. Transcription factor *Runx2* and its application to bone tissue engineering. *Stem Cell Rev* 2012; **8**: 891-897 [PMID: 22139789 DOI: 10.1007/s12015-011-9337-4]
- 8 **Otto F**, Lübbert M, Stock M. Upstream and downstream targets of *RUNX* proteins. *J Cell Biochem* 2003; **89**: 9-18 [PMID: 12682904 DOI: 10.1002/jcb.10491]
- 9 **Benítez-Burraco A**, Boeckx C. Possible functional links among brain- and skull-related genes selected in modern humans. *Front Psychol* 2015; **6**: 794 [PMID: 26136701 DOI: 10.3389/fpsyg.2015.00794]

- 10 **Otto F**, Thornell AP, Crompton T, Denzel A, Gilmour KC, Rosewell IR, Stamp GW, Beddington RS, Mundlos S, Olsen BR, Selby PB, Owen MJ. *Cbfa1*, a candidate gene for cleidocranial dysplasia syndrome, is essential for osteoblast differentiation and bone development. *Cell* 1997; **89**: 765-771 [PMID: 9182764 DOI: 10.1016/S0092-8674(00)80259-7]
- 11 **Valenti MT**, Mottes M, Cheri S, Deiana M, Micheletti V, Cosaro E, Davi MV, Francia G, Dalle Carbonare L. Runx2 overexpression compromises bone quality in acromegalic patients. *Endocr Relat Cancer* 2018; **25**: 269-277 [PMID: 29295822 DOI: 10.1530/ERC-17-0523]
- 12 **Ito Y**. RUNX genes in development and cancer: regulation of viral gene expression and the discovery of RUNX family genes. *Adv Cancer Res* 2008; **99**: 33-76 [PMID: 18037406 DOI: 10.1016/S0065-230X(07)99002-8]
- 13 **Ozaki T**, Nakamura M, Shimozato O. Novel Implications of DNA Damage Response in Drug Resistance of Malignant Cancers Obtained from the Functional Interaction between p53 Family and RUNX2. *Biomolecules* 2015; **5**: 2854-2876 [PMID: 26512706 DOI: 10.3390/biom5042854]
- 14 **Valenti MT**, Serafini P, Innamorati G, Gili A, Cheri S, Bassi C, Dalle Carbonare L. Runx2 expression: A mesenchymal stem marker for cancer. *Oncol Lett* 2016; **12**: 4167-4172 [PMID: 27895787 DOI: 10.3892/ol.2016.5182]
- 15 **Valenti MT**, Mori A, Malerba G, Dalle Carbonare L. Mesenchymal stem cells: A new diagnostic tool? *World J Stem Cells* 2015; **7**: 789-792 [PMID: 26131309 DOI: 10.4252/wjsc.v7.i5.789]
- 16 **Dalle Carbonare L**, Frigo A, Francia G, Davi MV, Donatelli L, Stranieri C, Brazzarola P, Zatelli MC, Menestrina F, Valenti MT. Runx2 mRNA expression in the tissue, serum, and circulating non-hematopoietic cells of patients with thyroid cancer. *J Clin Endocrinol Metab* 2012; **97**: E1249-E1256 [PMID: 22511796 DOI: 10.1210/jc.2011-2624]
- 17 **Baniwal SK**, Khalid O, Gabet Y, Shah RR, Purcell DJ, Mav D, Kohn-Gabet AE, Shi Y, Coetzee GA, Frenkel B. Runx2 transcriptome of prostate cancer cells: insights into invasiveness and bone metastasis. *Mol Cancer* 2010; **9**: 258 [PMID: 20863401 DOI: 10.1186/1476-4598-9-258]
- 18 **Riminucci M**, Corsi A, Peris K, Fisher LW, Chimenti S, Bianco P. Coexpression of bone sialoprotein (BSP) and the pivotal transcriptional regulator of osteogenesis, *Cbfa1/Runx2*, in malignant melanoma. *Calcif Tissue Int* 2003; **73**: 281-289 [PMID: 14667142 DOI: 10.1007/s00223-002-2134-y]
- 19 **Hofmann UB**, Westphal JR, Zendman AJ, Becker JC, Ruiter DJ, van Muijen GN. Expression and activation of matrix metalloproteinase-2 (MMP-2) and its co-localization with membrane-type 1 matrix metalloproteinase (MT1-MMP) correlate with melanoma progression. *J Pathol* 2000; **191**: 245-256 [PMID: 10878545 DOI: 10.1002/1096-9896(2000)9999:9999<::AID-PATH632>3.0.CO;2-#]
- 20 **Kiss T**, Ecsedi S, Vizkeleti L, Koroknai V, Emri G, Kovács N, Adany R, Balazs M. The role of osteopontin expression in melanoma progression. *Tumour Biol* 2015; **36**: 7841-7847 [PMID: 25944164 DOI: 10.1007/s13277-015-3495-y]
- 21 **Salvatierra E**, Alvarez MJ, Leishman CC, Rivas Baquero E, Lutzky VP, Chuluyan HE, Podhajcer OL. SPARC Controls Melanoma Cell Plasticity through Rac1. *PLoS One* 2015; **10**: e0134714 [PMID: 26248315 DOI: 10.1371/journal.pone.0134714]
- 22 **Lasfar A**, Cohen-Solal KA. Resistance to transforming growth factor β -mediated tumor suppression in melanoma: are multiple mechanisms in place? *Carcinogenesis* 2010; **31**: 1710-1717 [PMID: 20656791 DOI: 10.1093/carcin/bgq155]
- 23 **Mohammad KS**, Javelaud D, Fournier PG, Niewolna M, McKenna CR, Peng XH, Duong V, Dunn LK, Mauviel A, Guise TA. TGF-beta-RI kinase inhibitor SD-208 reduces the development and progression of melanoma bone metastases. *Cancer Res* 2011; **71**: 175-184 [PMID: 21084275 DOI: 10.1158/0008-5472.CAN-10-2651]
- 24 **Packer LM**, Pavey SJ, Boyle GM, Stark MS, Ayub AL, Rizos H, Hayward NK. Gene expression profiling in melanoma identifies novel downstream effectors of p14ARF. *Int J Cancer* 2007; **121**: 784-790 [PMID: 17450523 DOI: 10.1002/ijc.22725]
- 25 **Zelzer E**, Glotzer DJ, Hartmann C, Thomas D, Fukai N, Soker S, Olsen BR. Tissue specific regulation of VEGF expression during bone development requires *Cbfa1/Runx2*. *Mech Dev* 2001; **106**: 97-106 [PMID: 11472838 DOI: 10.1016/S0925-4773(01)00428-2]
- 26 **Yadav V**, Denning MF. Fyn is induced by Ras/PI3K/Akt signaling and is required for enhanced invasion/migration. *Mol Carcinog* 2011; **50**: 346-352 [PMID: 21480388 DOI: 10.1002/mc.20716]
- 27 **Pappalardo F**, Russo G, Candido S, Pennisi M, Cavalieri S, Motta S, McCubrey JA, Nicoletti F, Libra M. Computational Modeling of PI3K/AKT and MAPK Signaling Pathways in Melanoma Cancer. *PLoS One* 2016; **11**: e0152104 [PMID: 27015094 DOI: 10.1371/journal.pone.0152104]
- 28 **Tandon M**, Chen Z, Pratap J. Role of Runx2 in crosstalk between Mek/Erk and PI3K/Akt signaling in MCF-10A cells. *J Cell Biochem* 2014; **115**: 2208-2217 [PMID: 25147082 DOI: 10.1002/jcb.24939]
- 29 **Braeuer RR**, Watson IR, Wu CJ, Mobley AK, Kamiya T, Shoshan E, Bar-Eli M. Why is melanoma so metastatic? *Pigment Cell Melanoma Res* 2014; **27**: 19-36 [PMID: 24106873 DOI: 10.1111/pcmr.12172]
- 30 **Fon GT**, Wong WS, Gold RH, Kaiser LR. Skeletal metastases of melanoma: radiographic, scintigraphic, and clinical review. *AJR Am J Roentgenol* 1981; **137**: 103-108 [PMID: 6787858 DOI: 10.2214/ajr.137.1.103]
- 31 **Zekri J**, Marples M, Taylor D, Kandukurti K, McParland L, Brown JE. Complications of bone metastases from malignant melanoma. *J Bone Oncol* 2017; **8**: 13-17 [PMID: 28856087 DOI: 10.1016/j.jbo.2017.08.003]

P- Reviewer: Kiselev SL, Politi LE **S- Editor:** Ji FF **L- Editor:** A
E- Editor: Tan WW





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

