

Full Genome of *Phialocephala scopiformis* DAOMC 229536, a Fungal Endophyte of Spruce Producing the Potent Anti-Insectan Compound Rugulosin

Allison K. Walker,^{a*} Samantha L. Frasz,^a Keith A. Seifert,^b J. David Miller,^a Stephen J. Mondo,^c Kurt LaButti,^c Anna Lipzen,^c Rhyan B. Dockter,^c Megan C. Kennedy,^c Igor V. Grigoriev,^c Joseph W. Spatafora^d

Department of Chemistry, Carleton University, Ottawa, Ontario, Canada^a; Agriculture and Agri-Food Canada, Ottawa, Ontario, Canada^b; U.S. Department of Energy Joint Genome Institute, Walnut Creek, California, USA^c; Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA^d

* Present address: Allison K. Walker, Department of Biology, Acadia University, Wolfville, Nova Scotia, Canada.

We present the full genome of *Phialocephala scopiformis* DAOMC 229536 (Helotiales, Ascomycota), a foliar endophyte of white spruce from eastern Quebec. DAOMC 229536 produces the anti-insectan compound rugulosin, which inhibits a devastating forestry pest, the spruce budworm. This genome will enable fungal genotyping and host-endophyte evolutionary genomics in inoculated trees.

Received 30 December 2015 Accepted 11 January 2016 Published 25 February 2016

Citation Walker AK, Frasz SL, Seifert KA, Miller JD, Mondo SJ, LaButti K, Lipzen A, Dockter RB, Kennedy MC, Grigoriev IV, Spatafora JW. 2016. Full genome of *Phialocephala scopiformis* DAOMC 229536, a fungal endophyte of spruce producing the potent anti-insectan compound rugulosin. *Genome Announc* 4(1):e01768-15. doi:10.1128/genomeA.01768-15.

Copyright © 2016 Walker et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Allison K. Walker, Allison.Walker@acadiau.ca.

Phialocephala scopiformis (Helotiales, Ascomycota) strain DAOMC 229536 was isolated as a foliar endophyte of white spruce (*Picea glauca*) from eastern Quebec. It produces the anti-insectan secondary metabolite rugulosin (1), which inhibits feeding by spruce budworm, a devastating commercial forestry pest (2). Budworm damage results in profound habitat disturbance, reduces sustainable wood supply, and affects the carbon balance of the boreal forest. Inoculated trees are more tolerant of the spruce budworm both in nurseries and under field conditions. The growth rate of the insect pest is reduced, exposing larvae to increased disease and predation by birds and reducing the damage to trees (3–5). In recent years, millions of conifer seedlings in New Brunswick production nurseries were inoculated with this strain to reduce the need for spraying chemical insecticides (6), and beneficial effects have persisted more than a decade postinoculation (7). The genome sequence of *P. scopiformis* will enable studies of fungal genome evolution, fungal genotyping, and host-endophyte evolutionary genomics in the inoculated forests.

The *P. scopiformis* genome was sequenced using the Illumina platform. Genomic DNA was extracted using a modified Joint Genome Institute (JGI) cetyltrimethylammonium bromide (CTAB) and genomic tip (Qiagen) protocol; total RNA was extracted using a modified Trizol method. DNA was sheared and size selected to produce two 300 bp insert size fragments and 4 kb Cre-lox long-mate-pair libraries, and sequenced on Illumina HiSeq in 2 × 150 bp and 2 × 100 bp reads, respectively. Stranded cDNA libraries were generated using the Illumina TruSeq stranded RNA LT kit from mRNA fragmented and reverse transcribed using random hexamers and SSII (Invitrogen) followed by second strand synthesis. The fragmented

cDNA was treated with end-pair, A-tailing, adapter ligation, and 10 cycles of PCR, and sequenced on HiSeq in 2 × 150 bp reads format.

Illumina reads of stranded RNA-seq data were assembled into consensus sequences using Rnnotator (v2.5.6 or later) (8). Genomic data were filtered and assembled with AllPathsLG version R49403 (9) to produce a 48,876,257 bp assembly in 71 scaffolds ($N_{50} = 1.3$ Mb) and 302 contigs ($N_{50} = 433.5$ Kb). Mapped to the genome assembly were 99.42% of assembled consensus RNA sequences. A mitochondrion was assembled separately with AllPathsLG into a single 35.9 kb contig.

The genome of *P. scopiformis* was annotated using the JGI Annotation Pipeline (10) to identify 18,573 genes, over 80% of which are supported by the transcriptome. Genome assemblies and annotation are available via the JGI fungal portal MycoCosm (10).

Nucleotide sequence accession number. The genome of *Phialocephala scopiformis* DAOMC 229536 has been deposited in GenBank under the accession number [LKNI000000000](https://www.ncbi.nlm.nih.gov/nuccore/LKNI000000000).

ACKNOWLEDGMENTS

The assistance of B. Green (Carleton University) and E. Ponomareva, H.D.T. Nguyen, M. Balcerzak, T. Ouellet, T. Barasubiyev, G. Louis-Seize, and the Microbiology Molecular Technologies Laboratory (Agriculture and Agri-Food Canada, Eastern Cereal and Oilseed Research Centre, Ottawa) is gratefully acknowledged.

This work was supported by NSERC CRDPJ 421782-11 to J.D.M. and K.A.S. and by the 1000 Fungal Genomes Project. The work conducted by the U.S. Department of Energy Joint Genome Institute was supported by the Office of Science of the U.S. Department of Energy under contract DE-AC02-05CH11231.

FUNDING INFORMATION

Gouvernement du Canada | Natural Sciences and Engineering Research Council of Canada (NSERC) provided funding to Keith Seifert and J. David Miller under grant number CRDPJ 421782-11. This work was funded by U.S. Department of Energy (DOE) under grant DE-AC02-05CH11231.

REFERENCES

1. Calhoun LA, Findlay JA, Miller JD, Whitney NJ. 1992. Metabolites toxic to spruce budworm from balsam fir needle endophytes. *Mycol Res* 96: 281–286. [http://dx.doi.org/10.1016/S0953-7562\(09\)80939-8](http://dx.doi.org/10.1016/S0953-7562(09)80939-8).
2. Miller JD, Mackenzie S, Foto M, Adams GW, Findlay JA. 2002. Needles of white spruce inoculated with rugulosin-producing endophytes contain rugulosin reducing spruce budworm growth rate. *Mycol Res* 106: 471–479. <http://dx.doi.org/10.1017/S0953756202005671>.
3. Sumarah MW, Miller JD, Adams GW. 2005. Measurement of a rugulosin-producing endophyte in white spruce seedlings. *Mycologia* 97: 770–776. <http://dx.doi.org/10.3852/mycologia.97.4.770>.
4. Miller JD, Sumarah MW, Adams GW. 2008. Effect of a rugulosin-producing endophyte in *Picea glauca* on *Choristoneura fumiferana*. *J Chem Ecol* 34:362–368. <http://dx.doi.org/10.1007/s10886-007-9412-0>.
5. Miller JD, Cherid H, Sumarah MW, Adams GW. 2009. Horizontal transmission of the *Picea glauca* foliar endophyte *Phialocephala scopiformis* CBS 120377. *Fungal Ecol* 2:98–101. <http://dx.doi.org/10.1016/j.funeco.2009.01.002>.
6. Miller JD, Adams GW. 2012. Endophyte enhanced seedlings with increased pest tolerance and methods. Canadian patent 2562175; US20130219569.
7. Frasz SL, Walker AK, Nsiama TK, Adams GW, Miller JD. 2014. Distribution of the foliar fungal endophyte *Phialocephala scopiformis* and its toxin in the crown of a mature white spruce tree as revealed by chemical and qPCR analyses. *Can J For Res* 44:1138–1143. <http://dx.doi.org/10.1139/cjfr-2014-0171>.
8. Martin J, Bruno VM, Fang Z, Meng X, Blow M, Zhang T, Sherlock G, Snyder M, Wang Z. 2010. Rnnotator: an automated *de novo* transcriptome assembly pipeline from stranded RNA-Seq reads. *BMC Genomics* 11:663. <http://dx.doi.org/10.1186/1471-2164-11-663>.
9. Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci USA* 108:1513–1518. <http://dx.doi.org/10.1073/pnas.1017351108>.
10. Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otilar R, Riley R, Salamov A, Zhao X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic Acids Res* 42:D699–D704. <http://dx.doi.org/10.1093/nar/gkt1183>.

Erratum for Walker et al., Full Genome of *Phialocephala scopiformis* DAOMC 229536, a Fungal Endophyte of Spruce Producing the Potent Anti-Insectan Compound Rugulosin

 Allison K. Walker,^a Samantha L. Frasz,^a Keith A. Seifert,^b J. David Miller,^a Stephen J. Mondo,^c Kurt LaButti,^c Anna Lipzen,^c Rhyan B. Dockter,^c Megan C. Kennedy,^c Igor V. Grigoriev,^c Joseph W. Spatafora^d

Department of Chemistry, Carleton University, Ottawa, Ontario, Canada^a; Agriculture and Agri-Food Canada, Ottawa, Ontario, Canada^b; U.S. Department of Energy Joint Genome Institute, Walnut Creek, California, USA^c; Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA^d

Volume 4, no. 1, [e01768-15](#), 2016. Page 2: The final sentence in the Funding Information should read as follows. “This work was funded by U.S. Department of Energy (DOE) under grant DE-AC02-05CH11231.”

Published 26 May 2016

Copyright © 2016 Walker et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).