



Draft genome sequence of plant growth-promoting bacillus altitudinis strain pae4

Nahed Abdel Ghaffar Abdel Aziz Ibrahim, Mohamed Nabil Abdel Mageed
Omar, Ghada A. Abu El Heba, Yvan Moenne-Loccoz, Claire Prigent
Combaret, Daniel Muller

► To cite this version:

Nahed Abdel Ghaffar Abdel Aziz Ibrahim, Mohamed Nabil Abdel Mageed Omar, Ghada A. Abu El Heba, Yvan Moenne-Loccoz, Claire Prigent Combaret, et al.. Draft genome sequence of plant growth-promoting bacillus altitudinis strain pae4. Microbiology Resource Announcements, American Society for Microbiology, 2018, 7 (13), pp.1-2. 10.1128/MRA.00962-18 . hal-02159605

HAL Id: hal-02159605

<https://hal.archives-ouvertes.fr/hal-02159605>

Submitted on 18 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Draft Genome Sequence of Plant Growth-Promoting *Bacillus altitudinis* Strain PAE4

Nahed Abdel Ghaffar Abdel Aziz Ibrahim,^a Mohamed Nabil Abdel Mageed Omar,^b Ghada A. Abu El Heba,^a Yvan Moëgne-Loccoz,^c Claire Prigent-Combaret,^c  Daniel Muller^c

^aAgricultural Genetic Engineering Research institute, ARC, Giza, Egypt

^bSoil, Water and Environment Institute, ARC, Giza, Egypt

^cCNRS, Université de Lyon, Université Claude Bernard Lyon 1, INRA, VetAgro Sup, UMR 5557 Ecologie Microbienne, Villeurbanne, France

ABSTRACT We report here the draft genome of *Bacillus altitudinis* strain PAE4, a thermophilic plant growth-promoting rhizobacterium isolated from the coastal ridge of the Mediterranean Sea in Egypt. Besides heat shock protein genes, several genes encoding phytobeneficial properties were identified.

Bacillus altitudinis is a species encompassing strains that interact often with plants. Most studied strains are well described, as plant growth-promoting rhizobacteria (PGPRs) are known for their phytostimulation abilities (by producing phytohormones or enhancing mineral nutrition [1]), their biocontrol activities (by eliciting plant immune systemic responses or directly inhibiting phytopathogens [2]), and even their ability to help plants in phytoremediation systems (3).

B. altitudinis PAE4 is a bacterial strain isolated from the coastal Mediterranean Ridge in Egypt (4). Strain PAE4 is thermophilic, as it can grow in temperatures up to 70°C (4). As an inoculum, it improved maize and wheat grain yield in field conditions (5). The PII protein (gene *glnK*) that regulates nitrogen metabolism and the phosphotransferase system (PTS) interconnecting carbon and nitrogen metabolisms was shown to play a crucial role in its phytobeneficial activity (5).

Genomic DNA extraction was done from an overnight culture in LB medium (6) at 37°C using a Nucleospin tissue kit (catalog number 740952.50; Macherey-Nagel, Hoerd, France). Genomic DNA was sequenced at MR DNA (Shallowater, TX, USA) using Illumina MiSeq technology, generating a 2 × 300-bp paired-end library. NGen version 14 software (DNAStar, Inc.) was used for trimming sequences (default settings) and for *de novo* assembly (the average size of *B. altitudinis* genomes was used as an assembly parameter). Genome annotation was done with the MicroScope platform version 3.10.0 (7, 8).

A total of 5,940,024 paired-end reads were obtained, giving a coverage depth of 482×. The resulting assemblies generated 28 contigs. The maximum length and N_{50} values of the contigs were 962,863 bp and 877,514 bp, respectively, giving a genome 3,693,314 bp in size with a 36.39% G+C content.

The draft genome harbors 3 rRNAs (1 16S rRNA, 1 23S rRNA, and 1 5S rRNA gene), 70 tRNAs, 2,458 protein-coding genes with functional predictions, and 1,469 genes coding for hypothetical proteins. The *rrs* sequence identity was over 99% with *B. pumilus* and *B. altitudinis*. The Genome-to-Genome Distance Calculator version 2.1 (GGDC2.1) (9) gave a DNA-DNA hybridization estimate of 90.8% with *B. altitudinis* strain 41KF2b^T, showing that PAE4 belongs to the *B. altitudinis* species.

Besides general metabolism genes that were shown to be involved in the phyto-stimulation behavior of PAE4 (such as *glnK* and its downstream regulated genes coding for glutamine synthase, ammonium transport, and phosphoenolpyruvate carboxylase), several genes were present for the PAE4 genome code functions that are more

Received 9 July 2018 Accepted 13 September 2018 Published 4 October 2018

Citation Ibrahim NAGAA, Omar MNAM, El Heba GAA, Moëgne-Loccoz Y, Prigent-Combaret C, Muller D. 2018. Draft genome sequence of plant growth-promoting *Bacillus altitudinis* strain PAE4. Microbiol Resour Announc 7:e00962-18. <https://doi.org/10.1128/MRA.00962-18>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2018 Ibrahim et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Daniel Muller, daniel.muller@univ-lyon1.fr.

specifically involved in plant-microbe interactions (10, 11), such as auxin and cellulose production and xylanase, pectinesterase, amidase, and quercetin dioxygenase activities (12, 13), and that may be important for phytostimulation. In addition, PAE4 possesses a complete type VII secretion system for cell-to-cell molecular communication (14). Six biosynthesis gene clusters for secondary metabolites were predicted with antiSMASH version 4.0.2 (15) in the PAE4 genome. Two clusters are involved in spore coat component biosynthesis. Three other clusters are implicated in the biosynthesis of compounds with anti-phytopathogen activity (fungal or bacterial), i.e., bacilysin, surfactin, and alkylresorsinol (16, 17). Further experiments of plant protection and growth promotion will be performed to clarify the plant-beneficial potential of this bacterium.

Data availability. This draft genome sequence has been deposited at GenBank/ENA under BioProject number [PRJEB26379](https://www.ncbi.nlm.nih.gov/bioproject/PRJEB26379), BioSample number [ERS2486119](https://www.ncbi.nlm.nih.gov/biosample/ERS2486119), and SRA accession number [ERX2730420](https://www.ncbi.nlm.nih.gov/sra/ERX2730420). Contig sequences of the *de novo* assembly have been deposited at EMBL under the accession numbers [OVSL01000001](https://www.ebi.ac.uk/ena/record/OVSL01000001) to [OVSL01000028](https://www.ebi.ac.uk/ena/record/OVSL01000028).

ACKNOWLEDGMENTS

We acknowledge funding from Campus France Imothep and the Academy of Scientific Research and Technology in Egypt. We are grateful to the iBio platform (UMR CNRS 5557 Ecologie Microbienne) for helpful discussions. We also acknowledge the LABGeM and the National Infrastructure “France Génomique” for support within the MicroScope annotation platform.

REFERENCES

- Kumaravel S, Thankappan S, Raghupathi S, Uthandi S. 2018. Draft genome sequence of plant growth-promoting and drought-tolerant *Bacillus altitudinis* FD48, isolated from rice phylloplane. *Genome Announc* 6(9):e00019-18. <https://doi.org/10.1128/genomeA.00019-18>.
- Budiharjo A, Jeong H, Wulandari D, Lee S, Ryu C-M. 2017. Complete genome sequence of *Bacillus altitudinis*, a potential bioprotectant against *Xanthomonas oryzae* pv. *oryzae*, isolated from rice rhizosphere in Java, Indonesia. *Genome Announc* 5(48):e01388-17. <https://doi.org/10.1128/genomeA.01388-17>.
- Djedidi S, Kojima K, Yamaya H, Ohkama-Ohtsu N, Bellingrath-Kimura SD, Watanabe I, Yokoyama T. 2014. Stable cesium uptake and accumulation capacities of five plant species as influenced by bacterial inoculation and cesium distribution in the soil. *J Plant Res* 127:585–597. <https://doi.org/10.1007/s10265-014-0647-x>.
- Ibrahim NAA, El-Zawahry YA, Ahmed G, El-Moslami SH. 2007. Isolation, identification and molecular characterization of halo and thermo tolerant bacteria from coastal ridge of Mediterranean in Egypt. *Egypt J Genet Cytol* 36:341–354.
- Ibrahim NAA, Omar N. 2015. Thermophilic paenibacillus nitrogen fixation bacteria increase cereal crops productivity. *J Am Sci* 11:39–54.
- Bertani G. 1951. Studies on lysogeny I: the mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol* 62:293–300.
- Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S, Lajus A, Pascal G, Scarpelli C, Médigue C. 2006. MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res* 34:53–65. <https://doi.org/10.1093/nar/gkj406>.
- Vallenet D, Calteau A, Cruveiller S, Gachet M, Lajus A, Josso A, Mercier J, Renaux A, Rollin J, Rouy Z, Roche D, Scarpelli C, Médigue C. 2017. MicroScope in 2017: an expanding and evolving integrated resource for community expertise of microbial genomes. *Nucleic Acids Res* 45:D517–D528. <https://doi.org/10.1093/nar/gkw1101>.
- Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <https://doi.org/10.1186/1471-2105-14-60>.
- Bruto M, Prigent-Combaret C, Muller D, Moëgne-Loccoz Y. 2014. Analysis of genes contributing to plant-beneficial functions in plant growth-promoting rhizobacteria and related proteobacteria. *Sci Rep* 4:6261.
- Lemanceau P, Blouin M, Muller D, Moëgne-Loccoz Y. 2017. Let the core microbiota be functional. *Trends Plant Sci* 22:583–595. <https://doi.org/10.1016/j.tplants.2017.04.008>.
- Bowater L, Fairhurst SA, Just VJ, Bornemann S. 2004. *Bacillus subtilis* YxaG is a novel Fe-containing quercetin 2,3-dioxygenase. *FEBS Lett* 557:45–48. [https://doi.org/10.1016/S0014-5793\(03\)01439-X](https://doi.org/10.1016/S0014-5793(03)01439-X).
- Mierziak J, Kostyn K, Kulma A. 2014. Flavonoids as important molecules of plant interactions with the environment. *Molecules* 19:16240. <https://doi.org/10.3390/molecules191016240>.
- Ates LS, Houben ENG, Bitter W. 2016. Type VII secretion: a highly versatile secretion system, p 357–384. In Kudva I, Cornick N, Plummer P, Zhang Q, Nicholson T, Bannantine J, Bellaire B (ed), *Virulence mechanisms of bacterial pathogens*, 5th ed. ASM Press, Washington, DC. <https://doi.org/10.1128/microbiolspec.VMBF-0011-2015>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.
- Chen XH, Koumoutsis A, Scholz R, Eisenreich A, Schneider K, Heine-meyer I, Morgenstern B, Voss B, Hess WR, Reva O, Junge H, Voigt B, Jungblut PR, Vater J, Süßmuth R, Liesegang H, Strittmatter A, Gottschalk G, Borriss R. 2007. Comparative analysis of the complete genome sequence of the plant growth-promoting bacterium *Bacillus amyloliques* FZB42. *Nat Biotechnol* 25:1007. <https://doi.org/10.1038/nbt1325>.
- Cazorla FM, Duckett SB, Bergström ET, Noreen S, Odijk R, Lugtenberg BJJ, Thomas-Oates JE, Bloemberg GV. 2006. Biocontrol of avocado dematophora root rot by antagonistic *Pseudomonas fluorescens* PCL1606 correlates with the production of 2-hexyl 5-propyl resorcinol. *Mol Plant Microbe Interact* 19:418–428. <https://doi.org/10.1094/MPMI-19-0418>.