



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Functional anatomy of the macaque temporo-parieto-frontal connectivity.

This is the peer reviewed version of the following article:

Original

Functional anatomy of the macaque temporo-parieto-frontal connectivity / Borra, Elena; Luppino, Giuseppe. - In: CORTEX. - ISSN 0010-9452. - 97:(2017), pp. 306-326. [10.1016/j.cortex.2016.12.007]

Availability:

This version is available at: 11381/2822867 since: 2021-10-27T12:11:44Z

Publisher:

Masson SpA

Published

DOI:10.1016/j.cortex.2016.12.007

Terms of use:

openAccess

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

(Article begins on next page)

Manuscript Number: CORTEX-D-16-00546R1

Title: Functional anatomy of the macaque temporo-parieto-frontal connectivity

Article Type: Review

Keywords: parietal cortex; temporal cortex; inferior parietal lobule; motor control; sensorimotor transformations

Corresponding Author: Professor Giuseppe Luppino,

Corresponding Author's Institution: University of Parma

First Author: Elena Borra

Order of Authors: Elena Borra; Giuseppe Luppino

Abstract: The primate parietal lobe is primarily dedicated to the processing of sensory information for the guidance of motor behavior, based on the integration of sensory with motor signals (sensorimotor transformations), mediated by specific, strong, and reciprocal connections with the motor cortex. Sensorimotor transformations have been regarded as an automatic process carried out independently from the temporal cortex, which is considered the location where sensory information is used for perceptual processes. However, both human and non-human primate studies have shown interactions between these two regions in different aspects of sensorimotor and cognitive processes. Connectional studies in macaques have provided a detailed description of the possible neural substrate for these interactions. Specifically, temporo-parietal connections almost exclusively involve the inferior parietal lobule (IPL) and display a fine topographic organization, providing the substrate for the role of the macaque IPL in "perception-based" control of motor behavior. Particularly, more rostral IPL areas are involved in motor and cognitive motor functions related to hand action organization and oculomotor control as well as in action and intention understanding, whereas more caudal IPL areas are involved in multisensory integration for the construction of space representations for guiding arm and eye motor behavior. Temporal and IPL-interconnected areas also share connections with specific ventral frontal areas and are thus part of large-scale cortical networks in which the various nodes are linked through "dorsal" temporo-parieto-frontal and "ventral" temporo-frontal pathways. Anatomical and functional studies suggest homologies between human and macaque temporo-parieto-frontal connectivity; they also suggest that higher-order functions of the human IPL could have evolved from the exploitation and adaptation of phylogenetically older neural mechanisms that occur in macaque brains. Thus, connectional data from macaque studies appear essential for understanding human brain mechanisms, even in cases of cognitive abilities undeveloped in other animals, and for interpreting clinical data, including disconnection syndromes.



UNIVERSITA' DEGLI STUDI DI PARMA
DIPARTIMENTO DI NEUROSCIENZE

Via Volturmo 39 - I-43125 PARMA – Italy www.unipr.it

Segreteria amministrativa: ++39 0521 Tel. 903881 Fax 903900

Dear Dr. Catani,

Dear Dr. Ross,

We are herewith submitting a revised version of the manuscript “Functional anatomy of the macaque temporo-parieto-frontal connectivity” for publication in the forthcoming Cortex Special Issue dedicated to the temporal and parietal lobes.

In revising the manuscript, all the comments and suggestions of the reviewers have been carefully taken into account. Accordingly, the title and abstract have been modified in order to make them more congruent with the coverage of the article and several changes have been made throughout the manuscript. Furthermore, a new figure has been added and the manuscript has been revised by a professional company for editing and proofreading.

In the Response to Reviewers section, we describe in details the way in which we have responded to the comments of the referees.

We hope that, after these revisions, the present review article can be considered acceptable for publication in Cortex.

Sincerely yours,

Giuseppe Luppino

Functional anatomy of the macaque temporo-parieto-frontal connectivity

Elena Borra and Giuseppe Luppino

Department of Neuroscience, University of Parma, Parma, Italy

E mail addresses: elena.borra@unipr.it (E. Borra); luppino@unipr.it (G. Luppino)

Corresponding author: Prof. Giuseppe Luppino, Dipartimento di Neuroscienze, Sezione di Fisiologia, Università di Parma, Via Volturno 39, I-43100 Parma, Italy. Phone: +390521903879; Fax: +390521903900; E mail: luppino@unipr.it

We sincerely thank the Reviewers for taking their time in reviewing our manuscript and for their positive comments. Their comments and suggestions have been very helpful for improving the coverage of the topic of the present review and for highlighting some aspects that were underestimated in the previous version.

The following is the detailed description of the way in which the comments of the Reviewers have been addressed.

Reviewer #1:

Comment

The manuscript is well researched, with detailed description of the anatomy and very clear useful figures, however I feel the title and abstract do not fully reflect what is covered in the review. A notable example is that almost as much time is devoted to describing the role of the parieto-frontal connections as does the temporo-parietal connections, although this is hardly mentioned in the title, abstract or introduction.

Response

We agree with the Reviewer. The description of the functional anatomy of the parieto-frontal connectivity was necessary for the functional interpretation of the temporo-parietal connections and their involvement in large-scale cortical networks. However, it is true that the resulting coverage of the manuscript was not fully congruent with the title and abstract. Accordingly, we have modified the title and substantially modified the abstract.

Comment

The section on the human homologues is thin: it focuses mainly on homologous brain regions rather than connectivity per se. An example of this is page 18, line 55, whereby two possible white matter pathways are described as being homologous, yet no detail is given (for example the acronym MLF is not explained or expanded anywhere in the text).

Response

This section has been revised and slightly expanded in order to provide more details on the temporo-parietal connectivity of the human brain and on the possible homologies with the corresponding connectivity of the macaque brain.

Minor Comments

The English does not flow well throughout - it should be rechecked by a native English speaker.

Response

The manuscript has been revised by a professional company

Whilst the figures are generally clear and provide very useful guidance, it might be beneficial to add a figure showing possible homologous connections in humans.

Response

We have added a figure showing examples of large-scale cortical networks of the human brain, possibly evolved from phylogenetically older networks well characterized in the macaque brain.

Reviewer #2:

Comment

The paper's introduction, although thorough in providing a background on monkey connectivity studies, does not justify the central theme of the review. Why temporo-parietal connectivity? What is the gap/need in the literature to justify the review?

Response

In revising the manuscript we tried to declare more clearly the theme and aims of the present review article. This review, based on studies carried out mostly in the last decade, is to our knowledge the first in providing a comprehensive overview of the organization of the macaque temporo-parieto-frontal connectivity from the perspective of its specificity, participation to large-scale cortical networks, and possible homologies with the corresponding connectivity of the human brain.

Comment

This review provides little information on the methodological issues of the studies mentioned, or how the connectivity debate has evolved over the years. There is no critique on how studies limitations have or have not been addressed by other authors. Similarly, the concluding remarks do not offer an insight into future directions for the comparative field, and what important questions remain outstanding. This makes the paper largely read like a textbook.

Response

The aim of this review was to describe the temporo-parieto-frontal connectivity of the macaque brain and to argue that these data can be helpful for understanding the neural mechanisms underlying some high order functions of the human brain. Methodological issues have been addressed in this article (more in detail in the present revised version) only to note that non-invasive functional and connectional approaches used for studying the human brain still have technical limitations in terms of specificity and spatial resolution of the data, compared to the functional and connectional approaches that can be used in non-human primates. The last Section of the manuscript has been revised.

Minor comments:

- The title of the paper suggests a review on macaque temporo-parietal connectivity, when in fact, it also offers a broad review on overall parietal regions and connections associated with the fronto-parietal network and the dorsal and ventral streams. Perhaps a more encompassing title would prevent any confusion.

Response

We agree with the comment of the Reviewer. The Title (and Abstract) has been changed.

- Introduction, at the end of the second paragraph: "recent comparative studies in macaques have questioned its validity for precise in vivo tracing of point-to-point connectivity" - it's a vague statement and does not instruct the reader on what aspects of in-vivo tracing validity this could be referring to. This needs to be expanded a little more, regardless of providing references.

Response

The second paragraph of the Introduction has been revised and expanded in order to provide more details on the limitations of non-invasive MR-based approaches in studying brain connectivity.

- Introduction, the beginning of the 3rd paragraph states: "In this context, old world monkeys, such as macaques, are the primates most closely related to humans...", however it is not clear from the paragraph above, what exact context this is referring to. Also, does this statement refer to brain organisation or genetics? The paragraph needs to provide a background on why old world monkeys are the primates most closely related to humans (e.g. Kaas, 2013).

Response

This part of the Introduction (third-fifth paragraph) has been revised and reorganized also in order to eliminate some redundancies. In particular, we have specified that "Non-human primates such as macaques, are *the experimental model system phylogenetically closest to humans.....*".

- Introduction, 3rd paragraph, examples are given on the usefulness of monkey studies. References are missing after each example; multimodal architectonic studies, cortical maps, electrophysiological studies etc.

Response

We have provided some examples and references for these types of studies.

- The first sentence of the 4th paragraph on how non-human primates studies can be very helpful, is redundant, as this point has already been satisfactorily made.

Response

As mentioned above, this part of the introduction has been revised

- In section 2, " The macaque PPC", only the first paragraph is about the PPC, the second paragraph shifts to a description of parietal areas, which disrupts the flow. The descriptive anatomy should come first, then the subsections.

Response

The content of this Section has been reorganized

- In section 4, "The macaque temporal cortex", the 3rd paragraph briefly compares functional imaging studies in humans. The review thus far hasn't provided any comparisons to human studies, so this sentence seems out of place considering that other subsections don't follow this hierarchy.

Response

We fully agree with the Reviewer. Any reference to human studies in this Section was out of place and has been moved to Section 6.

- Section 6, page 19, "tool use" is discussed more than once in the same paragraph, which perhaps could be merged into one single point.

Response

This Section has been revised and reorganized. Tool use is now mentioned in only one paragraph (page 20).

Functional anatomy of the macaque temporo-parieto-frontal connectivity

Elena Borra and Giuseppe Luppino

Department of Neuroscience, University of Parma, Parma, Italy

E mail addresses: elena.borra@unipr.it (E. Borra); luppino@unipr.it (G. Luppino)

Corresponding author: Prof. Giuseppe Luppino, Dipartimento di Neuroscienze, Sezione di Fisiologia, Università di Parma, Via Volturno 39, I-43100 Parma, Italy. Phone: +390521903879; Fax: +390521903900; E mail: luppino@unipr.it

Abstract

The primate parietal lobe is primarily dedicated to the processing of sensory information for the guidance of motor behavior, based on the integration of sensory with motor signals (sensorimotor transformations), mediated by specific, strong, and reciprocal connections with the motor cortex. Sensorimotor transformations have been regarded as an automatic process carried out independently from the temporal cortex, which is considered the location where sensory information is used for perceptual processes. However, both human and non-human primate studies have shown interactions between these two regions in different aspects of sensorimotor and cognitive processes. Connectional studies in macaques have provided a detailed description of the possible neural substrate for these interactions. Specifically, temporo-parietal connections almost exclusively involve the inferior parietal lobule (IPL) and display a fine topographic organization, providing the substrate for the role of the macaque IPL in “perception-based” control of motor behavior. Particularly, more rostral IPL areas are involved in motor and cognitive motor functions related to hand action organization and oculomotor control as well as in action and intention understanding, whereas more caudal IPL areas are involved in multisensory integration for the construction of space representations for guiding arm and eye motor behavior. Temporal and IPL-interconnected areas also share connections with specific ventral frontal areas and are thus part of large-scale cortical networks in which the various nodes are linked through “dorsal” temporo-parieto-frontal and “ventral” temporo-frontal pathways. Anatomical and functional studies suggest homologies between human and macaque temporo-parieto-frontal connectivity; they also suggest that higher-order functions of the human IPL could have evolved from the exploitation and adaptation of phylogenetically older neural mechanisms that occur in macaque brains. Thus, connectional data from macaque studies appear essential for understanding human brain mechanisms, even in cases of cognitive abilities undeveloped in other animals, and for interpreting clinical data, including disconnection syndromes.

Keywords

parietal cortex; temporal cortex; inferior parietal lobule; motor control; sensorimotor transformations

1. Introduction

In neuroscience today, it is largely accepted that cortical functions are not confined to individual areas but result from the contribution of several, even distant, areas linked together by cortical connections and forming large-scale, functionally specialized networks (see, e.g., Catani et al., 2012). Accordingly, a full understanding of the neural mechanisms underlying a given cortical function also requires an anatomical definition of the underpinning neural substrate in terms of both the number and extent of areas potentially involved in that function and their cortical connectivity.

In recent years, the development of non-invasive functional and connectional imaging techniques has made it possible to address the issue of defining large-scale, functionally specialized networks in the human brain. Indeed, functionally distinct cortical sectors can be identified using functional magnetic resonance imaging (fMRI), and brain connectivity can be investigated using diffusion-weighted MRI (dMRI). Furthermore, intrinsic functional connectivity MRI (fcMRI) has recently emerged as a tool for mapping large-scale networks in the human brain. However, a detailed definition of cortical networks in the human brain is still prevented by several limitations of these techniques. First, fMRI is limited in spatio-temporal resolution and gives indirect information of neuronal activity only at the macroscale level. Further, the areal attribution of activation foci is at best based on probabilistic architectonic maps, which prevent univocal anatomo-functional correlations of experimental data. Second, recent studies of macaques in which well-known cortical pathways have been traced with dMRI have questioned the technique's validity for precise in vivo tracing of point-to-point connectivity (Reveley et al., 2015; Thomas et al., 2014). Indeed, these studies demonstrated that (i) an increase in the sensitivity of different methods seriously decreases specificity, and vice versa (Thomas et al., 2014); (ii) optimal parameters of the tractography algorithm vary according to the traced pathways (Thomas et al., 2014); and (iii) the complex arrangement of white matter fibers residing just under the cortical sheet poses severe challenges for long-range tractography (Reveley et al., 2015). Third, several observations have been made that functional connectivity is related to but distinct from anatomic connectivity, as it could be subserved by polysynaptic as well as monosynaptic anatomic circuits and can be modulated by the task performed by the subjects or several other factors, differently from structural connectivity (Biswal et al., 2010; Buckner, Krienen, & Yeo, 2013).

Non-human primates such as macaques are the experimental model system phylogenetically closest to humans for obtaining highly detailed anatomical and functional information about the neural pathways and mechanisms underlying sensory, motor, and cognitive functions (see, e.g., Passingham, 2009; Sereno & Tootell, 2005). Indeed, based on comparative observations, it is

1 largely agreed that the monkey and human brain share a common plan of anatomical and functional
2 organization of sensory, motor, and associated cortical areas (Caminiti, Innocenti, & Battaglia-
3 Mayer, 2015; Geyer, Matelli, Luppino, & Zilles, 2000; Mantini, Corbetta, Romani, Orban, &
4 Vanduffel, 2013; Orban, Van Essen, & Vanduffel, 2004).

5
6
7 In non-human primates, multimodal architectonic studies can provide reliable cortical maps that
8 can be used as anatomical frames of reference for the univocal attribution of specific functional and
9 connectional data, obtained in the same or different subjects, to specific cortical areas. Furthermore,
10 electrophysiological studies can provide information on the functional properties of cortical areas,
11 even at the level of individual neuronal activity. Finally, neural tracers, based on their anterograde
12 and/or retrograde axonal transport, can provide information on the afferent and efferent connections
13 of a given cortical sector at the cellular level and, via their laminar distribution, the hierarchical
14 relationships between connected areas (see Felleman & Van Essen, 1991). These connectional data
15 are crucial for assessing whether areas with common or similar functional properties belong to the
16 same or different networks and to make hypotheses on the potential information flow through a
17 given network. Thus, non-human primates provide invaluable data for building more
18 comprehensive, anatomically plausible models of cortical functions, which could in turn be used for
19 guiding anatomical and functional investigations in humans and supporting research-based clinical
20 practices. For example, based on cyto-, myelo-, chemo-, and receptor-architectonic observations,
21 the macaque inferior parietal convexity cortex has been subdivided into four different areas (Geyer,
22 Luppino, Ekamp, & Zilles, 2005; Gregoriou, Borra, Matelli, & Luppino, 2006), which later studies
23 have shown to be functionally and connectionally distinct (Rozzi et al., 2006; Rozzi, Ferrari,
24 Bonini, Rizzolatti, & Fogassi, 2008). These data have subsequently been used for suggesting
25 homologies with the corresponding region of the human brain (Caspers et al., 2006, 2011, 2013).

26
27
28 One potential challenge to the generalization of non-human primate data to the human brain is
29 that approximately 30 million years of independent evolution have resulted in significant
30 differences between the brains of monkeys and humans; for example, compared to monkeys, a
31 disproportionate enlargement of some cortical regions in the human brain is considered the neural
32 basis for the outstanding cognitive capabilities of humans (see, e.g., Chaplin, Yu, Soares, Gattass, &
33 Rosa, 2013; Passingham, 2009; Rilling, Glasser, Jbabdi, Andersson, & Preuss, 2012; Sereno &
34 Tootell, 2005). However, it is largely accepted that similar features present in different members of
35 a phylogenetic radiation can be explained as being retained from a common ancestor (see Kaas,
36 2013), and that evolutionary processes have more commonly involved the modification of existing
37 anatomical structures than the addition of new ones. Thus, novel functions have mostly evolved
38 from the adaptation of existing neural mechanisms (see, e.g., de Waal & Ferrari, 2010; Passingham,
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2009; Rilling et al., 2012). The macaque model could thus be essential for understanding the mechanisms of the human brain, even those cognitive abilities that have not developed in other animals.

The general aim of this review article is to argue that this could also be the case for higher-order cognitive functions resulting from the interplay between the parietal, frontal, and temporal cortex. Toward this end, we will first provide an updated overview of the functional anatomy of macaque temporo-parieto-frontal connectivity, based on data collected mostly in the last decade, showing that these connections are an ensemble of different streams linking specific temporal, parietal, and frontal areas and taking part in large-scale, functionally specialized networks for controlling different aspects of behavior. Then, based on anatomical and functional evidence from human studies, we will consider possible homologies with the corresponding connectivity of the human brain to suggest that in primate evolution, phylogenetically old neural pathways and mechanisms have been retained and, in the human lineage, have been exploited for the emergence of human-specific cognitive capabilities.

2. The macaque posterior parietal cortex (PPC)

The macaque posterior parietal cortex (PPC) is a large cortical region that, as in the human brain, is subdivided by the intraparietal sulcus (IP) into a superior (SPL) and inferior (IPL) parietal lobule. As a whole, the PPC is primarily dedicated to the analysis of diverse, high-order aspects of sensory information, with special emphasis on visual information. One main aim of this type of processing is to use sensory information for the guidance of motor behavior, based on the integration of sensory and motor signals mediated by strong and reciprocal connections with the motor (primary motor and premotor) cortex (see, e.g., Caminiti et al., 2015; Rizzolatti, Luppino, & Matelli, 1998). This integration is the basis of so-called “sensorimotor transformations,” a neural mechanism in which automatically processed sensory information leads to the activation, in the motor cortex, of representations of “potential actions” (see, e.g., Cisek & Kalaska, 2010; Cisek, 2007; Rizzolatti & Luppino, 2001).

For a long time, it was believed that the SPL was exclusively dedicated to the analysis of higher-order aspects of somatosensory information originating from the primary somatosensory cortex, which is located in the postcentral gyrus. In contrast, the IPL was believed to be the main target of the occipito-parietal visual information processing pathway, usually referred to as the “dorsal visual stream,” which is primarily involved in space processing (Ungerleider & Mishkin, 1982). Non-human primate research over the last two decades has led to a radically different view of the

1 organization of the PPC, involving the following major aspects (for reviews, see, e.g., Caminiti et
2 al., 2015; Colby, 1998; Rizzolatti et al., 1998; Rizzolatti, Fogassi, & Gallese, 1997). First, the PPC
3 is formed by a multiplicity of distinct sensorimotor areas (Figure 1), located in both the SPL and the
4 IPL, which are involved in the analysis of specific aspects of sensory information and are related to
5 the control of specific effectors. Second, a majority of these areas are involved in the analysis of
6 visual information, alone or in combination with somatosensory information. Third, visually related
7 PPC areas, fed by the dorsal visual stream, are located not only in the IPL but also in the SPL.
8 Finally, PPC areas of the IPL display robust, topographically organized connections with areas of
9 the temporal cortex.
10
11
12
13
14
15
16
17

18 **3. Parieto-frontal connectivity and general subdivision of the PPC**

19
20
21

22 Anatomical studies have shown that almost the entire extent of the PPC is linked by strong and
23 reciprocal connections to the motor cortex. Specifically, from early connectional studies of parieto-
24 frontal connectivity (e.g., Pandya & Kuypers, 1969; Petrides & Pandya, 1984), it has become clear
25 that these connections are topographically organized, with the SPL and IPL being mostly connected
26 with more dorsal (PMd) and more ventral (PMv) parts of the premotor cortex. In later studies, the
27 issue of the topographic organization of parieto-frontal connectivity was re-addressed in light of
28 data showing that both the PPC and the motor cortex are formed by several anatomically and
29 functionally independent areas (see, e.g., Rizzolatti & Luppino, 2001; Rizzolatti et al., 1998).
30
31
32
33
34
35

36 One major general finding of these studies was that parieto-frontal connections are organized in
37 a rather specific way, and that each parietal area appears to have a privileged target in the frontal
38 cortex. Based on these privileged connections, it is thus possible to identify a series of largely
39 segregated parieto-frontal circuits linking specific PPC areas with specific motor areas. Functional
40 evidence indicates that parietal and frontal areas forming a given circuit tend to have similar
41 functional properties and appear to be jointly involved in specific aspects of sensorimotor
42 transformations, based on the full integration of sensory and motor signals, at both the parietal and
43 frontal level. Accordingly, this anatomical organization of parieto-frontal connections represents the
44 substrate for a parallel processing of different aspects of sensorimotor transformations. It has been
45 proposed that these parieto-frontal circuits represent the functional units of the cortical motor
46 system (Rizzolatti et al., 1998).
47
48
49
50
51
52
53
54
55

56 Based on their general functional properties and patterns of frontal connectivity, the various PPC
57 areas can be grouped into a rostral and a caudal domain, in both the SPL and the IPL. In the SPL, a
58 more rostral domain includes two mostly somatomotor areas, PE and PEa. One major feature of the
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

neural activity in these areas is the integration of multi-joint sensory and motor signals related to arm/hand position and movement direction, which can be used for encoding egocentric representations of reaching, which are essential for guiding arm movements in space (e.g., Lacquaniti & Caminiti, 1998; Lacquaniti, Guigon, Bianchi, Ferraina, & Caminiti, 1995). Frontal connections of these areas (Figure 2A) involve the primary motor area F1 and the rostrally adjacent corticospinal areas F2 and F3 (SMA) located in the PMd and medial premotor cortex, respectively (Bakola, Passarelli, Gamberini, Fattori, & Galletti, 2013; Caminiti, Zeger, Johnson, Urbano, & Georgopoulos, 1985; Marconi et al., 2001; Matelli, Govoni, Galletti, Kutz, & Luppino, 1998; Tanné-Gariépy, Rouiller, & Boussaoud, 2002).

A more caudal SPL domain consists of a group of several visuomotor areas, such as V6A, PGm, PEc, and MIP, which are differentially connected (Figure 2B) with the PMd areas F7 and F2 (Bakola, Gamberini, Passarelli, Fattori, & Galletti, 2010; Gamberini et al., 2009; Johnson, Ferraina, Bianchi, & Caminiti, 1996; Johnson, Ferraina, & Caminiti, 1993; Marconi et al., 2001; Matelli et al., 1998; Tanné-Gariépy et al., 2002). Two of these areas—V6A and PGm—are also connected to the dorsal part of prefrontal area 46 (46d; Gamberini et al., 2009; Leichnetz, 2001; Saleem, Miller, & Price, 2014), a region originally considered to be involved in encoding visuospatial information in working memory (see, e.g., Levy & Goldman-Rakic, 2000). These caudal SPL areas, in general, combine visuospatial information with gaze- and arm-related signals for visuomotor control of arm-reaching movements (for reviews, see, e.g., Caminiti et al., 2015; Fattori, Breveglieri, Bosco, Gamberini, & Galletti, 2015). Visual information sent to these areas mostly originates from the extrastriate motion-sensitive area V6 (Galletti et al., 2001), where the representation of the visual periphery is relatively emphasized (Galletti, Fattori, Battaglini, Shipp, & Zeki, 1996). Altogether, these data have supported the notion that caudal SPL–PMd connectivity is primarily involved in visuomotor control of reaching behavior. The finding that reaching neurons in area V6A can also be modulated by grasping movements has suggested its broader role in visuomotor control of reaching-to-grasp actions. Parieto-frontal connectivity of the areas of both the rostral and caudal SPL domains is supported by the first branch of the superior longitudinal fasciculus (SLF I; Schmahmann et al., 2007).

In the IPL, a more rostral domain includes the somato- and visuo-motor areas PF, PFG, and AIP, which are devoted to the control of goal-directed hand and mouth actions. These areas are tightly connected (Figure 2C) to the various subdivisions of the PMv area F5 (F5p, F5c, and F5a/44) and to the ventrolateral prefrontal (VLPF) areas ventral 46 (46v) and 12r (Borra et al., 2008; Cavada & Goldman-Rakic, 1989b; Petrides & Pandya, 1984; Rozzi et al., 2006). Specifically, areas AIP and PFG are two hand-related areas and are targets of dorsal visual stream information originating from

1 the motion-sensitive area MST and the caudal intraparietal (CIP) area, in which neurons code 3D
2 objects' visual features (Sakata, Tsutsui, & Taira, 2005). AIP and PFG, together with F5, play a
3 crucial role in selecting and organizing object-oriented hand actions (for a review, see Rizzolatti,
4 Cattaneo, Fabbri-Destro, & Rozzi, 2014). Specifically, different lines of evidence have clearly
5 shown that AIP and F5 are jointly involved in visuomotor transformations for grasping; that is, the
6 selection of appropriate patterns of distal hand configurations based on visual coding of an object's
7 physical properties. Furthermore, PFG and F5 appear to be jointly involved in selecting hand
8 actions according to the goal or motor intention, reflecting sequential action organization. Finally,
9 areas PFG and AIP, as well as F5, host visuomotor hand-related "mirror" neurons that activate
10 during action observation (Fogassi et al., 2005; Maeda, Ishida, Nakajima, Inase, & Murata, 2015;
11 Rizzolatti et al., 2014) and represent the parietal node of the so-called "mirror system," in which
12 observed actions are mapped onto the corresponding motor representations and which is deemed to
13 be the basis of action and intention understanding (Rizzolatti & Craighero, 2004). A further IPL
14 area that could be included within this domain is area LIP, located more caudally in the IP. Indeed,
15 area LIP is an oculomotor area, selectively connected (Figure 2C) with the frontal eye field (FEF)
16 and the adjacent oculomotor caudal prefrontal areas 45B, 8r, and caudal 46 (Andersen, Asanuma,
17 Essick, & Siegel, 1990; Blatt, Andersen, & Stoner, 1990; Cavada & Goldman-Rakic, 1989b), which
18 plays a crucial role in visuomotor transformations for controlling oculomotor behavior (e.g., Blatt et
19 al., 1990). The connectivity of these more rostral IPL (including LIP) areas with PMv, prearcuate
20 oculomotor, and VLPF areas appears to be supported mostly by the third branch of the superior
21 longitudinal fasciculus (SLF III; Schmahmann et al., 2007).

22 Finally, a more caudal domain in the IPL includes two areas, PG and Opt, which are targets of
23 strong projections from the motion-sensitive area MST (Rozzi et al., 2006) and correspond to the
24 cortical sector originally defined as area 7a by Vogt and Vogt (1919). Differently from most other
25 PPC areas, PG and Opt, through the second branch of the superior longitudinal fasciculus (SLF II;
26 Schmahmann et al., 2007), display relatively widespread frontal connectivity (Figure 2D), as they
27 are connected to different parts of both PMd and PMv and 46d and 46v and are strongly and
28 reciprocally connected to rostral IPL and caudal SPL visuomotor areas (Cavada & Goldman-Rakic,
29 1989a, 1989b; Petrides & Pandya, 1984; Rozzi et al., 2006). Neurons in this parietal sector have
30 large visual receptive fields, modulated by the orbital position of the eye or by the position of the
31 head, and respond to different patterns of optic flow stimuli (Merchant, Battaglia-Mayer, &
32 Georgopoulos, 2001). Most neurons also display movement-related activity, mostly related to arm-
33 reaching in the rostral area PG and to saccades in the caudal area Opt (Battaglia-Mayer, Mascarò,
34 Brunamonti, & Caminiti, 2005; Hyvärinen, 1981; MacKay, 1992; Mountcastle, Lynch,
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Georgopoulos, Sakata, & Acuna, 1975). However, there are also neurons encoding both eye- and
2 arm-directional visuomotor signals related to the contralateral space (Battaglia-Mayer et al., 2005),
3 suggesting a role of this sector in representing action space for intended eye and/or arm movements
4 toward the contralateral space. Furthermore, this sector hosts neurons that encode the location of the
5 most salient stimulus and are involved in redirecting attention to stimuli appearing at novel,
6 unattended locations (Constantinidis & Steinmetz, 2001a, 2001b, 2005; Mountcastle, Andersen, &
7 Motter, 1981). Finally, visually responsive neurons that appear to support an object-referenced
8 representation of space also appear in this sector (Chafee, Averbeck, & Crowe, 2007) and can be
9 modulated by world-referenced information (Snyder, Grieve, Brotchie, & Andersen, 1998).
10 Altogether, these data suggest that this caudal IPL sector, based on multisensory integration,
11 contributes to the construction of representations of surrounding space and to spatial attentional
12 processes for guiding the motor behavior of arms and/or eyes.
13
14
15
16
17
18
19
20

21 The notion that the PPC, or most of it, is primarily involved in sensorimotor (mostly visuomotor)
22 transformations was formalized in the 1990s by Milner and Goodale (Milner & Goodale, 1993,
23 1995), who proposed that the dorsal visual stream could actually be considered, as a whole, a
24 “vision for action” pathway. In this view, visuomotor transformations are mostly meant as fast,
25 automatic, and unconscious processes carried out independently from perceptual processes, which
26 are considered the exclusive province of the occipito-temporal “ventral visual stream,” leaving open
27 the question of how and where perceptual and cognitive process modulate or steer voluntary motor
28 behavior.
29
30
31
32
33
34
35

36 A different model of the general organization of visual information processing in the macaque
37 PPC was subsequently proposed by Rizzolatti and Matelli (2003). According to this model, the
38 various parieto-frontal circuits, fed by the dorsal visual stream, can be grouped into two
39 anatomically and functionally distinct main components: the dorso-dorsal stream, formed by
40 circuits linking SPL with PMd areas; and the ventro-dorsal stream, formed by circuits linking IPL
41 with PMv areas and the FEF. This proposal was mostly founded on two sets of connectional data.
42 First, caudal SPL and IPL visuomotor areas are targets of dorsal visual stream information
43 originating mostly from different extrastriate visual areas—that is, areas V6 and MT/MST,
44 respectively. Second, as noted since earlier tract-tracing studies (e.g., Seltzer & Pandya, 1984,
45 1994), the IPL, but not the SPL, is robustly connected with the temporal cortex. More recent studies
46 focused on the cortical connectivity of the various visuomotor parietal areas have described in detail
47 the organization of temporo-parietal connectivity, providing new insights into their potential
48 contribution to motor and cognitive motor function.
49
50
51
52
53
54
55
56
57
58
59

60 **4. The macaque temporal cortex**

1 The macaque temporal cortex can be anatomically subdivided by the superior temporal sulcus
2 (STS) into an inferior and superior temporal gyrus and functionally subdivided into four main
3 domains: a purely visual domain, located in the inferior temporal gyrus (inferotemporal cortex, IT);
4 a multisensory domain, located along the upper bank of the STS; an auditory domain, located in the
5 superior temporal gyrus (STG) and the planum temporale; and a memory-related domain, located in
6 the medial temporal cortex.
7

8
9 The first domain—the IT cortex—includes areas, located at different rostrocaudal levels, which
10 represent the progression in the temporal lobe of the ventral visual stream: the occipito-temporal
11 visual information processing pathways devoted to the analysis of object qualities and identities
12 (Ungerleider & Mishkin, 1982). This pathway comprises a set of bidirectional projections that, from
13 visual areas of the occipital cortex, run in a rostrocaudal direction to the temporal pole through the
14 IT areas TEO and TE (see, e.g., Kravitz, Saleem, Baker, Ungerleider, & Mishkin, 2013). This
15 connectivity is supported by the inferior longitudinal fasciculus (ILF; Schmahmann et al., 2007).
16 Visual information processing along this pathway was originally viewed as a serially and
17 hierarchically organized sequence of processing stages in which simpler features are aggregated to
18 obtain more complex object representations (see, e.g., Kravitz et al., 2013). This view has been
19 challenged by more recent data showing that the ventral visual stream is actually a complex
20 recurrent network supporting the parallel processing of different features used to build up
21 representations of what could be better defined as Real World Entities (RWEs)—that is, objects,
22 plants, animals, and conspecifics (see Orban, Zhu, & Vanduffel, 2014). Specifically, it is possible to
23 identify three parallel substreams running in the ventral bank of the STS (TEa/m) and dorsal and
24 ventral parts of TE that are preferentially involved in the analysis of 3D shape, 2D shape, and
25 material properties, respectively. These substreams converge in the rostral part of the IT, where
26 representations of known RWEs are generated.
27

28
29 Connections of the IT cortex involve both temporal and extratemporal areas (Figure 3A).
30 Specifically, rostral IT areas display connections with the perirhinal cortex in the MTL, which can
31 be used for encoding visual information related to RWEs in long-term memory (see Yoshida, Naya,
32 & Miyashita, 2003) and with the upper bank of the STS (Saleem, Suzuki, Tanaka, & Hashikawa,
33 2000). Extratemporal connections involve the frontal and parietal cortex (Saleem, Kondo, & Price,
34 2008; Seltzer & Pandya, 1989, 1994; Webster, Bachevalier, & Ungerleider, 1994) and run through
35 the extreme capsule/uncinate fasciculus and the ILF, respectively (Schmahmann et al., 2007). In the
36 frontal cortex, IT connections involve orbitofrontal areas 11, 13, and 12o, where visual information
37 can be used for the computation of good identities and subjective values in an abstract
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 representation (see, e.g., Padoa-Schioppa & Cai, 2011). Furthermore, IT areas display robust
2 connections (Gerbella, Belmalih, Borra, Rozzi, & Luppino, 2010; Saleem et al., 2014; Schall,
3 Morel, King, & Bullier, 1995; Stanton, Bruce, & Goldberg, 1995) with caudal prefrontal
4 oculomotor areas (FEF, 45B) and, especially for area TEa/m, with VLPF areas—mostly area 12r—
5 originally considered to be involved in working memory for objects and shapes (Wilson, Scalaidhe,
6 & Goldman-Rakic, 1993) and in conditional learning based on object identity (Passingham, Toni, &
7 Rushworth, 2000). This last connection is generally viewed as the progression in the frontal lobe of
8 the ventral visual stream (Kravitz et al., 2013). Finally, the IT cortex, especially area TEa/m of the
9 lower bank of the STS, is also robustly connected with the IPL, much more to the two intraparietal
10 areas, the oculomotor area LIP, and the hand-related area AIP; and much less to IPL convexity
11 visuomotor areas, except for the caudal-most area Opt (Borra et al., 2008; Cavada & Goldman-
12 Rakic, 1989b; Rozzi et al., 2006; Seltzer & Pandya, 1994; Webster et al., 1994).

13
14
15
16
17
18
19
20
21
22 The second domain corresponds to a relatively large cortical sector, including the rostral two-
23 thirds of the upper bank of the STS, and is commonly referred to as the superior temporal
24 polysensory area (STP; Bruce, Desimone, & Gross, 1981). Area STP is a site of convergence for
25 visual information originating from both more caudal motion-sensitive and inferotemporal areas,
26 auditory information originating from dorsally adjacent auditory areas of the STG, and
27 somatosensory information likely originating from IPL areas (for reviews, see Cusick, 1997;
28 Felleman & Van Essen, 1991). Indeed, most STP neurons are visually responsive, but over half of
29 them also exhibit somatosensory and/or auditory responses (Bruce et al., 1981). It has thus been
30 suggested that this area plays a role in the integration of information within and across modalities,
31 subserving orienting behavior to novel stimuli (Baylis, Rolls, & Leonard, 1987; Bruce et al., 1981).
32 Visually responsive neurons may have complex functional properties and have been extensively
33 studied by Perrett and colleagues (for a review, see, e.g., Carey, Perrett, & Oram, 1997). In
34 particular, STP visual neurons respond to various forms of biological motion, including the
35 direction of gaze and head, mouth, or hand movements; they can also differentiate between self-
36 produced actions and actions made by others, and they appear to code the intentionality of actions
37 as well (Jellema, Baker, Wicker, & Perrett, 2000; Jellema & Perrett, 2003). Altogether, these data
38 suggest a key role of this temporal sector in social cognition. More recent functional imaging and
39 electrophysiological data have shown that biological motion processing involves areas of both the
40 upper and lower bank of the STS. Specifically, the upper bank appears to be more sensitive to
41 motion patterns—that is, to kinematic aspects of the actions—whereas the lower bank appears to be
42 more sensitive to configuration changes—that is, the deformation of body shapes (Jastorff,
43 Popivanov, Vogels, Vanduffel, & Orban, 2012; Orban, Zhu, & Vanduffel, 2014).

1 Intratemporal connections of area STP, in addition to the IT cortex (Saleem et al., 2000) and the
2 STG (Seltzer & Pandya, 1978), involve the MTL (Insausti & Amaral, 2008). Extratemporal
3 connectivity (Figure 3B) involves both the frontal cortex and the IPL, with a distribution mostly
4 complementary to that of the IT cortex (Seltzer & Pandya, 1989, 1994). In the frontal lobe, area
5 STP, mostly the rostral part, is connected to the medial prefrontal and orbitofrontal areas of the so-
6 called “medial prefrontal network” (Saleem et al., 2008) and to the rostral part of the VLPF areas
7 12r and 46v. Furthermore, there are strong connections between nearly the entire extent of the STP
8 and dorsal prearcuate oculomotor areas, rostral dorsolateral prefrontal areas, and caudal VLPF area
9 45A. These frontal connections run mostly through the EC. In the IPL, connections running into the
10 middle longitudinal fasciculus (MdLF; Schmahmann et al., 2007) link the STP mostly with
11 visuomotor areas PFG, PG, and Opt of the convexity cortex (Cavada & Goldman-Rakic, 1989a;
12 Rozzi et al., 2006).

13
14
15
16
17
18
19
20
21
22 The third, more dorsal, domain corresponds to the auditory cortex, defined as the cortical region
23 receiving significant thalamic input from one or more divisions of the medial geniculate complex
24 (Hackett, 2011). It is well known that this domain displays a complex anatomical and functional
25 organization and consists of several distinct areas (for reviews, see Hackett, 2011; Kaas & Hackett,
26 2000). Based on common anatomical and functional features, these areas are usually grouped into
27 three regions. The primary areas are located in the planum temporale and are grouped into a
28 centrally located “core” region, whereas the secondary areas are assigned to “belt” or “parabelt”
29 regions, which are located around the core and extend into the STG. Auditory information from the
30 core region progresses serially across regions along the medio-lateral axis, i.e., from core to belt, to
31 the parabelt. Furthermore, the rostral parabelt (RPB) largely connects with the rostral belt areas,
32 whereas the caudal parabelt (CPB) largely connects with the caudal belt areas. The rostrocaudal
33 topography of the connections of the auditory areas represents the substrate for a rostrocaudal
34 functional specialization of the parabelt region. Specifically, there is evidence that neurons located
35 more rostrally in the auditory cortex are preferentially tuned for the type of auditory stimuli,
36 whereas neurons located more caudally are preferentially tuned for the spatial localization of
37 auditory stimuli. These data represent the functional basis for the organization of auditory
38 information processing into “what” and “where” streams analogous to the cortical visual system.

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53 Intratemporal projections from auditory areas are directed medially, toward the insular and
54 retroinsular cortex; laterally, toward the STP; caudally, toward the caudal temporal area Tpt; and
55 rostrally, ultimately targeting the MTL (Hackett, 2011; Muñoz-López, Insausti, Mohedano-
56 Moriano, Mishkin, & Saunders, 2015). Extratemporal projections from the auditory cortex (Figure
57 3C) originate from the belt and parabelt areas and, through the EC, mostly target the prefrontal
58
59
60
61
62
63
64
65

1 cortex (Romanski et al., 1999). Specifically, the rostral parabelt is mostly connected to orbitofrontal
2 areas 11 and 12o, to the frontal pole (area 10), and to VLPF areas 45A and 12l. In contrast, the
3 caudal parabelt is mostly connected to dorsal prearcuate oculomotor areas, rostral area 46d, and area
4 45A. There are few if any projections from the belt and parabelt areas to the PPC. However, the
5 auditory-related area Tpt, located just caudal to the belt, displays relatively robust projections to the
6 IPL that appear to selectively involve area PG (Rozzi et al., 2006).
7
8
9

10 The fourth domain is represented by a set of areas located in the MTL, associated with the limbic
11 system, which play a key role in the formation and consolidation of conscious or declarative
12 memories based on convergent projections from sensory, polymodal, and reward-related cortical
13 areas. Recent functional data have shown that sectors of the parahippocampal cortex are
14 preferentially active for images of scenes (Kornblith, Cheng, Ohayon, & Tsao, 2013). It has been
15 proposed that these sectors are involved in a distinct multisynaptic visual information processing
16 pathway, stemming from area V4v and ultimately targeting the hippocampus and the temporal pole,
17 which could play an important role in spatial memory and navigation (see Kravitz, Saleem, Baker,
18 & Mishkin, 2011). The parahippocampal cortex is also a target of projections from the posterior
19 cingulate cortex and from the IPL area Opt (Cavada & Goldmann-Rakic, 1989a; Rozzi et al., 2006),
20 suggesting a contribution of visuospatial dorsal visual stream information to this function.
21
22
23
24
25
26
27
28
29
30

31 **5. Temporo-parietal connections and large-scale cortical networks for motor and cognitive** 32 **motor functions** 33 34 35 36 37

38 Data so far reviewed show that there is robust connectivity in the macaque brain between the
39 PPC and the temporal lobe that almost exclusively involves the IPL. Furthermore, studies aiming to
40 describe the cortical connectivity of individual anatomically and functionally characterized IPL
41 areas (Cavada & Goldmann-Rakic, 1989a; Blatt et al., 1990; Rozzi et al., 2006; Borra et al., 2008)
42 have provided clear evidence for a topographic organization of IPL–temporal connectivity (Figure
43 4). Specifically, the two main intraparietal areas, AIP and LIP, are preferentially connected with
44 areas of the IT cortex, mostly area TEa/m of the lower bank of the STS; whereas visuomotor IPL
45 convexity areas PFG, PG, and Opt are preferentially connected with the multisensory area STP.
46
47
48
49
50
51
52
53 Furthermore, area PG is also a target of the auditory-related area Tpt.

54 Altogether, these data have strongly contributed to further supporting the notion of a dual-stream
55 organization of visuomotor processing in the macaque PPC, in which SPL and IPL areas take part
56 in a dorso-dorsal and ventro-dorsal stream, respectively. In this context, the dorso-dorsal stream is a
57 relatively “short” visuomotor pathway that, in a few steps from the visual extrastriate area V6
58
59
60
61
62
63
64
65

1 through the caudal SPL, conveys visual information with an emphasis on the visual periphery to
2 arm-related fields of the PMd. In this pathway, visual information is integrated with gaze and arm-
3 related somatosensory and motor signals for relatively fast feed-forward guidance and online
4 monitoring of reach-to-grasp actions (Fattori et al., 2015). In contrast, the macaque IPL is a region
5 where both dorsal and ventral visual stream information and higher-order multisensory information
6 from the STP is integrated with information on motor programs. This type of integration could
7 provide the neural substrate for the role of the ventro-dorsal stream in what could be defined as a
8 “perception-based” control of motor behavior (Rizzolatti & Matelli, 2003; see also Fogassi &
9 Luppino, 2005). The rostral IPL domain appears to be involved in motor and cognitive motor
10 functions related to the selection and organization of hand and mouth actions (in AIP and PFG) and
11 to oculomotor control (in LIP) and in action and intention understanding. The caudal IPL domain
12 appears to be involved in multisensory integration for the construction of representations of
13 surrounding space for guiding arm and eye motor behavior.
14
15
16
17
18
19
20
21
22

23 In this context, the hand-related area AIP is extensively connected to several inferotemporal
24 areas, including strong connections to a specific sector of area TEa/m, located in its intermediate
25 part (Borra et al., 2008). Functional imaging and electrophysiological studies have provided clear
26 evidence for a role of this inferotemporal sector in 3D-shape coding and perception (Janssen,
27 Vogels, Liu, & Orban, 2001; Uka, Tanaka, Yoshiyama, Kato, & Fujita, 2000) and visual coding of
28 hand-object interactions (Barraclough, Keith, Xiao, Oram, & Perrett, 2009; Nelissen et al., 2011;
29 Perrett et al., 1989). The connections between the AIP and intermediate area TEa/m provide the
30 substrate for the potential interaction and exchange of visual and motor signals between these two
31 areas. Indeed, there is evidence for the synchronization of neural activity in the intermediate TEa/m
32 and AIP during 3D-shape discrimination (Verhoef, Vogels, & Janssen, 2011) as well as for the
33 contribution of AIP neurons to 3D-shape perception (Verhoef, Vogels, & Janssen, 2015). Thus, it is
34 possible that ventral visual stream information about object identity is broadcast from
35 inferotemporal areas directly to the AIP. Accordingly, visuomotor transformations for grasping,
36 carried out by the AIP-F5 parieto-frontal circuit, would rely on information related not only to the
37 visual coding of an object’s physical properties, i.e., “how” an object is made, but also to “what” the
38 object target of the action is. The connectivity between the AIP and inferotemporal cortex also
39 provides a direct link between components of the hand motor system and areas responsible for
40 object recognition—that is, an access of signals related to motor and haptic representations of hand
41 actions to the representations of object identity. Accordingly, these connections could play a crucial
42 role in the neural mechanisms underlying tactile object recognition.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Differently from the AIP, the temporal connections of area PFG mostly involve sectors of the
2 upper bank of the STS, corresponding to area STP. As mentioned in Section 4, this area integrates
3 information within and across modalities (Baylis et al., 1987; Bruce et al., 1981) and hosts visual
4 neurons (see Carey et al., 1997) coding biological motion, differentiating between self-produced
5 actions and actions made by others, and coding the intentionality of actions (Jellema et al., 2000;
6 Jellema & Perrett, 2003). This area was originally considered the major source of visual action
7 information to the parieto-frontal mirror system (Rizzolatti & Craighero, 2004). However,
8 functional imaging data in conscious macaques determined that the observation of object-oriented
9 actions activates cortical sectors located in both area STP and TEa/m (Nelissen et al., 2011).
10 Correlations with connectional data showed that one activated sector of area TEa/m corresponded to
11 the intermediate sector of this area connected to the AIP, whereas an activated sector of area STP
12 corresponded to a sector of this area connected to area PFG. These data suggest that visual action
13 information, encoded in the STS, is forwarded to parietal areas of the mirror system along two
14 distinct functional routes. One route, from the TEa/m to AIP area, could provide visual descriptions
15 of the type and immediate goal of hand actions made by others; the other route, from the STP to
16 PFG area, could be involved in extracting the intention behind the observed motor act (Nelissen et
17 al., 2011).

18 Recent connectional evidence in macaques has pointed out that the functional role of these
19 temporo-parietal connections must be considered in the framework of a large-scale, functionally
20 specialized cortical network. Indeed, connectional studies of VLPF areas have shown that a specific
21 sector of area 12r, located in its intermediate part, is connected to area AIP, the anterior sector of
22 area F5 (F5a), and the intermediate sector of area TEa/m, which, in turn, is connected to area AIP
23 (Borra, Gerbella, Rozzi, & Luppino, 2011). These data suggest that these areas altogether are
24 involved in a large-scale cortical network that provides the potential neural substrate for generating
25 purposeful hand actions (Figure 5A). This network has been designated a “lateral grasping network”
26 (Borra, Gerbella, Rozzi, Tonelli, & Luppino, 2014). In this context, area 12r, together with the
27 adjacent interconnected area 46v, could represent the specific substrate for interfacing perceptual,
28 cognitive, and sensorimotor hand-related processes. Specifically, area 12r is a possible substrate for
29 the integration of the ventral visual stream with sensorimotor hand-related information in the
30 prefrontal cortex. These data, in addition to data on the temporal connectivity of area AIP, provide
31 strong support for the idea that the properties of ventral stream processing may not have originally
32 evolved to play a role in pure perception, but may instead reveal its earlier and more fundamental
33 role in collecting information useful for action selection (Cisek & Kalaska, 2010).

1 Inferotemporal areas also project to the oculomotor intraparietal area LIP. These projections
2 originate from sectors of the TEa/m and inferotemporal convexity cortex located more caudally
3 than those connected to the AIP area (Cavada & Goldman-Rakic, 1989a; Blatt et al., 1990;
4 unpublished observations). Indeed, the LIP hosts neurons showing shape selectivity, encoding non-
5 spatial attributes of the stimuli, and activating during the observation of 2D shapes, thus likely
6 reflecting input from ventral visual stream areas (Serenó & Maunsell, 1998; Toth & Assad, 2002).
7 As for the temporal connectivity of area AIP, the functional role of these connections should be
8 considered in the framework of a large-scale cortical network (Figure 5B). Indeed, the ventral part
9 of the FEF, where small amplitude saccades are represented (Bruce, Goldberg, Bushnell, & Stanton,
10 1985), and even more so the ventral prearcuate area 45B, both robustly connected to the LIP, are
11 also robustly connected to those IT sectors that serve as sources of projections to the LIP (Gerbella
12 et al., 2010; Nelissen et al., 2011). In this network, area 45B would represent a prefrontal site of
13 integration of spatial and non-spatial visual information on RWEs used for guiding oculomotor
14 behavior (Gerbella et al., 2010).
15
16
17
18
19
20
21
22
23
24

25 The two caudal IPL convexity areas PG and Opt are robustly connected to area STP.
26 Furthermore, area PG is a target of projections from the auditory-related area Tpt. As noted in
27 Section 3, since early electrophysiological observations, it has been proposed that area STP may be
28 involved in multisensory integration subserving orienting behavior to novel stimuli (Baylis et al.,
29 1987; Bruce et al., 1981). In accord with this view, Perrett and colleagues have proposed that STP
30 neurons belong to a system that detects unexpected and hence behaviorally relevant sensory events
31 (Hietanen & Perrett, 1993; Mistlin & Perrett, 1990). Indeed, lesions involving area STP produce
32 impairment in orientation toward stimuli in the contralateral space (Luh, Butter, & Buchtel, 1986).
33 Furthermore, gaze- and body-part-direction-sensitive neurons can contribute to spatial attentional
34 processes that provide information about the intentions of others (see Carey et al., 1997).
35 Accordingly, STP input, as well as auditory-spatial information from area Tpt, could substantially
36 contribute to the role of caudal IPL areas PG and Opt in higher-order visuospatial processing. As
37 noted above, both caudal IPL areas and area STP have relatively widespread cortical connectivity,
38 which could provide the substrate for their participation in different cortical networks. In this
39 context, there is evidence that prefrontal area 46d, which was originally considered to be involved
40 in visuospatial processing, is connected to both area STP and PG/Opt (Borra, Gerbella, Giorgetti,
41 Rozzi, & Luppino, 2014; Petrides & Pandya, 1999), thus suggesting that temporo-parietal
42 connections between the caudal IPL and area STP are also part of a large-scale, functionally
43 specialized cortical network involved in spatial and attentional processing (Figure 5C). Caudal IPL
44 areas, especially area Opt, are also connected to parahippocampal areas (Cavada & Goldman-Rakic,
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1989a; Rozzi et al., 2006) of the medial temporal lobe that are involved in a scene-processing visual pathway that ultimately targets the hippocampus. Specifically, it has been proposed that the caudal IPL is at the origin of a parieto-medial temporal pathway that, directly or indirectly through posterior cingulate areas, contributes to the complex spatial processing required for navigating through the environment (Kravitz et al., 2013).

6. Possible homologies between the macaque and human temporo-parieto-frontal connectivity

The possible homologies between the macaque and human PPC have been a matter of debate (Luppino, 2005). In humans, lesions involving the PPC produce a large variety of clinical syndromes, according to their localization in the SPL versus the IPL and in the right versus the left hemisphere. Specifically, lesions centered on the SPL, independently from the side of the lesion, can determine optic ataxia (Perenin & Vighetto, 1988; Ratcliff & Davies-Jones, 1972), which can basically be defined as a disorder of visually guided movements of the arms toward a target. In contrast, lesions of the IPL can determine deficits that vary according to the side of the lesion. Lesions of the right IPL, especially its lower part, can produce neglect (Perenin & Vighetto, 1988; Vallar & Perani, 1986), which mostly manifests as the inability of a patient to process and perceive stimuli on the contralateral side of the body or space. Lesions of the rostral part of the left IPL, in correspondence to the supramarginal gyrus (SMG), can cause ideomotor apraxia, a higher-order motor deficit manifesting as a failure to transform the internal representation of a gesture into the appropriate motor actions. This failure is the foundation for the inability to replicate the same gesture following a verbal command, imitate in the absence of an object, and distinguish well-performed from poorly performed movements (De Renzi & Faglioni, 1999; Heilman, Rothi, & Valenstein, 1982). Indeed, Jeannerod and Jacob (2005) have proposed that, in humans, the dorsal visual stream is not a unitary functional system. Whereas the SPL is mostly involved in visuomotor transformations, the IPL is involved in high-level visuomotor representations that contribute, in the right hemisphere, to the perception of spatial relationships; and in the left hemisphere, to the storage of complex representations of actions. This view was difficult to reconcile with the notion that the macaque PPC is part of a “vision for action” visual stream involved in automatic visuomotor processing. As reviewed above, more recent data on macaques have highlighted a dual organization of the dorsal visual stream in which a dorso-dorsal “vision for action” stream is mostly involved in fast feed-forward guidance and online monitoring of reach-to-grasp actions, and a ventro-dorsal stream is involved in “perception-based” motor control. These data, together with functional and

1 structural data from human studies, have provided converging evidence for a general homology
2 between the macaque and human PPC.

3 Indeed, the human rostral IPL, as in macaques, is involved in the neural mechanisms for
4 selecting and controlling goal-directed hand actions and in action and intention understanding
5 (Caspers, Zilles, Laird, & Eickhoff, 2010; Grosbras, Beaton, & Eickhoff, 2012; Molenberghs, Sale,
6 & Mattingley, 2012; Rizzolatti et al., 2014). Specifically, two rostral IPL sectors appear more
7 specifically engaged during object grasping and manipulation, visuomotor transformations for
8 grasping, and planning of sequential goal-directed hand actions. The first, located in the lateral bank
9 of the IP, is considered the human homolog of the macaque AIP (hAIP; see, e.g., Davare, Kraskov,
10 Rothwell, & Lemon, 2011; Grafton, 2010). The second, located in the rostral SMG, corresponds to
11 an architectonic area—PFt—considered the putative homolog of the macaque area PF/PFG
12 (Caspers et al., 2006, 2010). These same IPL areas also activate during observation of object-
13 oriented actions (e.g., Caspers et al., 2010; Grosbras et al., 2012; Molenberghs et al., 2012) and are
14 considered a parietal node of the human mirror system involved in coding goals and intentions of
15 others' actions (see Rizzolatti et al., 2014).

16 As in macaques, a branch of the SLF (SLFIII) in humans connects the rostral intraparietal sulcus
17 and the SMG with the ventral premotor cortex and the inferior frontal gyrus (Makris et al., 2005;
18 Thiebaut de Schotten, Dell'Acqua, & Valabregue, 2012). Thus the frontal target/origin of the
19 SLFIII includes areas involved in hand action control and action and intention understanding, which
20 can be considered at least in part homologous to the macaque area F5 (Rizzolatti & Arbib, 1998;
21 Rizzolatti et al., 2014). Indeed, dMRI evidence for a linkage between the PMv/area 44 and anterior
22 intraparietal sulcus/SMG (Caspers et al., 2011; Ruschel et al., 2014; Schubotz, Anwander, Knösche,
23 von Cramon, & Tittgemeyer, 2010), an increase in effective connectivity between the human AIP
24 and PMv when grasping small objects (Grol et al., 2007), and a reduction in PMv–M1 interactions
25 during grasping preparation after a virtual lesion of the human AIP (Davare, Rothwell, & Lemon,
26 2010) strongly favors the existence of IPL-PMv circuits for the visuomotor control of hand actions
27 homologous to those within the macaque brain.

28 The human rostral IPL also displays an extensive connectivity with the temporal lobe. This
29 connectivity appears to be mostly supported by a fiber system identified as the posterior segment of
30 the arcuate fasciculus (AF) or the SLF (SLFV), linking the posterior part of the superior (STG),
31 middle (MTG), and inferior (ITG) temporal gyrus with the SMG (Catani, Jones, & Ffytche, 2005;
32 Martino et al., 2013; Wu, Sun, Wang, Wang, & Wang, 2016). Recent dMRI evidence has shown
33 that connectivity between the STG and SMG could also be supported by the middle longitudinal
34 fasciculus (MdLF; Makris, Zhu, Papadimitriou, Mouradian, & Ng, 2016). Note that in all these
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 studies, connectivity of the anterior intraparietal sulcus with the temporal cortex was not explicitly
2 investigated.

3 Functional imaging studies in humans have provided clear evidence for a cortical sector located
4 in the posterior ITG and fusiform gyrus (the lateral occipital complex, or LOC) in which different
5 sites are specifically active during the visual processing of shapes, faces, and actions (e.g., Denys et
6 al., 2004; Kanwisher, McDermott, & Chun, 1997; Malach et al., 1995; see also, Orban et al., 2014).
7 Based on comparative fMRI observations, Denis et al. (2004) proposed that the LOC could be the
8 human homolog of the lower bank of the STS (area TEa/m) and the laterally adjacent IT convexity
9 cortex in macaques.
10

11 Notably, in humans, visuomotor processing for selecting and controlling hand actions carried out
12 in the rostral IPL is influenced by processing taking place in ventral visual stream areas, mostly in
13 the LOC (Bracci & Peelen, 2013; Gallivan, Cant, Goodale, & Flanagan, 2014; Majdandžić et al.,
14 2007; Verhagen, Dijkerman, Grol, & Toni, 2008). Furthermore, there is evidence that visual and
15 haptic coding of objects activates both the hAIP (Grefkes, Weiss, Zilles, & Fink, 2002) and LOC
16 (Amedi, Jacobson, Hendler, Malach, & Zohary, 2002; James et al., 2002; Reed, Shoham, &
17 Halgren, 2004), suggesting the multimodal representation of objects in the human ventral visual
18 stream and interactions between the hAIP and LOC for tactile object recognition (Lacey, Tal,
19 Amedi, & Sathian, 2009; Tal & Amedi, 2009).
20

21 Dorsal to the LOC and in proximity to the posterior part of the STS (pSTS) is a region that
22 activates for visual, auditory, and somatosensory stimulation, suggesting multisensory integration
23 and homologies with the STP area in macaques (Beauchamp, Yasar, Frye, & Ro, 2008).
24 Furthermore, the human STS, especially the caudal part, is responsive to diverse types of biological
25 motion and is considered a key node of the so-called “social brain” (Allison, Puce, & McCarthy,
26 2000; Frith & Frith, 2007). Biological motion processing in this region appears to concern mostly
27 kinematic aspects, whereas in the LOC it appears to concern mostly configuration changes of the
28 observed actions, further supporting the homology of this region with the STP in macaques and,
29 also as in macaques, suggesting a dual-stream processing of action observations (Jastorff, Kourtzi,
30 & Giese, 2009). These sectors of the posterior MTG/STS and LOC are considered the major source
31 of visual action information to the mirror system (Caspers et al., 2010; Grosbras et al., 2012;
32 Molenberghs et al., 2012; Rizzolatti et al., 2014). Combined, these data suggest that within the
33 general framework of the temporo-parietal connectivity of the human rostral IPL, there are
34 components equivalent to the macaque pathways connecting the TEa/m and AIP, and the STP and
35 PFG, respectively.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

However, in the left hemisphere, the SMG, posterior MTG/ITG, and PMv/area 44 are also involved in cognitive abilities unique or almost unique to humans, such as higher-order aspects of organizing object-oriented actions, including tool use (Johnson-Frey, Newman-Norlund, & Grafton, 2005; Peeters et al., 2009; Peeters, Rizzolatti, & Orban, 2013; Rilling et al., 2012) and imitation (Buccino et al., 2004; Caspers et al., 2010; Rizzolatti et al., 2014). These cognitive abilities could have emerged in primate evolution from the exploitation and adaptation of the phylogenetically older neural mechanisms and pathways for hand action organization and action and intention understanding that take place in the lateral grasping network of macaques, resulting in the expansion and differentiation of the homologous regions and pathways of the human brain (Figure 6A; e.g., Hecht et al., 2013; Johnson-Frey et al., 2005; Johnson-Frey, 2004; Orban & Caruana, 2014; Ramayya, Glasser, & Rilling, 2010; Rizzolatti et al., 2014).

Specifically, based on dMRI observations, Hecht et al. (2013) suggested that a stronger and more extensive connectivity of MTG regions involved in the visual coding of actions and ITG regions involved in shape and action coding with the SMG, as well as stronger connectivity between the SMG and PMv/area 44, differentiate the cortical mirror network of humans from macaques and could have contributed to the emergence of the role of this system in imitation and imitation learning (see also Rizzolatti et al., 2014). Furthermore, the differentiation of MTG areas responsible for storing conceptual and semantic information about tools, and of a rostral SMG sector devoted to tool use and their interconnectivity, could have contributed to the emergence of tool use based on the exploitation of the phylogenetically older neural mechanisms that occur in the lateral grasping network (Orban & Caruana, 2014; Peeters et al., 2009, 2013; Ramayya et al., 2010).

In the right hemisphere of human brains, a cortical region that includes the ventral part of the angular gyrus (AG), the SMG, and the adjacent caudal temporal cortex (temporo-parietal junction, TPJ) plays an important role in higher-order aspects of visuospatial and attentional processing (Corbetta, Patel, & Shulman, 2008). Specifically, it has been proposed that in the human brain, there are two main large-scale cortical networks involved in attending to environmental stimuli, designated the dorsal and ventral attention network (Corbetta, Patel, & Shulman, 2008; see also Bartolomeo, Schotten, & Chica, 2012; Corbetta & Shulman, 2011). The dorsal attention network includes oculomotor and visuospatial processing areas of the medial IP, SPL, and frontal oculomotor areas and would be more involved in spatial orienting. Considering that the human homolog of the macaque LIP appears to be located in the medial IP (Grefkes & Fink, 2005; Orban, 2016), this network, to some extent, could represent the human equivalent of the oculomotor parieto-frontal circuitry in macaques, linking area LIP to frontal oculomotor areas. The ventral attention network includes the TPJ/IPL and ventral frontal cortex. This network could play a role in

1 detecting salient and behaviorally relevant stimuli in the environment, especially when unattended.
2 These systems dynamically interact during normal perception to determine where and what to
3 attend to. Connectional evidence from dMRI observations has shown that the parieto-frontal
4 connectivity of the dorsal network is supported by the SLFI, whereas that of the ventral network is
5 supported by the SLFIII and is more developed in the right hemisphere (Thiebaut de Schotten et al.,
6 2011). Furthermore, the TPJ/IPL, through the SLFII, is also connected to the frontal components of
7 the dorsal network. These connections of the human TPJ/IPL with both dorsal and ventral frontal
8 areas closely resemble those of the caudal IPL in macaques (areas PG and Opt). Similar to the
9 caudal IPL in macaques, which displays temporal connectivity with the auditory STG area Tpt,
10 STP, and IT, the TPJ/IPL is also connected with the STG, MTG, and ITG through the posterior
11 segment of the AF and the MdLF (Catani et al., 2005; Makris et al., 2016; Martino et al., 2013; Wu
12 et al., 2016). In the right hemisphere, the STG/MTG, TPJ/IPL, and IFG/PMv are the most frequent
13 anatomical correlates of spatial unilateral neglect (Corbetta & Shulman, 2011; Halligan, Fink,
14 Marshall, & Vallar, 2003), and have been included by Karnath and Rorden (2012) in a so-called
15 “cortical network for spatial neglect” (Figure 6B). Indeed, it has long been proposed that attentional
16 spatial processes that may be disrupted in neglect do not result from the activity of single-brain
17 areas but rather emerge from the interaction of large-scale networks (see, e.g., Bartolomeo,
18 Thiebaut De Schotten, & Doricchi, 2007; Heilman, Bowers, Valenstein, & Watson, 1993; Mesulam,
19 1981). In this context, it has been suggested that the parietal component of the ventral attention
20 network includes evolutionarily new areas, not present in macaques, which could be responsible of
21 higher-order aspects of attentional control, including non-spatial attentional processes considered
22 unique or almost unique to humans (Bartolomeo et al., 2012; Corbetta & Shulman, 2011; Patel et
23 al., 2015). Altogether, these data favor topological and connectional homologies between the human
24 TPJ/IPL and the macaque caudal IPL (areas PG and Opt) and suggest that higher-order supramodal
25 attentional processing of the human right IPL, more detached from action, could have evolved from
26 the exploitation and adaptation of the neural mechanisms underlying spatial processing for eye
27 and/or arm movements that occur in caudal IPL areas in macaques.

51 **7. Concluding remarks**

52 The primary aim of this review was to describe the detailed organization of the temporo-parieto-
53 frontal connectivity of the macaque brain, based on tract-tracing studies of the connectivity of
54 individual anatomically and functionally defined areas. These studies show that such connections
55 link specific temporal areas with specific IPL and ventral frontal areas, which are in turn
56
57
58
59
60

1 interconnected, thus providing the neural substrate for large-scale cortical networks for controlling
2 higher-order aspects of motor behavior and cognitive motor functions, in which the various nodes
3 are linked through “dorsal” temporo-parieto-frontal and “ventral” temporo-frontal pathways. This is
4 the case, for example, in the large-scale cortical network for action organization and action and
5 intention understanding involving the rostral IPL, as well as the large-scale cortical network for
6 visuospatial and attentional processing involving the caudal IPL.
7
8
9

10 This organizational picture appears to have been conserved in primate evolution. Indeed, there is
11 clear evidence for human equivalents of these networks (Caspers et al., 2010; Corbetta & Shulman,
12 2011; Grosbras et al., 2012; Karnath & Rorden, 2012; Molenberghs et al., 2012; Rizzolatti et al.,
13 2014), including both “dorsal” and “ventral” pathways, in which the macaque neural mechanisms
14 for action organization, action and intention understanding, and visuospatial and attentional
15 processing have been conserved and exploited for the emergence of human-specific cognitive
16 capabilities, such as tool use, imitation learning, and non-spatial attentional processes. Accordingly,
17 fine-grained connectional data from macaque studies could be essential for understanding human
18 brain mechanisms, even in cases of cognitive abilities not developed in other animals. Furthermore,
19 these data could be helpful for the interpretation of clinical data, including disconnection
20 syndromes, considering that even more modern dMRI approaches can fail in solving fiber crossings
21 and following bundles within fiber tracts, thus preventing the identification of the specific
22 components forming a given fiber tract.
23
24
25
26
27
28
29
30
31
32
33

34 In this context, some features of human temporo-parieto-frontal connectivity still need to be
35 interpreted in light of the general homologies between the macaque and human brain. This is the
36 case, for example, with the temporal connectivity of the SPL, for which there is evidence from both
37 dMRI and fiber dissection observations, although it still lacks a univocal definition. Specifically, it
38 has been reported (although with some differences across studies) that these connections involve, in
39 the temporal cortex, rostral and caudal regions located in the STG, MTG, and ITG, including the
40 fusiform gyrus and, in the SPL, the caudal part corresponding to Brodmann’s area 7 (Hecht et al.,
41 2013; Kamali, Sair, Radmanesh, & Hasan, 2014; Makris et al., 2009, 2013, 2016; Wang et al.,
42 2013; Wu et al., 2016). These connections appear to run through either the MdLF (Kamali et al.,
43 2014; Makris et al., 2009, 2013, 2016; Wang et al., 2013), a fiber system considered a component of
44 the AF/SLF (Kamali et al., 2014), or an independent bundle (Wu et al., 2016). This connection has
45 been described as evolutionarily new, not present in the macaque brain (e.g., Hecht et al., 2013;
46 Wang et al., 2013), and as a possible substrate for a role of the human SPL in action observation
47 (Abdollahi, Jastorff, & Orban, 2013) or visuo-auditory attentional processing (Hecht et al., 2013;
48 Wang et al., 2013; Makris et al., 2016). Furthermore, Hecht et al. (2013) described a connection
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 between the SPL and inferior frontal areas, not present in the macaque brain; however, this
2 connection has not been observed in other studies. Yet, it should be noted that Brodmann’s area 7 is
3 a large, heterogeneous cortical region that includes not just sectors potentially homologous to the
4 macaque visuomotor area V6A, PEc, and MIP, but also extends medially into the precuneus, which
5 likely hosts sectors homologous to the macaque area PGM and posterior cingulate cortex (see, e.g.,
6 Pitzalis, Fattori, & Galletti, 2015); and laterally into the IP, which hosts the putative homolog of the
7 macaque LIP (see, e.g., Orban, 2016). As mentioned in Section 4, in the macaque, area LIP and the
8 posterior cingulate cortex are provided with connections to the IT and medial temporal areas,
9 respectively. Accordingly, it is possible that the temporo–SPL connectivity described in the human
10 brain at least in part includes equivalents of these connections in the macaque brain. Thus, it is
11 possible that the temporal connections of the putative human homolog of the macaque LIP are part
12 of a cortical network equivalent to the “oculomotor network” described in Section 5 (Figure 5B).
13 Future studies are needed to better define the various, potentially phylogenetically older and newer
14 components of human temporo–SPL connectivity.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

The work from our laboratory has been supported by Ministero dell'Istruzione, dell'Università e della Ricerca (grant number: PRIN 2010, 2010MEFNF7_005) and by the European Commission Grant Cogsystems FP7-250013.

References

- 1
2
3
4 Abdollahi, R. O., Jastorff, J., & Orban, G. A. (2013). Common and segregated processing of
5 observed actions in human SPL. *Cerebral Cortex*, *23*(11), 2734–53.
6 <http://doi.org/10.1093/cercor/bhs264>
7
8
9 Allison, T., Puce, A., & McCarthy, G. (2000). Social perception from visual cues: role of the STS
10 region. *Trends in Cognitive Sciences*, *4*(7), 267–278. <http://doi.org/10.1016/S1364->
11 [6613\(00\)01501-1](http://doi.org/10.1016/S1364-6613(00)01501-1)
12
13
14 Amedi, A., Jacobson, G., Hendler, T., Malach, R., & Zohary, E. (2002). Convergence of visual and
15 tactile shape processing in the human lateral occipital complex. *Cerebral Cortex*, *12*(11),
16 1202–12. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12379608>
17
18
19 Andersen, R. A., Asanuma, C., Essick, G., & Siegel, R. M. (1990). Corticocortical connections of
20 anatomically and physiologically defined subdivisions within the inferior parietal lobule. *The*
21 *Journal of Comparative Neurology*, *296*(1), 65–113. <http://doi.org/10.1002/cne.902960106>
22
23
24 Bakola, S., Gamberini, M., Passarelli, L., Fattori, P., & Galletti, C. (2010). Cortical connections of
25 parietal field PEc in the macaque: Linking vision and somatic sensation for the control of limb
26 action. *Cerebral Cortex*, *20*(11), 2592–2604. <http://doi.org/10.1093/cercor/bhq007>
27
28
29
30 Bakola, S., Passarelli, L., Gamberini, M., Fattori, P., & Galletti, C. (2013). Cortical connectivity
31 suggests a role in limb coordination for macaque area PE of the superior parietal cortex. *The*
32 *Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *33*(15), 6648–
33 58. <http://doi.org/10.1523/JNEUROSCI.4685-12.2013>
34
35
36
37 Barraclough, N. E., Keith, R. H., Xiao, D., Oram, M. W., & Perrett, D. I. (2009). Visual adaptation
38 to goal-directed hand actions. *Journal of Cognitive Neuroscience*, *21*(9), 1806–1820.
39 <http://doi.org/10.1162/jocn.2008.21145>
40
41
42
43 Bartolomeo, P., Thiebaut de Schotten, M., & Chica, A. B. (2012). Brain networks of visuospatial
44 attention and their disruption in visual neglect. *Frontiers in Human Neuroscience*, *6*(May),
45 110. <http://doi.org/10.3389/fnhum.2012.00110>
46
47
48 Battaglia-Mayer, A., Mascaro, M., Brunamonti, E., & Caminiti, R. (2005). The over-representation
49 of contralateral space in parietal cortex: a positive image of directional motor components of
50 neglect? *Cerebral Cortex*, *15*(5), 514–25. <http://doi.org/10.1093/cercor/bhh151>
51
52
53
54 Baylis, G. C., Rolls, E. T., & Leonard, C. M. (1987). Functional subdivisions of the temporal lobe
55 neocortex. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*,
56 *7*(2), 330–342.
57
58
59
60 Beauchamp, M. S., Yasar, N. E., Frye, R. E., & Ro, T. (2008). Touch, sound and vision in human
61
62
63
64
65

superior temporal sulcus. *NeuroImage*, 41(3), 1011–1020.

<http://doi.org/10.1016/j.neuroimage.2008.03.015>

Belmalih, A., Borra, E., Contini, M., Gerbella, M., Rozzi, S., & Luppino, G. (2009). Multimodal architectonic subdivision of the rostral part (area F5) of the macaque ventral premotor cortex. *Journal of Comparative Neurology*, 512(2), 183–217.

Biswal, B. B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S. M., ... Milham, M. P. (2010). Toward discovery science of human brain function. *Proceedings of the National Academy of Sciences of the United States of America*, 107(10), 4734–9.

<http://doi.org/10.1073/pnas.0911855107>

Blatt, G. J., Andersen, R. A., & Stoner, G. R. (1990). Visual receptive field organization and cortico-cortical connections of the lateral intraparietal area (area LIP) in the macaque. *The Journal of Comparative Neurology*, 299(4), 421–45. <http://doi.org/10.1002/cne.902990404>

Borra, E., Belmalih, A., Calzavara, R., Gerbella, M., Murata, A., Rozzi, S., & Luppino, G. (2008). Cortical connections of the macaque anterior intraparietal (AIP) area. *Cerebral Cortex*, 18(5), 1094–1111.

Borra, E., Gerbella, M., Giorgetti, V., Rozzi, S., & Luppino, G. (2014). Connectional heterogeneity of the macaque dorsal prefrontal area 46 (46d) (p. Program No 249.12, 2014 Neuroscience Meeting Plann). Washington, DC: Society for Neuroscience, Online.

Borra, E., Gerbella, M., Rozzi, S., & Luppino, G. (2011). Anatomical evidence for the involvement of the macaque ventrolateral prefrontal area 12r in controlling goal-directed actions. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 31(34), 12351–12363.

Borra, E., Gerbella, M., Rozzi, S., Tonelli, S., & Luppino, G. (2014). Projections to the superior colliculus from inferior parietal, ventral premotor, and ventrolateral prefrontal areas involved in controlling goal-directed hand actions in the macaque. *Cerebral Cortex*, 24(4), 1054–1065. <http://doi.org/10.1093/cercor/bhs392>

Boussaoud, D., Ungerleider, L. G., & Desimone, R. (1990). Pathway for motion analysis: cortical connections of the medial superior temporal and fundus of the superior temporal visual areas in the macaque. *The Journal of Comparative Neurology*, 296(3), 462–495.

<http://doi.org/10.1002/cne.902960311>

Bracci, S., & Peelen, M. V. (2013). Body and object effectors: the organization of object representations in high-level visual cortex reflects body-object interactions. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 33(46), 18247–58.

<http://doi.org/10.1523/JNEUROSCI.1322-13.2013>

- 1
2 Bruce, C. J., Desimone, R., & Gross, C. G. (1981). Visual properties of neurons in a polysensory
3 area in superior temporal sulcus of the macaque. *Journal of Neurophysiology*, *46*(2), 369–84.
4 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6267219>
- 5 Bruce, C. J., Goldberg, M. E., Bushnell, M. C., & Stanton, G. B. (1985). Primate frontal eye fields.
6 II. Physiological and anatomical correlates of electrically evoked eye movements. *Journal of*
7 *Neurophysiology*, *54*(3), 714–34. Retrieved from
8
9 <http://www.ncbi.nlm.nih.gov/pubmed/4045546>
- 10
11
12 Buccino, G., Vogt, S., Ritzl, A., Fink, G. R., Zilles, K., Freund, H.-J. J., & Rizzolatti, G. (2004).
13 Neural circuits underlying imitation learning of hand actions: An event-related fMRI study.
14 *Neuron*, *42*(2), 323–334. [http://doi.org/10.1016/S0896-6273\(04\)00181-3](http://doi.org/10.1016/S0896-6273(04)00181-3)
- 15
16
17
18 Buckner, R. L., Krienen, F. M., & Yeo, B. T. T. (2013). Opportunities and limitations of intrinsic
19 functional connectivity MRI. *Nature Neuroscience*, *16*(7), 832–7.
20
21 <http://doi.org/10.1038/nn.3423>
- 22
23 Caminiti, R., Innocenti, G. M., & Battaglia-Mayer, A. (2015). Organization and evolution of
24 parieto-frontal processing streams in macaque monkeys and humans. *Neuroscience &*
25 *Biobehavioral Reviews*, *56*, 73–96. <http://doi.org/10.1016/j.neubiorev.2015.06.014>
- 26
27
28
29 Caminiti, R., Zeger, S., Johnson, P. B., Urbano, A., & Georgopoulos, a. P. (1985). Corticocortical
30 efferent systems in the monkey: A quantitative spatial analysis of the tangential distribution of
31 cells of origin. *Journal of Comparative Neurology*, *241*(4), 405–419.
32
33 <http://doi.org/10.1002/cne.902410402>
- 34
35
36 Carey, D. P., Perrett, D. I., & Oram, M. W. (1997). Recognizing, understanding and reproducing
37 action. In F. Boller & J. Grafman (Eds.), *Handbook of Neuropsychology*, *volIII* (pp. 111–130).
38
39 Amsterdam: Elsevier.
- 40
41
42 Carmichael, S. T., & Price, J. L. (1994). Architectonic subdivision of the orbital and medial
43 prefrontal cortex in the macaque monkey. *The Journal of Comparative Neurology*, *346*(3),
44 366–402. <http://doi.org/10.1002/cne.903460305>
- 45
46
47 Caspers, S., Eickhoff, S. B., Rick, T., von Kapri, A., Kuhlen, T., Huang, R., ... Zilles, K. (2011).
48 Probabilistic fibre tract analysis of cytoarchitectonically defined human inferior parietal lobule
49 areas reveals similarities to macaques. *NeuroImage*, *58*(2), 362–380.
50
51 <http://doi.org/10.1016/j.neuroimage.2011.06.027>
- 52
53
54 Caspers, S., Geyer, S., Schleicher, A., Mohlberg, H., Amunts, K., & Zilles, K. (2006). The human
55 inferior parietal cortex: Cytoarchitectonic parcellation and interindividual variability.
56
57 *NeuroImage*, *33*(2), 430–448. <http://doi.org/10.1016/j.neuroimage.2006.06.054>
- 58
59
60 Caspers, S., Schleicher, A., Bacha-Trams, M., Palomero-Gallagher, N., Amunts, K., & Zilles, K.

- (2013). Organization of the human inferior parietal lobule based on receptor architectonics. *Cerebral Cortex*, 23(3), 615–28. <http://doi.org/10.1093/cercor/bhs048>
- Caspers, S., Zilles, K., Laird, A. R., & Eickhoff, S. B. (2010). ALE meta-analysis of action observation and imitation in the human brain. *NeuroImage*, 50(3), 1148–1167. <http://doi.org/10.1016/j.neuroimage.2009.12.112>
- Catani, M., Dell'Acqua, F., Bizzi, A., Forkel, S. J., Williams, S. C., Simmons, A., ... Thiebaut de Schotten, M. (2012). Beyond cortical localization in clinico-anatomical correlation. *Cortex*, 48(10), 1262–1287. <http://doi.org/10.1016/j.cortex.2012.07.001>
- Catani, M., Jones, D. K., & Ffytche, D. H. (2005). Perisylvian language networks of the human brain. *Annals of Neurology*, 57(1), 8–16. <http://doi.org/10.1002/ana.20319>
- Cavada, C., & Goldman-Rakic, P. S. (1989a). Posterior parietal cortex in rhesus monkey: I. Parcellation of areas based on distinctive limbic and sensory corticocortical connections. *The Journal of Comparative Neurology*, 287(4), 393–421. <http://doi.org/10.1002/cne.902870402>
- Cavada, C., & Goldman-Rakic, P. S. (1989b). Posterior parietal cortex in rhesus monkey: II. Evidence for segregated corticocortical networks linking sensory and limbic areas with the frontal lobe. *The Journal of Comparative Neurology*, 287(4), 422–445. <http://doi.org/10.1002/cne.902870403>
- Chafee, M. V, Averbeck, B. B., & Crowe, D. A. (2007). Representing spatial relationships in posterior parietal cortex: single neurons code object-referenced position. *Cerebral Cortex*, 17(12), 2914–32. <http://doi.org/10.1093/cercor/bhm017>
- Chaplin, T. a., Yu, H.-H., Soares, J. G. M., Gattass, R., & Rosa, M. G. P. (2013). A Conserved Pattern of Differential Expansion of Cortical Areas in Simian Primates. *Journal of Neuroscience*, 33(38), 15120–15125. <http://doi.org/10.1523/JNEUROSCI.2909-13.2013>
- Cisek, P. (2007). Cortical mechanisms of action selection: the affordance competition hypothesis. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 362(1485). <http://doi.org/10.1098/rstb.2007.2054>
- Cisek, P., & Kalaska, J. F. (2010). Neural mechanisms for interacting with a world full of action choices. *Annual Review of Neuroscience*, 33, 269–298. <http://doi.org/10.1146/annurev.neuro.051508.135409>
- Colby, C. L. (1998). Action-Oriented Spatial Reference Frames in Cortex. *Neuron*, 20(1), 15–24. [http://doi.org/10.1016/S0896-6273\(00\)80429-8](http://doi.org/10.1016/S0896-6273(00)80429-8)
- Constantinidis, C., & Steinmetz, M. A. (2001a). Neuronal responses in area 7a to multiple-stimulus displays: I. neurons encode the location of the salient stimulus. *Cerebral Cortex*, 11(7), 581–91. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11415960>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Constantinidis, C., & Steinmetz, M. A. (2001b). Neuronal responses in area 7a to multiple stimulus displays: II. responses are suppressed at the cued location. *Cerebral Cortex*, *11*(7), 592–7. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11415961>
- Constantinidis, C., & Steinmetz, M. A. (2005). Posterior parietal cortex automatically encodes the location of salient stimuli. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *25*(1), 233–8. <http://doi.org/10.1523/JNEUROSCI.3379-04.2005>
- Corbetta, M., Patel, G., & Shulman, G. L. (2008). The Reorienting System of the Human Brain: From Environment to Theory of Mind. *Neuron*, *58*(3), 306–324. <http://doi.org/10.1016/j.neuron.2008.04.017>
- Corbetta, M., & Shulman, G. L. (2011). Spatial neglect and attention networks. *Annual Review of Neuroscience*, *34*, 569–599. <http://doi.org/10.1146/annurev-neuro-061010-113731>
- Cusick, C. G. (1997). The superior temporal polysensory region in monkeys. In J. Kaas, K. Rockland, & A. Peters (Eds.), *Cerebral cortex Vol. 12* (pp. 435–468). New York: Plenum Press.
- Davare, M., Kraskov, A., Rothwell, J. C., & Lemon, R. N. (2011). Interactions between areas of the cortical grasping network. *Current Opinion in Neurobiology*, *21*(4), 565–570. <http://doi.org/10.1016/j.conb.2011.05.021>
- Davare, M., Rothwell, J. C., & Lemon, R. N. (2010). Causal Connectivity between the Human Anterior Intraparietal Area and Premotor Cortex during Grasp. *Current Biology*, *20*(2), 176–81.
- De Renzi, E., & Faglioni, P. (1999). Apraxia. In D. G. Pizzamiglio (Ed.), *Clinical and experimental neuropsychology*. East Sussex, UK: Psychology Press.
- de Waal, F. B. M., & Ferrari, P. F. (2010). Towards a bottom-up perspective on animal and human cognition. *Trends in Cognitive Sciences*, *14*(5), 201–207. <http://doi.org/10.1016/j.tics.2010.03.003>
- Denys, K., Vanduffel, W., Fize, D., Nelissen, K., Sawamura, H., Georgieva, S., ... Orban, G. A. (2004). Visual activation in prefrontal cortex is stronger in monkeys than in humans. *Journal of Cognitive Neuroscience*, *16*(9), 1505–16. <http://doi.org/10.1162/0898929042568505>
- Fattori, P., Breveglieri, R., Bosco, A., Gamberini, M., & Galletti, C. (2015). Vision for Prehension in the Medial Parietal Cortex. *Cerebral Cortex*, bhv302. <http://doi.org/10.1093/cercor/bhv302>
- Felleman, D. J., & Van Essen, D. C. (1991). Distributed hierarchical processing in the primate cerebral cortex. *Cerebral Cortex*, *1*(1), 1–47. <http://doi.org/10.1093/cercor/1.1.1>
- Fogassi, L., Ferrari, P. F., Gesierich, B., Rozzi, S., Chersi, F., & Rizzolatti, G. (2005). Parietal lobe: from action organization to intention understanding. *Science*, *308*(5722), 662–667.

<http://doi.org/10.1126/science.1106138>

- 1
2 Fogassi, L., & Luppino, G. (2005). Motor functions of the parietal lobe. *Current Opinion in*
3 *Neurobiology*, 15(6), 626–631. <http://doi.org/10.1016/j.conb.2005.10.015>
4
5 Frith, C. D., & Frith, U. (2007). Social Cognition in Humans. *Current Biology*, 17(16), R724–R732.
6 <http://doi.org/10.1016/j.cub.2007.05.068>
7
8 Galletti, C., Fattori, P., Battaglini, P. P., Shipp, S., & Zeki, S. (1996). Functional demarcation of a
9 border between areas V6 and V6A in the superior parietal gyrus of the macaque monkey. *The*
10 *European Journal of Neuroscience*, 8(1), 30–52. Retrieved from
11 <http://www.ncbi.nlm.nih.gov/pubmed/8713448>
12
13 Galletti, C., Gamberini, M., Kutz, D. F., Fattori, P., Luppino, G., & Matelli, M. (2001). The cortical
14 connections of area V6: an occipito-parietal network processing visual information. *The*
15 *European Journal of Neuroscience*, 13(8), 1572–88. Retrieved from
16 <http://www.ncbi.nlm.nih.gov/pubmed/11328351>
17
18 Gallivan, J. P., Cant, J. S., Goodale, M. A., & Flanagan, J. R. (2014). Representation of Object
19 Weight in Human Ventral Visual Cortex. *Current Biology*, 24(16), 1866–1873.
20
21 Gamberini, M., Passarelli, L., Fattori, P., Zucchelli, M., Bakola, S., Luppino, G., & Galletti, C.
22 (2009). Cortical connections of the visuomotor parietooccipital area V6Ad of the macaque
23 monkey. *Journal of Comparative Neurology*, 513(6), 622–642.
24 <http://doi.org/10.1002/cne.21980>
25
26 Gerbella, M., Belmalih, A., Borra, E., Rozzi, S., & Luppino, G. (2007). Multimodal architectonic
27 subdivision of the caudal ventrolateral prefrontal cortex of the macaque monkey. *Brain*
28 *Structure and Function*, 212(3–4), 269–301.
29
30 Gerbella, M., Belmalih, A., Borra, E., Rozzi, S., & Luppino, G. (2010). Cortical Connections of the
31 Macaque Caudal Ventrolateral Prefrontal Areas 45A and 45B. *Cerebral Cortex*, 20(1), 141–
32 168. <http://doi.org/10.1093/cercor/bhp087>
33
34 Geyer, S., Luppino, G., Ekamp, H., & Zilles, K. (2005). The macaque inferior parietal lobule:
35 cytoarchitecture and distribution pattern of serotonin 5-HT1A binding sites. *Anatomy and*
36 *Embryology*, 210(5–6), 353–62. <http://doi.org/10.1007/s00429-005-0026-4>
37
38 Geyer, S., Matelli, M., Luppino, G., & Zilles, K. (2000). Functional neuroanatomy of the primate
39 isocortical motor system. *Anatomy and Embryology*, 202(6), 443–474.
40 <http://doi.org/10.1007/s004290000127>
41
42 Grafton, S. T. (2010). The cognitive neuroscience of prehension: recent developments.
43 *Experimental Brain Research*, 204(4), 475–91. <http://doi.org/10.1007/s00221-010-2315-2>
44
45 Grefkes, C., & Fink, G. R. (2005). The functional organization of the intraparietal sulcus in humans

and monkeys. *Journal of Anatomy*, 207(1), 3–17. <http://doi.org/10.1111/j.1469-7580.2005.00426.x>

- 1
2
3
4 Grefkes, C., Weiss, P. H., Zilles, K., & Fink, G. R. (2002). Crossmodal processing of object
5 features in human anterior intraparietal cortex: An fMRI study implies equivalencies between
6 humans and monkeys. *Neuron*, 35(1), 173–184. [http://doi.org/10.1016/S0896-6273\(02\)00741-](http://doi.org/10.1016/S0896-6273(02)00741-9)
7
8
9
10
11 Gregoriou, G. G., Borra, E., Matelli, M., & Luppino, G. (2006). Architectonic organization of the
12 inferior parietal convexity of the macaque monkey. *Journal of Comparative Neurology*,
13
14 496(3), 422–451.
15
16 Grol, M. J., Majdandzic, J., Stephan, K. E., Verhagen, L., Dijkerman, H. C., Bekkering, H., ...
17
18 Toni, I. (2007). Parieto-frontal connectivity during visually guided grasping. *The Journal of*
19
20 *Neuroscience : The Official Journal of the Society for Neuroscience*, 27(44), 11877–11887.
21
22 <http://doi.org/10.1523/JNEUROSCI.3923-07.2007>
23
24 Grosbras, M.-H., Beaton, S., & Eickhoff, S. B. (2012). Brain regions involved in human movement
25 perception: a quantitative voxel-based meta-analysis. *Human Brain Mapping*, 33(2), 431–54.
26
27 <http://doi.org/10.1002/hbm.21222>
28
29 Hackett, T. A. (2011). Information flow in the auditory cortical network. *Hearing Research*, 271(1–
30
31 2), 133–146. <http://doi.org/10.1016/j.heares.2010.01.011>
32
33 Hackett, T. A., Stepniewska, I., & Kaas, J. H. (1998). Subdivisions of auditory cortex and ipsilateral
34 cortical connections of the parabelt auditory cortex in macaque monkeys. *The Journal of*
35
36 *Comparative Neurology*, 394(4), 475–95. Retrieved from
37
38 <http://www.ncbi.nlm.nih.gov/pubmed/9590556>
39
40 Halligan, P. W., Fink, G. R., Marshall, J. C., & Vallar, G. (2003). Spatial cognition: Evidence from
41
42 visual neglect. *Trends in Cognitive Sciences*, 7(3), 125–133. [http://doi.org/10.1016/S1364-](http://doi.org/10.1016/S1364-6613(03)00032-9)
43
44 6613(03)00032-9
45
46 Hecht, E. E., Gutman, D. a., Preuss, T. M., Sanchez, M. M., Parr, L. a., & Rilling, J. K. (2013).
47
48 Process versus product in social learning: Comparative diffusion tensor imaging of neural
49
50 systems for action execution-observation matching in macaques, chimpanzees, and humans.
51
52 *Cerebral Cortex*, 23(5), 1014–1024. <http://doi.org/10.1093/cercor/bhs097>
53
54 Heilman, K. M., Bowers, D., Valenstein, E., & Watson, R. T. (1993). Disorders of visual attention.
55
56 *Baillière's Clinical Neurology*, 2(2), 389–413. Retrieved from
57
58 <http://www.ncbi.nlm.nih.gov/pubmed/8137006>
59
60 Heilman, K. M., Rothi, L. J., & Valenstein, E. (1982). Two forms of ideomotor apraxia. *Neurology*,
61
62 32(4), 342–6. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7199656>
63
64
65

- 1 Hietanen, J. K., & Perrett, D. I. (1993). Motion sensitive cells in the macaque superior temporal
2 polysensory area. I. Lack of response to the sight of the animal's own limb movement.
3 *Experimental Brain Research*, 93(1), 117–28. Retrieved from
4 <http://www.ncbi.nlm.nih.gov/pubmed/8467881>
5
6
7 Hyvärinen, J. (1981). Regional distribution of functions in parietal association area 7 of the
8 monkey. *Brain Research*, 206(2), 287–303.
9
10 Insausti, R., & Amaral, D. G. (2008). Entorhinal cortex of the monkey: IV. Topographical and
11 laminar organization of cortical afferents. *The Journal of Comparative Neurology*, 509(6),
12 608–41. <http://doi.org/10.1002/cne.21753>
13
14
15
16 James, T. W., Humphrey, G. K., Gati, J. S., Servos, P., Menon, R. S., & Goodale, M. A. (2002).
17 Haptic study of three-dimensional objects activates extrastriate visual areas.
18 *Neuropsychologia*, 40(10), 1706–1714. [http://doi.org/10.1016/S0028-3932\(02\)00017-9](http://doi.org/10.1016/S0028-3932(02)00017-9)
19
20
21 Janssen, P., Vogels, R., Liu, Y., & Orban, G. A. (2001). Macaque inferior temporal neurons are
22 selective for three-dimensional boundaries and surfaces. *The Journal of Neuroscience : The*
23 *Official Journal of the Society for Neuroscience*, 21(23), 9419–29. Retrieved from
24 <http://www.ncbi.nlm.nih.gov/pubmed/11717375>
25
26
27
28
29 Jastorff, J., Kourtzi, Z., & Giese, M. A. (2009). Visual learning shapes the processing of complex
30 movement stimuli in the human brain. *The Journal of Neuroscience: The Official Journal of*
31 *the Society for Neuroscience*, 29(44), 14026–38. [http://doi.org/10.1523/JNEUROSCI.3070-](http://doi.org/10.1523/JNEUROSCI.3070-09.2009)
32
33
34
35
36
37 Jastorff, J., Popivanov, I. D., Vogels, R., Vanduffel, W., & Orban, G. A. (2012). Integration of
38 shape and motion cues in biological motion processing in the monkey STS. *NeuroImage*,
39 60(2), 911–921. <http://doi.org/10.1016/j.neuroimage.2011.12.087>
40
41
42 Jeannerod, M., & Jacob, P. (2005). Visual cognition: A new look at the two-visual systems model.
43 *Neuropsychologia*, 43(2 SPEC. ISS.), 301–312.
44 <http://doi.org/10.1016/j.neuropsychologia.2004.11.016>
45
46
47 Jellema, T., Baker, C. I. I., Wicker, B., & Perrett, D. I. I. (2000). Neural representation for the
48 perception of the intentionality of actions. *Brain and Cognition*, 44(2), 280–302.
49 <http://doi.org/10.1006/brcg.2000.1231>
50
51
52 Jellema, T., & Perrett, D. I. (2003). Perceptual history influences neural responses to face and body
53 postures. *Journal of Cognitive Neuroscience*, 15(7), 961–71.
54 <http://doi.org/10.1162/089892903770007353>
55
56
57
58 Johnson-Frey, S. H. (2004). The neural bases of complex tool use in humans. *Trends in Cognitive*
59 *Sciences*, 8(2), 71–78. <http://doi.org/10.1016/j.tics.2003.12.002>
60
61
62
63
64
65

- 1 Johnson-Frey, S. H., Newman-Norlund, R., & Grafton, S. T. (2005). A distributed left hemisphere
2 network active during planning of everyday tool use skills. *Cerebral Cortex*, *15*(6), 681–695.
3 <http://doi.org/10.1093/cercor/bhh169>
4
- 5 Johnson, P. B., Ferraina, S., Bianchi, L., & Caminiti, R. (1996). Cortical networks for visual
6 reaching: Physiological and anatomical organization of frontal and parietal lobe arm regions.
7 *Cerebral Cortex*, *6*(2), 102–119. <http://doi.org/10.1093/cercor/6.2.102>
8
- 9 Johnson, P. B., Ferraina, S., & Caminiti, R. (1993). Cortical networks for visual reaching.
10 *Experimental Brain Research*, *97*(2), 361–5. Retrieved from
11 <http://www.ncbi.nlm.nih.gov/pubmed/7512050>
12
- 13 Kaas, J. H. (2013). The evolution of brains from early mammals to humans. *Wiley Interdisciplinary*
14 *Reviews: Cognitive Science*, *4*(1), 33–45. <http://doi.org/10.1002/wcs.1206>
15
- 16 Kaas, J. H., & Hackett, T. A. (2000). Subdivisions of auditory cortex and processing streams in
17 primates. *Proceedings of the National Academy of Sciences of the United States of America*,
18 *97*(22), 11793–9. <http://doi.org/10.1073/pnas.97.22.11793>
19
- 20 Kamali, A., Sair, H. I., Radmanesh, A., & Hasan, K. M. (2014). Decoding the superior parietal
21 lobule connections of the superior longitudinal fasciculus/arcuate fasciculus in the human
22 brain. *Neuroscience*, *277*, 577–83. <http://doi.org/10.1016/j.neuroscience.2014.07.035>
23
- 24 Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The fusiform face area: a module in human
25 extrastriate cortex specialized for face perception. *The Journal of Neuroscience : The Official*
26 *Journal of the Society for Neuroscience*, *17*(11), 4302–11. Retrieved from
27 <http://www.ncbi.nlm.nih.gov/pubmed/9151747>
28
- 29 Karnath, H. O., & Rorden, C. (2012). The anatomy of spatial neglect. *Neuropsychologia*, *50*(6),
30 1010–1017. <http://doi.org/10.1016/j.neuropsychologia.2011.06.027>
31
- 32 Kornblith, S., Cheng, X., Ohayon, S., & Tsao, D. Y. Y. (2013). A network for scene processing in
33 the macaque temporal lobe. *Neuron*, *79*(4), 766–781.
34 <http://doi.org/10.1016/j.neuron.2013.06.015>
35
- 36 Kravitz, D. J., Saleem, K. S., Baker, C. I., & Mishkin, M. (2011). A new neural framework for
37 visuospatial processing. *Nature Reviews. Neuroscience*, *12*(4), 217–30.
38 <http://doi.org/10.1038/nrn3008>
39
- 40 Kravitz, D. J., Saleem, K. S., Baker, C. I., Ungerleider, L. G., & Mishkin, M. (2013). The ventral
41 visual pathway: An expanded neural framework for the processing of object quality. *Trends in*
42 *Cognitive Sciences*, *17*(1), 26–49. <http://doi.org/10.1016/j.tics.2012.10.011>
43
- 44 Lacey, S., Tal, N., Amedi, A., & Sathian, K. (2009). A putative model of multisensory object
45 representation. *Brain Topography*, *21*(3–4), 269–74. <http://doi.org/10.1007/s10548-009-0087->
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2 Lacquaniti, F., & Caminiti, R. (1998). Visuo-motor transformations for arm reaching. *The*
3 *European Journal of Neuroscience*, *10*(1), 195–203. Retrieved from
4 <http://www.ncbi.nlm.nih.gov/pubmed/9753127>
5
6
7 Lacquaniti, F., Guigon, E., Bianchi, L., Ferraina, S., & Caminiti, R. (1995). Representing spatial
8 information for limb movement: role of area 5 in the monkey. *Cerebral Cortex*, *5*(5), 391–409.
9 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8547787>
10
11
12 Leichnetz, G. R. (2001). Connections of the medial posterior parietal cortex (area 7m) in the
13 monkey. *The Anatomical Record*, *263*(2), 215–36. Retrieved from
14 <http://www.ncbi.nlm.nih.gov/pubmed/11360237>
15
16
17
18 Levy, R., & Goldman-Rakic, P. S. (2000). Segregation of working memory functions within the
19 dorsolateral prefrontal cortex. *Experimental Brain Research*, *133*(1), 23–32.
20 <http://doi.org/10.1086/425589>
21
22
23
24 Luh, K. E., Butter, C. M., & Buchtel, H. A. (1986). Impairments in orienting to visual stimuli in
25 monkeys following unilateral lesions of the superior sulcal polysensory cortex.
26 *Neuropsychologia*, *24*(4), 461–470. [http://doi.org/10.1016/0028-3932\(86\)90091-6](http://doi.org/10.1016/0028-3932(86)90091-6)
27
28
29 Luppino, G. (2005). Organization of the posterior parietal lobe and of parietofrontal connections. In
30 S. Dehaene, J.-R. Duhamel, M. D. Hauser, & G. Rizzolatti (Eds.), *From Monkey Brain to*
31 *Human Brain: A Fyssen Foundation Symposium* (pp. 235–252). Cambridge, MA: MIT Press,
32 Bradford Books.
33
34
35
36 Luppino, G., Ben Hamed, S., Gamberini, M., Matelli, M., & Galletti, C. (2005). Occipital (V6) and
37 parietal (V6A) areas in the anterior wall of the parieto-occipital sulcus of the macaque: A
38 cytoarchitectonic study. *European Journal of Neuroscience*, *21*(11), 3056–3076.
39 <http://doi.org/10.1111/j.1460-9568.2005.04149.x>
40
41
42
43 MacKay, W. A. (1992). Properties of reach-related neuronal activity in cortical area 7A. *Journal of*
44 *Neurophysiology*, *67*(5), 1335–45. Retrieved from
45 <http://www.ncbi.nlm.nih.gov/pubmed/1597716>
46
47
48
49 Maeda, K., Ishida, H., Nakajima, K., Inase, M., & Murata, A. (2015). Functional properties of
50 parietal hand manipulation-related neurons and mirror neurons responding to vision of own
51 hand action. *Journal of Cognitive Neuroscience*, *27*(3), 560–72.
52 http://doi.org/10.1162/jocn_a_00742
53
54
55
56 Majdandžić, J., Grol, M. J., van Schie, H. T., Verhagen, L., Toni, I., & Bekkering, H. (2007). The
57 role of immediate and final goals in action planning: An fMRI study. *NeuroImage*, *37*(2), 589–
58 598. <http://doi.org/10.1016/j.neuroimage.2007.04.071>
59
60
61
62
63
64
65

- 1 Makris, N., Kennedy, D. N., McInerney, S., Sorensen, A. G., Wang, R., Caviness, V. S., & Pandya,
2 D. N. (2005). Segmentation of subcomponents within the superior longitudinal fascicle in
3 humans: a quantitative, in vivo, DT-MRI study. *Cerebral Cortex*, *15*(6), 854–69.
4 <http://doi.org/10.1093/cercor/bhh186>
5
6
7 Makris, N., Papadimitriou, G. M., Kaiser, J. R., Sorg, S., Kennedy, D. N., & Pandya, D. N. (2009).
8 Delineation of the middle longitudinal fascicle in humans: a quantitative, in vivo, DT-MRI
9 study. *Cerebral Cortex*, *19*(4), 777–85. <http://doi.org/10.1093/cercor/bhn124>
10
11
12 Makris, N., Preti, M. G., Asami, T., Pelavin, P., Campbell, B., Papadimitriou, G. M., ... Kubicki,
13 M. (2013). Human middle longitudinal fascicle: variations in patterns of anatomical
14 connections. *Brain Structure & Function*, *218*(4), 951–68. <http://doi.org/10.1007/s00429-012->
15
16
17
18
19
20 Makris, N., Zhu, A., Papadimitriou, G. M., Mouradian, P., & Ng, I. (2016). Mapping temporo-
21 parietal and temporo-occipital cortico-cortical connections of the human middle longitudinal
22 fascicle in subject-specific , probabilistic , and stereotaxic Talairach spaces. *Brain Imaging and*
23
24
25
26
27
28
29 Malach, R., Reppas, J. B., Benson, R. R., Kwong, K. K., Jiang, H., Kennedy, W. A., ... Tootell, R.
30 B. (1995). Object-related activity revealed by functional magnetic resonance imaging in
31 human occipital cortex. *Proceedings of the National Academy of Sciences of the United States*
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Matelli, M., Luppino, G., & Rizzolatti, G. (1991). Architecture of superior and mesial area 6 and the adjacent cingulate cortex in the macaque monkey. *The Journal of Comparative Neurology*, *311*(4), 445–62. <http://doi.org/10.1002/cne.903110402>
- Merchant, H., Battaglia-Mayer, A., & Georgopoulos, A. P. (2001). Effects of optic flow in motor cortex and area 7a. *Journal of Neurophysiology*, *86*(4), 1937–54. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11600652>
- Mesulam, M. M. (1981). A cortical network for directed attention and unilateral neglect. *Annals of Neurology*, *10*(4), 309–25. <http://doi.org/10.1002/ana.410100402>
- Milner, D., & Goodale, M. A. (1993). Visual pathways to perception and action. *Progress in Brain Research*, *95*, 317–37. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8493342>
- Milner, D., & Goodale, M. A. (1995). *The visual brain in action*. Oxford: Oxford University Press.
- Mistlin, A. J., & Perrett, D. I. (1990). Visual and somatosensory processing in the macaque temporal cortex: the role of “expectation”. *Experimental Brain Research*, *82*(2), 437–50. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2286243>
- Molenberghs, P., Sale, M. V., & Mattingley, J. B. (2012). Is there a critical lesion site for unilateral spatial neglect? A meta-analysis using activation likelihood estimation. *Frontiers in Human Neuroscience*, *6*, 78. <http://doi.org/10.3389/fnhum.2012.00078>
- Mountcastle, V. B., Andersen, R. A., & Motter, B. C. (1981). The influence of attentive fixation upon the excitability of the light-sensitive neurons of the posterior parietal cortex. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *1*(11), 1218–25. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6796657>
- Mountcastle, V. B., Lynch, J. C., Georgopoulos, A., Sakata, H., & Acuna, C. (1975). Posterior parietal association cortex of the monkey: command functions for operations within extrapersonal space. *Journal of Neurophysiology*, *38*(4), 871–908. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/808592>
- Muñoz-López, M., Insausti, R., Mohedano-Moriano, A., Mishkin, M., & Saunders, R. C. (2015). Anatomical pathways for auditory memory II: information from rostral superior temporal gyrus to dorsolateral temporal pole and medial temporal cortex. *Frontiers in Neuroscience*, *9*, 158. <http://doi.org/10.3389/fnins.2015.00158>
- Nelissen, K., Borra, E., Gerbella, M., Rozzi, S., Luppino, G., Vanduffel, W., ... Orban, G. A. (2011). Action observation circuits in the macaque monkey cortex. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *31*(10), 3743–3756.
- Orban, G. a. (2016). Functional definitions of parietal areas in human and non-human primates. *Proceedings. Biological Sciences / The Royal Society*, *283*(1828).

<http://doi.org/10.1098/rspb.2016.0118>

- 1
2 Orban, G. A., & Caruana, F. (2014). The neural basis of human tool use. *Frontiers in Psychology*,
3 5(APR), 1–12. <http://doi.org/10.3389/fpsyg.2014.00310>
4
- 5 Orban, G. a., Van Essen, D., & Vanduffel, W. (2004). Comparative mapping of higher visual areas
6 in monkeys and humans. *Trends in Cognitive Sciences*, 8(7), 315–324.
7
8 <http://doi.org/10.1016/j.tics.2004.05.009>
9
- 10 Orban, G. A., Zhu, Q., & Vanduffel, W. (2014). The transition in the ventral stream from feature to
11 real-world entity representations. *Frontiers in Psychology*, 5(JUL), 1–9.
12
13 <http://doi.org/10.3389/fpsyg.2014.00695>
14
15
- 16 Padoa-Schioppa, C., & Cai, X. (2011). The orbitofrontal cortex and the computation of subjective
17 value: consolidated concepts and new perspectives. *Annals of the New York Academy of*
18 *Sciences*, 1239, 130–7. <http://doi.org/10.1111/j.1749-6632.2011.06262.x>
19
20
- 21 Pandya, D. N., & Kuypers, H. G. J. M. (1969). Cortico-cortical connections in the rhesus monkey.
22 *Brain Research*, 13(1), 13–36. [http://doi.org/10.1016/0006-8993\(69\)90141-3](http://doi.org/10.1016/0006-8993(69)90141-3)
23
24
- 25 Pandya, D. N., & Seltzer, B. (1982). Intrinsic connections and architectonics of posterior parietal
26 cortex in the rhesus monkey. *The Journal of Comparative Neurology*, 204(2), 196–210.
27
28 <http://doi.org/10.1002/cne.902040208>
29
30
- 31 Passingham, R. E. (2009). How good is the macaque monkey model of the human brain? *Current*
32 *Opinion in Neurobiology*, 19(1), 6–11. <http://doi.org/10.1016/j.conb.2009.01.002>
33
34
- 35 Passingham, R. E., Toni, I., & Rushworth, M. F. (2000). Specialisation within the prefrontal cortex:
36 the ventral prefrontal cortex and associative learning. *Experimental Brain Research*, 133(1),
37 103–113. <http://doi.org/10.1007/s002210000405>
38
39
- 40 Patel, G. H., Yang, D., Jamerson, E. C., Snyder, L. H., Corbetta, M., & Ferrera, V. P. (2015).
41 Functional evolution of new and expanded attention networks in humans. *Proceedings of the*
42 *National Academy of Sciences of the United States of America*, 112(30), 9454–9.
43
44 <http://doi.org/10.1073/pnas.1420395112>
45
46
- 47 Peeters, R. R., Rizzolatti, G., & Orban, G. A. (2013). Functional properties of the left parietal tool
48 use region. *NeuroImage*, 78, 83–93. <http://doi.org/10.1016/j.neuroimage.2013.04.023>
49
50
- 51 Peeters, Simone, L., Nelissen, K., Fabbri-Destro, M., Vanduffel, W., Rizzolatti, G., & Orban, G. a.
52 (2009). The representation of tool use in humans and monkeys: common and uniquely human
53 features. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*,
54 29(37), 11523–39. <http://doi.org/10.1523/JNEUROSCI.2040-09.2009>
55
56
57
- 58 Perenin, M., & Vighetto, A. (1988). Optic ataxia: a specific disruption in visuomotor mechanisms.
59 I. Different aspects of the deficit in reaching for objects. *Brain*, 111, 643–74.
60
61
62
63
64
65

- 1 Perrett, D. I., Harries, M. H., Bevan, R., Thomas, S., Benson, P. J., Mistlin, A. J., ... Ortega, J. E.
2 (1989). Frameworks of analysis for the neural representation of animate objects and actions.
3 *The Journal of Experimental Biology*, *146*, 87–113. Retrieved from
4 <http://www.ncbi.nlm.nih.gov/pubmed/2689570>
5
6
7 Petrides, M., & Pandya, D. N. (1984). Projections to the frontal cortex from the posterior parietal
8 region in the rhesus monkey. *The Journal of Comparative Neurology*, *228*(1), 105–116.
9 <http://doi.org/10.1002/cne.902280110>
10
11
12 Petrides, M., & Pandya, D. N. (1999). Dorsolateral prefrontal cortex: comparative cytoarchitectonic
13 analysis in the human and the macaque brain and corticocortical connection patterns. *The*
14 *European Journal of Neuroscience*, *11*(3), 1011–1036. <http://doi.org/10.1046/j.1460->
15 [9568.1999.00518.x](http://doi.org/10.1046/j.1460-9568.1999.00518.x)
16
17
18
19
20 Pitzalis, S., Fattori, P., & Galletti, C. (2015). The human cortical areas V6 and V6A. *Visual*
21 *Neuroscience*, *32*, E007. <http://doi.org/10.1017/S0952523815000048>
22
23
24 Ramayya, A. G., Glasser, M. F., & Rilling, J. K. (2010). A DTI investigation of neural substrates
25 supporting tool use. *Cerebral Cortex*, *20*(3), 507–516. <http://doi.org/10.1093/cercor/bhp141>
26
27
28 Ratcliff, G., & Davies-Jones, G. A. (1972). Defective visual localization in focal brain wounds.
29 *Brain*, *95*(1), 49–60. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/5023090>
30
31
32 Reed, C. L., Shoham, S., & Halgren, E. (2004). Neural substrates of tactile object recognition: an
33 fMRI study. *Human Brain Mapping*, *21*(4), 236–46. <http://doi.org/10.1002/hbm.10162>
34
35
36 Reveley, C., Seth, A. K., Pierpaoli, C., Silva, A. C., Yu, D., Saunders, R. C., ... Ye, F. Q. (2015).
37 Superficial white matter fiber systems impede detection of long-range cortical connections in
38 diffusion MR tractography. *Proceedings of the National Academy of Sciences*, *112*(21),
39 E2820–E2828. <http://doi.org/10.1073/pnas.1418198112>
40
41
42 Rilling, J. K., Glasser, M. F., Jbabdi, S., Andersson, J., & Preuss, T. M. (2012). Continuity,
43 Divergence, and the Evolution of Brain Language Pathways. *Frontiers in Evolutionary*
44 *Neuroscience*, *3*, 11. <http://doi.org/10.3389/fnevo.2011.00011>
45
46
47 Rizzolatti, G., & Arbib, M. A. (1998). Language within our grasp. *Trends in Neurosciences*, *21*(5),
48 188–194. [http://doi.org/10.1016/S0166-2236\(98\)01260-0](http://doi.org/10.1016/S0166-2236(98)01260-0)
49
50
51 Rizzolatti, G., Cattaneo, L., Fabbri-Destro, M., & Rozzi, S. (2014). Cortical mechanisms underlying
52 the organization of goal-directed actions and mirror neuron-based action understanding.
53 *Physiological Reviews*, *94*(2), 655–706. <http://doi.org/10.1152/physrev.00009.2013>
54
55
56 Rizzolatti, G., & Craighero, L. (2004). the Mirror-Neuron System. *Annual Review of Neuroscience*,
57 *27*(1), 169–192. <http://doi.org/10.1146/annurev.neuro.27.070203.144230>
58
59
60 Rizzolatti, G., Fogassi, L., & Gallese, V. (1997). Parietal cortex: From sight to action. *Current*
61

Opinion in Neurobiology, 7(4), 562–567. [http://doi.org/10.1016/S0959-4388\(97\)80037-2](http://doi.org/10.1016/S0959-4388(97)80037-2)

1 Rizzolatti, G., & Luppino, G. (2001). The cortical motor system. *Neuron*, 31(6), 889–901.

2 [http://doi.org/10.1016/S0896-6273\(01\)00423-8](http://doi.org/10.1016/S0896-6273(01)00423-8)

3 Rizzolatti, G., Luppino, G., & Matelli, M. (1998). The organization of the cortical motor system:

4 New concepts. *Electroencephalography and Clinical Neurophysiology*, 106(4), 283–296.

5 [http://doi.org/10.1016/S0013-4694\(98\)00022-4](http://doi.org/10.1016/S0013-4694(98)00022-4)

6 Rizzolatti, G., & Matelli, M. (2003). Two different streams form the dorsal visual system: Anatomy

7 and functions. *Experimental Brain Research*, 153(2), 146–157. <http://doi.org/10.1007/s00221->

8 003-1588-0

9 Rizzolatti, G., & Sinigaglia C. (2016). The mirror mechanism: a basic principle of brain function.

10 *Nature Review Neuroscience*, 17(12), 757–765. doi: 10.1038/nrn.2016.135.

11 Romanski, L. M., Tian, B., Fritz, J., Mishkin, M., Goldman-Rakic, P. S., & Rauschecker, J. P.

12 (1999). Dual streams of auditory afferents target multiple domains in the primate prefrontal

13 cortex. *Nature Neuroscience*, 2(12), 1131–1136. <http://doi.org/10.1038/16056>

14 Rozzi, S., Calzavara, R., Belmalih, A., Borra, E., Gregoriou, G. G., Matelli, M., & Luppino, G.

15 (2006). Cortical connections of the inferior parietal cortical convexity of the macaque monkey.

16 *Cerebral Cortex*, 16(10), 1389–1417.

17 Rozzi, S., Ferrari, P. F., Bonini, L., Rizzolatti, G., & Fogassi, L. (2008). Functional organization of

18 inferior parietal lobule convexity in the macaque monkey: Electrophysiological

19 characterization of motor, sensory and mirror responses and their correlation with

20 cytoarchitectonic areas. *European Journal of Neuroscience*, 28(8), 1569–1588.

21 <http://doi.org/10.1111/j.1460-9568.2008.06395.x>

22 Ruschel, M., Knösche, T. R., Friederici, A. D., Turner, R., Geyer, S., & Anwender, A. (2014).

23 Connectivity architecture and subdivision of the human inferior parietal cortex revealed by

24 diffusion MRI. *Cerebral Cortex*, 24(9), 2436–48. <http://doi.org/10.1093/cercor/bht098>

25 Sakata, H., Tsutsui, K.-I., & Taira, M. (2005). Toward an understanding of the neural processing for

26 3D shape perception. *Neuropsychologia*, 43(2), 151–161.

27 <http://doi.org/10.1016/j.neuropsychologia.2004.11.003>

28 Saleem, K. S., Kondo, H., & Price, J. L. (2008). Complementary circuits connecting the orbital and

29 medial prefrontal networks with the temporal, insular, and opercular cortex in the macaque

30 monkey. *The Journal of Comparative Neurology*, 506(4), 659–93.

31 <http://doi.org/10.1002/cne.21577>

32 Saleem, K. S., Miller, B., & Price, J. L. (2014). Subdivisions and connective networks of the

33 lateral prefrontal cortex in the macaque monkey. *Journal of Comparative Neurology*, 522(7),

34 1641–1690. <http://doi.org/10.1002/cne.23498>

- 1 Saleem, K. S., Suzuki, W., Tanaka, K., & Hashikawa, T. (2000). Connections between anterior
2 inferotemporal cortex and superior temporal sulcus regions in the macaque monkey. *The*
3 *Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 20(13), 5083–
4 5101. <http://doi.org/20/13/5083> [pii]
- 5
6
7 Schall, J. D., Morel, A., King, D. J., & Bullier, J. (1995). Topography of visual cortex connections
8 with frontal eye field in macaque: convergence and segregation of processing streams. *The*
9 *Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, 15(6), 4464–
10 4487. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7540675>
- 11
12
13 Schmahmann, J. D., Pandya, D. N., Wang, R., Dai, G., D'Arceuil, H. E., De Crespigny, A. J., &
14 Wedeen, V. J. (2007). Association fibre pathways of the brain: Parallel observations from
15 diffusion spectrum imaging and autoradiography. *Brain*, 130(3), 630–653.
16
17 <http://doi.org/10.1093/brain/awl359>
- 18
19
20 Schubotz, R. I., Anwander, A., Knösche, T. R., von Cramon, D. Y., & Tittgemeyer, M. (2010).
21 Anatomical and functional parcellation of the human lateral premotor cortex. *NeuroImage*,
22 50(2), 396–408. <http://doi.org/10.1016/j.neuroimage.2009.12.069>
- 23
24
25 Seltzer, B., & Pandya, D. N. (1978). Afferent cortical connections and architectonics of the superior
26 temporal sulcus and surrounding cortex in the rhesus monkey. *Brain Research*, 149(1), 1–24.
27
28 [http://doi.org/10.1016/0006-8993\(78\)90584-X](http://doi.org/10.1016/0006-8993(78)90584-X)
- 29
30
31 Seltzer, B., & Pandya, D. N. (1984). Further observations on parieto-temporal connections in the
32 rhesus monkey. *Experimental Brain Research*, 55(2), 301–12. Retrieved from
33
34 <http://www.ncbi.nlm.nih.gov/pubmed/6745368>
- 35
36
37 Seltzer, B., & Pandya, D. N. (1989). Frontal lobe connections of the superior temporal sulcus in the
38 rhesus monkey. *The Journal of Comparative Neurology*, 281(1), 97–113.
39
40 <http://doi.org/10.1002/cne.902810108>
- 41
42
43 Seltzer, B., & Pandya, D. N. (1994). Parietal, temporal, and occipital projections to cortex of the
44 superior temporal sulcus in the rhesus monkey: a retrograde tracer study. *The Journal of*
45 *Comparative Neurology*, 343(3), 445–63. <http://doi.org/10.1002/cne.903430308>
- 46
47
48 Sereno, A. B., & Maunsell, J. H. (1998). Shape selectivity in primate lateral intraparietal cortex.
49 *Nature*, 395(6701), 500–503. <http://doi.org/10.1038/26752>
- 50
51
52 Sereno, M. I., & Tootell, R. B. (2005). From monkeys to humans: what do we now know about
53 brain homologies? *Current Opinion in Neurobiology*, 15(2), 135–144.
54
55 <http://doi.org/10.1016/j.conb.2005.03.014>
- 56
57
58 Snyder, L. H., Grieve, K. L., Brotchie, P., & Andersen, R. A. (1998). Separate body- and world-
59 referenced representations of visual space in parietal cortex. *Nature*, 394(6696), 887–91.
60
61
62
63
64
65

<http://doi.org/10.1038/29777>

- 1
2 Stanton, G. B., Bruce, C. J., & Goldberg, M. E. (1995). Topography of projections to posterior
3 cortical areas from the macaque frontal eye fields. *The Journal of Comparative Neurology*,
4 353(2), 291–305. <http://doi.org/10.1002/cne.903530210>
5
6
7 Tal, N., & Amedi, A. (2009). Multisensory visual-tactile object related network in humans: insights
8 gained using a novel crossmodal adaptation approach. *Experimental Brain Research*, 198(2–
9 3), 165–82. <http://doi.org/10.1007/s00221-009-1949-4>
10
11
12 Tanné-Gariépy, J., Rouiller, E. M., & Boussaoud, D. (2002). Parietal inputs to dorsal versus ventral
13 premotor areas in the macaque monkey: Evidence for largely segregated visuomotor pathways.
14 *Experimental Brain Research*, 145(1), 91–103. <http://doi.org/10.1007/s00221-002-1078-9>
15
16
17 Thiebaut de Schotten, M., Dell’Acqua, F., Forkel, S. J., Simmons, A., Vergani, F., Murphy, D. G.
18 M., & Catani, M. (2011). A lateralized brain network for visuospatial attention. *Nature*
19 *Neuroscience*, 14(10), 1245–1246. <http://doi.org/10.1038/nn.2905>
20
21
22 Thiebaut de Schotten, M., Dell’Acqua, F., & Valabregue, R. (2012). Monkey to human comparative
23 anatomy of the frontal lobe association tracts. *Cortex*, 48(1), 82–96.
24
25
26 <http://doi.org/10.1016/j.cortex.2011.10.001>
27
28
29 Thomas, C., Ye, F. Q., Irfanoglu, M. O., Modi, P., Saleem, K. S., Leopold, D. A., & Pierpaoli, C.
30 (2014). Anatomical accuracy of brain connections derived from diffusion MRI tractography is
31 inherently limited. *Proceedings of the National Academy of Sciences*, 111(46), 16574–16579.
32
33
34 <http://doi.org/10.1073/pnas.1405672111>
35
36
37 Toth, L. J., & Assad, J. A. (2002). Dynamic coding of behaviourally relevant stimuli in parietal
38 cortex. *Nature*, 415(6868), 165–8. <http://doi.org/10.1038/415165a>
39
40
41 Uka, T., Tanaka, H., Yoshiyama, K., Kato, M., & Fujita, I. (2000). Disparity selectivity of neurons
42 in monkey inferior temporal cortex. *Journal of Neurophysiology*, 84(1), 120–32. Retrieved
43 from <http://www.ncbi.nlm.nih.gov/pubmed/10899190>
44
45
46 Ungerleider, L. G., & Mishkin, M. (1982). Two visual systems. In D. J. Ingle, M. A. Goodale, & R.
47 J. W. Mansfield (Eds.), *Analysis of visual behavior*. (pp. 549--586). Cambridge: MIT Press.
48
49
50 Vallar, G., & Perani, D. (1986). The anatomy of unilateral neglect after right-hemisphere stroke
51 lesions. A clinical/CT-scan correlation study in man. *Neuropsychologia*, 24(5), 609–622.
52
53 [http://doi.org/10.1016/0028-3932\(86\)90001-1](http://doi.org/10.1016/0028-3932(86)90001-1)
54
55
56 Verhagen, L., Dijkerman, H. C., Grol, M. J., & Toni, I. (2008). Perceptuo-motor interactions during
57 prehension movements. *The Journal of Neuroscience : The Official Journal of the Society for*
58 *Neuroscience*, 28(18), 4726–4735. <http://doi.org/10.1523/JNEUROSCI.0057-08.2008>
59
60
61 Verhoef, B.-E., Vogels, R., & Janssen, P. (2011). Synchronization between the end stages of the
62
63
64
65

dorsal and the ventral visual stream. *Journal of Neurophysiology*, 105(5), 2030–2042.

<http://doi.org/10.1152/jn.00924.2010>

Verhoef, B.-E., Vogels, R., & Janssen, P. (2015). Effects of Microstimulation in the Anterior Intraparietal Area during Three-Dimensional Shape Categorization. *Plos One*, 10(8),

e0136543. <http://doi.org/10.1371/journal.pone.0136543>

Vogt, O., & Vogt, C. (1919). Ergebnisse unserer Hirnforschung. *J Psychol Neurol*, (25), 277–462.

Wang, Y., Fernández-Miranda, J. C., Verstynen, T., Pathak, S., Schneider, W., & Yeh, F.-C. (2013). Rethinking the role of the middle longitudinal fascicle in language and auditory pathways.

Cerebral Cortex, 23(10), 2347–56. <http://doi.org/10.1093/cercor/bhs225>

Webster, M. J., Bachevalier, J., & Ungerleider, L. G. (1994). Connections of inferior temporal areas TEO and TE with parietal and frontal cortex in macaque monkeys. *Cerebral Cortex*, 4(5),

470–83. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7530521>

Wilson, F. A., Scaldidhe, S. P., & Goldman-Rakic, P. S. (1993). Dissociation of object and spatial processing domains in primate prefrontal cortex. *Science*, 260(5116), 1955–8. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/8316836>

Wu, Y., Sun, D., Wang, Y., Wang, Y., & Wang, Y. (2016). Tracing short connections of the temporo-parieto-occipital region in the human brain using diffusion spectrum imaging and fiber dissection. *Brain Research*, 1646, 152–9. <http://doi.org/10.1016/j.brainres.2016.05.046>

Yoshida, M., Naya, Y., & Miyashita, Y. (2003). Anatomical organization of forward fiber projections from area TE to perirhinal neurons representing visual long-term memory in monkeys. *Proceedings of the National Academy of Sciences of the United States of America*, 100(7), 4257–62. <http://doi.org/10.1073/pnas.0736457100>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legends

Figure 1. Schematic view of the lateral, medial, and orbitofrontal cortical surface of the macaque brain showing the areal subdivision of the posterior parietal, temporal and frontal cortex. The superior (SA) and inferior (IA) arcuate, the intraparietal (IP) and the superior temporal (ST) sulci are shown unfolded to display the location of cortical areas buried in their banks. P, C, L, and Lu on the lateral view of the hemisphere indicate the principal, central, lateral, and lunate sulcus, respectively. Cg and PO on the medial view of the hemisphere indicate the cingulate and the parieto-occipital sulcus, respectively. LO and MO in the orbito-frontal cortex indicate the lateral and medial orbital sulcus, respectively. Parietal areas are defined according to architectonic criteria described by Pandya and Seltzer (1982), Luppino, Ben Hamed, Gamberini, Matelli and Galletti (2005), and Gregoriou, Borra, Matelli and Luppino (2006), except for the areas of the IP, defined according to functional and connectional criteria (see Colby, 1998; Rizzolatti et al., 1998). Temporal areas are defined according to architectonic and functional criteria according to Seltzer and Pandya (1978) and Boussaoud, Ungerleider and Desimone (1990) and Hackett, Stepniewska and Kaas (1998). Agranular frontal (motor) areas are defined according to architectonic criteria described by Matelli, Luppino and Rizzolatti (1991) and Belmalih et al. (2009). Prefrontal areas are defined according to architectonic criteria described by Carmichael and Price (1994) and Gerbella, Belmalih, Borra, Rozzi and Luppino (2007).

Figure 2. Summary view of the main frontal connections of the macaque PPC. A: Connections of rostral SPL areas. B: Connections of caudal SPL areas. C: Connections of rostral IPL areas. D: Connections of caudal IPL areas.

Figure 3. Summary view of the main frontal and parietal connections of the macaque temporal cortex. A: Connections of the ventral visual stream areas of the inferotemporal cortex. B: Connections of the multisensory area STP. C: Connections of the auditory parabelt areas of the superior temporal gyrus.

Figure 4. Summary view of the topography of the temporo-parietal connectivity of the macaque brain.

Figure 5. Examples of large-scale functionally specialized temporo-parieto-frontal networks, based on tract tracing connectional data in the macaque brain. A: Network for controlling hand actions. B: Network for controlling oculomotor behavior. C: Network for visuospatial and attentional processing.

Figure 6. Cortical sectors of the human brain involved in action observation and imitation (A) and most frequently correlated with spatial unilateral neglect (B) and their possible interconnections. Based on data from Casper et al. (2010), Rizzolatti & Sinigallia (2016), Corbetta & Shulman, 2011) and Karnath & Rorden (2012)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

*Highlights

Robust connections link the macaque temporal and inferior parietal cortex

These connections are part of large-scale cortical networks involved in motor and cognitive motor functions

This organizational picture appears to have been conserved in primate evolution

Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)







Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)






Figure
[Click here to download high resolution image](#)

