



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk

This is the peer reviewed version of the following article:

Original

Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk / Grosso, Giuseppe; Bella, Francesca; Godos, Justyna; Sciacca, Salvatore; Del Rio, Daniele; Ray, Sumantra; Galvano, Fabio; Giovannucci, Edward L.. - In: NUTRITION REVIEWS. - ISSN 0029-6643. - 75:6(2017), pp. 405-419. [[10.1093/nutrit/nux012](https://doi.org/10.1093/nutrit/nux012)]

Availability:

This version is available at: 11381/2829032 since: 2021-10-05T09:16:44Z

Publisher:

Oxford University Press

Published

DOI:[10.1093/nutrit/nux012](https://doi.org/10.1093/nutrit/nux012)

Terms of use:

openAccess

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

(Article begins on next page)

Journal: *Nutrition Reviews*
Article Doi: 10.1093/nutrit/nux012
Article Title: Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk
First Author: Giuseppe Grosso
Corr. Author: Giuseppe Grosso

OXFORD
UNIVERSITY PRESS

INSTRUCTIONS

We encourage you to use Adobe's editing tools (please see the next page for instructions). If this is not possible, please send a list of corrections (in an email or Word attachment) listing each change in the following manner: line number, current text, change to be made.

Changes should be corrections of typographical errors only. Changes that contradict journal style will not be made.

These proofs are for checking purposes only. They should not be considered as final publication format. The proof must not be used for any other purpose. In particular we request that you do not post them on your personal/institutional web site, and do not print and distribute multiple copies. Neither excerpts nor all of the article should be included in other publications written or edited by yourself until the final version has been published and the full citation details are available. You will be sent these when the article is published, along with an author PDF of the final article.

- 1. License to Publish:** If you have not already done so, please visit the link in your Welcome email and complete your License to Publish online.
 - 2. Author groups:** Please check that all names have been spelled correctly and appear in the correct order. Please also check that complete first and last names appear and that any initials are present. Please check that the author surnames (family name) have been correctly identified by a pink background. If this is incorrect, please identify the full surname of the relevant authors. Occasionally, the distinction between surnames and forenames can be ambiguous, and this is to ensure that the authors' full surnames and forenames are tagged correctly, for accurate indexing online. Please also check all author affiliations.
 - 3. Funding:** Please ensure that a Funding statement appears in the Acknowledgments, detailing any funding that was received or indicating "no external funding was provided". Remember that any funding used while completing this work should be highlighted in a separate Funding section. Please ensure that you use the full official name of the funding body, and if your paper has received funding from any institution, such as NIH, please inform us of the grant number to go into the funding section. We use the institution names to tag NIH funded articles so they are deposited at PMC. If we already have this information, we will have tagged it and it will appear as colored text in the funding paragraph. Please check the information is correct.
 - 4. Declaration of interest:** All authors must make a formal statement indicating any potential conflict of interest that might constitute an embarrassment to any of the authors if it were not to be declared and were to emerge after publication. Such conflicts might include, but are not limited to, shareholding in or receipt of a grant or consultancy fee from a company whose product features in the submitted manuscript or which manufactures a competing product. The following statement has been added to your proof: "Declaration of interest. The authors have no relevant interests to declare." If this is incorrect please supply the necessary text to identify the declaration of interest.
-


MAKING CORRECTIONS TO YOUR PROOF

These instructions show you how to mark changes or add notes to your proofs using Adobe Acrobat Professional versions 7 and onwards, or Adobe Reader DC. To check what version you are using go to **Help** then **About**. The latest version of Adobe Reader is available for free from get.adobe.com/reader.

DISPLAYING THE TOOLBARS


Adobe Reader DC

In Adobe Reader DC, the Comment toolbar can be found by clicking 'Comment' in the menu on the right-hand side of the page (shown below).



Acrobat Professional 7, 8, and 9

In Adobe Professional, the Comment toolbar can be found by clicking 'Comment(s)' in the top toolbar, and then clicking 'Show Comment & Markup Toolbar' (shown below).



The toolbar shown below will then be displayed along the top.




The toolbar shown below will then display along the top.



USING TEXT EDITS AND COMMENTS IN ADOBE ACROBAT

This is the quickest, simplest and easiest method both to make corrections, and for your corrections to be transferred and checked.



1. Click **Text Edits**
2. Select the text to be annotated or place your cursor at the insertion point and start typing.
3. Click the **Text Edits** drop down arrow and select the required action.

You can also right click on selected text for a range of commenting options, or add sticky notes.

SAVING COMMENTS

In order to save your comments and notes, you need to save the file (**File, Save**) when you close the document.


USING COMMENTING TOOLS IN ADOBE READER

All commenting tools are displayed in the toolbar. You cannot use text edits, however you can still use highlighter, sticky notes, and a variety of insert/replace text options.




POP-UP NOTES

In both Reader and Acrobat, when you insert or edit text a pop-up box will appear. In **Acrobat** it looks like this:



In **Reader** it looks like this, and will appear in the right-hand pane:







DO NOT MAKE ANY EDITS DIRECTLY INTO THE TEXT, USE COMMENTING TOOLS ONLY.

Author Query Form

Journal: *Nutrition Reviews*
Article Doi: 10.1093/nutrit/nux012
Article Title: Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk
First Author: Giuseppe Grosso
Corr. Author: Giuseppe Grosso

AUTHOR QUERIES – TO BE ANSWERED BY THE CORRESPONDING AUTHOR

The following queries have arisen during the typesetting of your manuscript. Please click on each query number and respond by indicating the change required within the text of the article. If no change is needed please add a note saying "No change."

- AQ1:** Please check that all names have been spelled correctly and appear in the correct order. Please also check that all initials are present. Please check that the author surnames (family name) have been correctly identified by a pink background. If this is incorrect, please identify the full surname of the relevant authors. Occasionally, the distinction between surnames and forenames can be ambiguous, and this is to ensure that the authors' full surnames and forenames are tagged correctly, for accurate indexing online. Please also check all author affiliations. 
- AQ2:** Please add 1-2 sentences describing data extraction (guidelines for abstracting data and assessing data quality). 
- AQ3:** Please confirm the addition of "hormone" is acceptable. 
- AQ4:** AMA defines "sex" as the classification of living things as male or female. "Gender" is defined as a person's self-representation as man or woman. There are six instances of "gender" in this article. Please check all six and see if "gender" should be changed to "sex" and make revisions, if necessary. 
- AQ5:** Should "3 or more" be "3 or fewer" since "insufficient evidence" is being defined? Please change if necessary. 
- AQ6:** Is "gender" correct, or should it be "sex"? 
- AQ7:** Information in parentheses in this sentence has been revised slightly; eg, reversed: 17.1 SD changed to SD17.1. 
- AQ8:** Footnote "a" does not appear in the table. Please place it where it belongs. 
- AQ9:** Please confirm this sentence is correct. No revisions have been made since copyediting, but a query about this sentence was overlooked. 
- AQ10:** Our journal prefers as few abbreviations as possible, thus "MD" has been reverted to "Mean difference" 
- AQ11:** Please confirm that it is okay to update this reference to the print version. 
- AQ12:** Please confirm the addition of "RR" here is correct, or revise as needed. 
- AQ13:** Could "most likely" be deleted? If yes, please delete. 
- AQ14:** Please confirm this sentence is correct. No further revisions were made since copyediting, but a query about this sentence was overlooked. 
- AQ15:** In the original sentence, "their" referred to "permeability." Please confirm the revised sentence is correct, or revise as needed. 
- AQ16:** Please confirm the slightly revised sentence is correct ("sites" was changed to "types"). 
- AQ17:** Please check this reference. The URL provided was invalid and was changed to the one shown here. Please confirm this reference is correct. 

*Lead Article***Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk**

Giuseppe **Grosso**, Francesca **Bella**, Justyna **Godos**, Salvatore **Sciacca**, Daniele **Del Rio**, Sumantra **Ray**, Fabio **Galvano**, and Edward L. **Giovannucci**

Context: Evidence of an association between dietary patterns derived a posteriori and risk of cancer has not been reviewed comprehensively. **Objective:** The aim of this review was to investigate the relation between a posteriori-derived dietary patterns, grouped as healthy or unhealthy, and cancer risk. The relation between cancer risk and background characteristics associated with adherence to dietary patterns was also examined. **Data Sources:** PubMed and Embase electronic databases were searched. **Study Selection:** A total of 93 studies including over 85 000 cases, 100 000 controls, and 2 000 000 exposed individuals were selected. **Data Extraction. ●●●. Results:** The most convincing evidence (significant results from prospective cohort studies) supported an association between healthy dietary patterns and decreased risk of colon and breast cancer, especially in post-menopausal, hormone receptor-negative women, and an association between unhealthy dietary patterns and increased risk of colon cancer. Limited evidence of a relation between an unhealthy dietary pattern and risk of upper aerodigestive tract, pancreatic, ovarian, endometrial, and prostatic cancers relied only on case-control studies. Unhealthy dietary patterns were associated with higher body mass index and energy intake, while healthy patterns were associated with higher education, physical activity, and less smoking. Potential differences across geographical regions require further evaluation. **Conclusions:** The results suggest a potential role of diet in certain cancers, but the evidence is not conclusive and may be driven or mediated by lifestyle factors.

INTRODUCTION

Cancer is among the leading causes of mortality worldwide, representing a major public health issue. The Global Burden of Disease evaluation estimated the

incidence of cancer at 14.9 million cases, accounting for 8.2 million deaths and 196.3 million disability-adjusted life-years, with prostate cancer and breast cancer causing the greatest impact in men and women, respectively.¹ Ongoing research on modifiable risk factors,

Affiliation: G. Grosso, F. Bella, J. Godos, and S. Sciacca are with the Integrated Cancer Registry of Catania-Messina-Siracusa-Enna, Azienda Ospedaliero-Universitaria Policlinico-Vittorio Emanuele, Catania, Italy. F. Galvano is with the Department of Biomedical and Biotechnological Sciences, University of Catania, Catania, Italy. D. Del Rio is with the Department of Food Science, University of Parma, Parma, Italy. G. Grosso, D. Del Rio, and S. Ray are with the NNEdPro Global Centre for Nutrition and Health, St John's Innovation Centre, Cambridge, United Kingdom. E.L. Giovannucci is with the Department of Epidemiology and the Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; and the Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA.

Correspondence: G. Grosso, Integrated Cancer Registry of Catania-Messina-Siracusa-Enna, Via S. Sofia 85, 95123 Catania, Italy. Email: giuseppe.grosso@studium.unict.it. Phone: +39-095-3782182.

Key words: cancer risk, case-control studies, dietary patterns, meta-analysis, prospective studies.

© The Author(s) 2017. Published by Oxford University Press on behalf of the International Life Sciences Institute. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

doi: 10.1093/nutrit/nux012

Nutrition Reviews® Vol. 0(0):1–15

including dietary habits, estimated that a large percentage of cancer is preventable by behavior modification. Numerous studies have explored and produced evidence of an association between individual nutrients or foods and risk of cancer. While the study of single foods or nutrients provides evidence of the effects of individual components of the diet, it fails to encompass the complexity of the whole diet, including the interactions between all dietary components. The examination of existing dietary patterns and their association with cancer risk could provide a wider and more realistic estimation of the relation between food habits and health in a particular population.

A number of studies have been conducted on the potential association between *a posteriori*-derived dietary patterns and certain types of cancer, such as esophageal,² gastric,³ colorectal,⁴ and breast.⁵ However, using principal component analysis and exploratory factor analysis, no comprehensive evaluation of all cancer sites has been conducted. Moreover, an association between cancer and unhealthy behaviors has been previously discussed but not quantitatively investigated. Thus, the aim of the present study was threefold. The first aim was to update and summarize current knowledge on the relation between adherence to *a posteriori*-derived healthy or unhealthy dietary patterns (characterized by, but not limited to, red and processed meat, sugary drinks and salty snacks, starchy foods, and refined carbohydrates) and cancer risk in epidemiological studies. The second aim was to provide risk estimation of different cancer types associated with similar dietary patterns to test consistency of findings. The final aim was to explore the association between the dietary patterns investigated and factors potentially clustering together to mediate or confound the resulting association.

METHODS

Study selection

The design, analysis, and reporting of this study is compliant with the MOOSE (Meta-analysis of Observational Studies in Epidemiology) guidelines.⁶ A systematic search of the PubMed and Embase databases for all English-language studies published up to December 2015 was performed. The search terms used for the study selection were “dietary patterns” associated with the target cancer sites, including “esophagus, esophageal,” “stomach, gastric,” “colon, rectal, rectum, colorectal, colorectum,” “liver/hepatic,” “pancreas/pancreatic,” “ovarian,” “breast,” “endometrial, endometrium,” “prostate, prostatic,” “lung,” and “thyroid.” Studies were included if they met the following inclusion criteria (Table 1): (1) case-control or

Table 1 PICOS criteria for inclusion and exclusion of studies

Criteria	Description
Participants	General population
Intervention/exposure	Adherence to <i>a posteriori</i> -derived dietary pattern
Comparison	Highest vs lowest category of exposure
Outcome	Cancer risk/incidence
Study design	Cohort and case-control studies

prospective study design; (2) dietary patterns derived empirically with *a posteriori* methods using principal component analysis and exploratory factor analysis; and (3) estimated odds ratios (ORs) or hazard ratios for a well-defined outcome, namely a specific cancer site. Exclusion criteria were the lack of statistics or results and the lack of an assessed composite outcome (eg, overall cancer mortality). Reference lists of included manuscripts were also checked for studies not identified previously. If more than 1 study was conducted on the same population, only the most recent was included. However, all patterns identified in each study conducted on the same population were considered, and those that were more consistent and comparable with others were included. If more than 1 article was published that used the same sample or cohort, only the study that included the entire cohort or the study with the longest follow-up or the most comparable dietary patterns identified was included.

Data extraction

Data were abstracted from each identified study by using a standardized form. The following information was collected: tumor site; name of first author, year of publication; study location; study design; name of study cohort (when available); number of participants/controls; number of cases; age range and body mass index (BMI) at baseline; gender of participants; dietary assessment method; dietary patterns identified; food items included; energy intake (in kilocalories per day) and amount of main food group consumed (in grams per day) in the highest and the lowest categories of exposure; covariates used in adjustments; and ORs or hazard ratios and 95% CIs for the highest vs the lowest category of exposure. This process was performed independently by two authors (G.G. and J.G.), and discrepancies were discussed and resolved by consensus.

Level of evidence

In accordance with the modified Joint World Health Organization–Food and Agriculture Expert Consultation criteria for evidence in nutrition,⁷ a

“possible” association was considered when results were supported by meta-analysis of prospective studies with no evidence of significant heterogeneity or publication bias, a “limited” association was considered when results were supported by meta-analysis of prospective studies with high heterogeneity or meta-analysis of case-control studies, and “insufficient evidence” was considered when meta-analysis included 3 or more studies.

10 Statistical analysis

The mean amount of alcohol, fruit, vegetable, and meat consumption per day in the highest category of exposure to healthy and unhealthy dietary patterns was calculated. Random-effects models were used to calculate pooled relative risks (RRs) and 95% CIs for the highest vs the lowest category of exposure. Prospective and case-control studies were considered separately. The risk estimate from the most fully adjusted models in the analysis of the pooled RR was used. Heterogeneity was assessed by using the Q test and the I^2 statistic. The level of significance for the Q test was defined as $P < 0.10$. The I^2 statistic represented the amount of total variation that could be attributed to heterogeneity. I^2 values $\geq 50\%$ indicated significant heterogeneity. To test for the potential source of heterogeneity, several subgroup analyses were performed. First, since the definition of an unhealthy dietary pattern was substantially arbitrary and included different key characteristics, further subgroup analyses were conducted in which foods were grouped into the following dietary patterns: “alcohol” pattern, including foods associated with alcohol drinking; “carbohydrate/refined” pattern, including refined sugars, sugary drinks, and sweets; “fat/salty” pattern, including fatty or salty snacks and fast foods; “animal/meat” pattern, including mainly red and processed meat; and “Western” pattern, when defined as such in the reference. Second, the possible influence of geographical area on the relation between dietary patterns and cancer risk was tested. The possible presence of publication bias was tested through visual examination of funnel plots. Publication bias was further evaluated by performing a sensitivity analysis using a single dataset that referred to 1 dietary pattern per study, in order to avoid to prevent individual publications from being weighted too heavily.

To identify a potential relation between dietary patterns and lifestyle factors, data on baseline BMI values and relative standard deviations (SDs) by category of exposure (quantiles of dietary patterns) were extracted. Missing SDs were imputed by considering the average of existing values according to guidelines in the *Cochrane Handbook for Systematic Reviews of*

*Interventions*⁸; the difference between the highest and the lowest quantiles of each dietary pattern investigated was calculated, and values were subsequently meta-analyzed to estimate the mean difference using random-effects models. To identify other factors potentially related to dietary patterns, data on smoking status, education, and physical activity were collected, the ORs between the highest and the lowest (reference) quantiles of each dietary pattern investigated were calculated, and the results were meta-analyzed using random-effects models. The same strategies to control for heterogeneity and publication bias were adopted. All analyses were conducted by using Review Manager (RevMan) software, version 5.2 (Nordic Cochrane Centre, Copenhagen, Denmark).

RESULTS

Study characteristics

The literature search process is shown in Figure 1. The search strategy led to full-text examination of articles reporting 141 studies, 51 of which were excluded for the following reasons: 15 used a priori dietary pattern scores; 12 explored the association with cancer risk between dietary patterns (not adherence within individual patterns); 7 were conducted on the same population; 6 were conducted on the same cohort but had shorter follow-up; 5 used other methods to assess dietary patterns; and 4 explored the association with cancer survival. One study was added through manual search of references; thus, 93 studies were included in the quantitative analysis, divided by site of cancer as follows: mouth/pharynx (n = 3),⁹⁻¹¹ larynx (n = 2),^{9,10} esophagus (n = 8),^{9,12-18} stomach (n = 9),^{9,12,14,19-24} colorectum (n = 21),²⁵⁻⁴⁵ pancreas (n = 5),⁴⁶⁻⁵⁰ liver (n = 1),⁵¹ breast (n = 31),^{9,31,52-80} endometrium (n = 5),⁸¹⁻⁸⁵ ovarian (n = 3),^{55,86,87} lung (n = 4),^{9,88-90} thyroid (n = 1),⁹¹ urinary tract (n = 2),^{9,92} prostate (n = 10).^{9,93-101}

The main characteristics of the studies included, grouped by cancer site, are reported in Table S1 in the Supporting Information online. Studies included have been conducted in various geographic areas, including the United States, South America, Europe, Asia, and Australia. The average age range of the population included was 40–80 years. Most of the studies included covariates that may significantly influence cancer risk, such as age, gender (when not analyzed separately), and education, as well as factors that may be strongly associated with healthy/unhealthy dietary patterns (ie, BMI) and more general lifestyle behaviors (ie, physical activity and smoking status) (see Table S1 in the Supporting Information online). A total of over 85 000 cases,

AQ6




Figure 1 Flow diagram of the literature search process.

100 000 controls, and 2 000 000 exposed individuals were considered in the analyses (see Table S2 in the Supporting Information online). The number of cases varied greatly, according to the primary site of cancer; cancers of the breast, colorectum, and prostate were the most commonly studied (see Table S2 in the Supporting Information online).

On the basis of food or nutrient constituents of the dietary patterns described in the studies, 315 datasets on healthy patterns and 358 datasets on unhealthy patterns were identified. The former included mainly fruit- and vegetable-based dietary patterns, named “healthy” or “prudent,” and some “traditional” patterns (see Table S3 in the Supporting Information online). Traditional

patterns were a more heterogeneous group of dietary patterns and were described as “Western,” “animal,” “fat and salty,” or “refined” and included not only animal products but also salty/sweet snacks, fatty foods, and refined foods (see Table S3 in the Supporting Information online). Only 6 studies provided individual amounts of main food groups by category of exposure.^{9,34,46,70,73,75} On average, the highest quantile of healthy and unhealthy dietary patterns contained, respectively, 11.5 (SD = 17.1) and 26.4 (SD = 41.6) grams of alcohol per day, 390.6 (SD = 494.9) and 230.5 (SD = 194.8) grams of fruit per day, 287 (SD = 263) and 175.9 (SD = 113) grams of vegetables per day, and 67.8 (SD = 42.5) and 95.8 (SD = 67.6) grams of meat per day.

AQ7

Table 2 Summary of results on the association between high adherence to healthy dietary patterns and risk of cancer, by type of cancer and study design

Type of cancer	Study design						
	Case-control			Prospective			
	No. of datasets	RR (95%CI)	<i>I</i> ² (%)	No. of datasets	RR (95%CI)	95%CI	<i>I</i> ² (%)
Oral/pharyngeal	3	0.63 (0.42–0.94)	69	NA	NA	NA	NA
Laryngeal	2	0.50 (0.36–0.71)	10	NA	NA	NA	NA
Esophageal squamous cell carcinoma	18	0.65 (0.51–0.82)	76	NA	NA	NA	NA
Esophageal squamous cell carcinoma, M	5	0.67 (0.41–1.09)	86	NA	NA	NA	NA
Esophageal squamous cell carcinoma, F	4	0.76 (0.45–1.30)	79	NA	NA	NA	NA
Esophageal adenocarcinoma	7	0.98 (0.63–1.54)	83	NA	NA	NA	NA
Stomach	13	0.72 (0.58–0.90)	70	4	0.98 (0.74–1.29)		43
Stomach, gastric cardia	5	0.86 (0.68–1.09)	28	NA	NA	NA	NA
Stomach, M	2	0.64 (0.23–1.75)	93	2	1.14 (0.89–1.46)		0
Stomach, F	2	0.81 (0.40–1.67)	73	2	0.76 (0.44–1.33)		59
Colorectal	13	0.48 (0.27–0.84)	97	15	0.89 (0.83–0.95)		12
Colorectal, M	3	0.53 (0.40–0.69)	0	5	0.89 (0.78–1.01)		43
Colorectal, F	2	0.45 (0.30–0.67)	5	8	0.92 (0.83–1.02)		0
Colon	9	0.79 (0.68–0.92)	27	14	0.87 (0.81–0.94)		0
Colon, M	2	0.62 (0.41–0.93)	0	6	0.87 (0.79–0.97)		0
Colon, F	1	0.45 (0.24–0.84)	NA	7	0.86 (0.76–0.98)		0
Colon, proximal	3	0.97 (0.67–1.39)	19	10	0.89 (0.79–1.00)		0
Colon, distal	3	0.75 (0.43–1.31)	70	9	0.85 (0.75–0.96)		0
Rectal	10	0.76 (0.59–0.98)	72	13	0.95 (0.78–1.16)		59
Rectal, M	2	0.55 (0.38–0.80)	0	5	0.90 (0.64–1.26)		74
Rectal, F	1	0.66 (0.33–1.32)	NA	6	1.11 (0.92–1.34)		0
Pancreatic	6	0.67 (0.50–0.91)	52	7	1.09 (0.92–1.31)		12
Pancreatic, M	2	0.53 (0.37–0.76)	0	1	1.88 (1.06–3.33)		NA
Pancreatic, F	2	0.73 (0.36–1.49)	67	6	1.04 (0.87–1.24)		0
Endometrial	6	0.83 (0.67–1.04)	32	2	0.96 (0.82–1.12)		0
Ovarian	3	0.86 (0.73–1.00)	0	2	1.33 (0.89–1.97)		46
Lung	3	0.86 (0.52–1.40)	73	4	0.78 (0.65–0.93)		0
Liver	NA	NA	NA	4	0.80 (0.50–1.29)		68
Thyroid	1	0.68 (0.36–1.28)	NA	NA	NA	NA	NA
Urinary tract	5	0.88 (0.61–1.25)	24	NA	NA	NA	NA
Breast	20	0.66 (0.55–0.80)	87	13	0.90 (0.84–0.96)		26
Breast, premenopausal	7	0.64 (0.43–0.94)	80	10	0.95 (0.83–1.10)		27
Breast, postmenopausal	10	0.73 (0.55–0.96)	89	13	0.93 (0.87–1.00)		19
Breast, receptor positive	2	0.40 (0.05–2.88)	97	8	0.93 (0.85–1.02)		35
Breast, receptor negative	2	0.42 (0.08–2.31)	92	8	0.79 (0.65–0.95)		41
Prostate	9	0.99 (0.85–1.15)	18	5	0.99 (0.90–1.08)		0
Prostate, low grade	3	1.07 (0.76–1.51)	0	4	0.97 (0.87–1.09)		13
Prostate, high grade	3	1.13 (0.82–1.55)	0	4	1.00 (0.82–1.21)		0

Abbreviations: F, female; M, male; NA, not applicable; RR, relative risk.

Healthy dietary patterns and cancer risk

The comprehensive evaluation of the association between greater adherence to a healthy dietary pattern and cancer risk by study design (case-control/prospective) is presented in Table 2. Among prospective studies, a higher adherence to healthy dietary patterns was associated with a decreased risk of colorectal cancer (RR = 0.89; 95%CI 0.83–0.95), with no evidence of heterogeneity or publication bias (see Figure S1 and Figure S2 in the Supporting Information online). When analysis was restricted to a specific digestive tract location, a significantly decreased risk was found for the colon (RR = 0.87; 95%CI 0.81–0.94; *I*² = 27%). A decreased

risk was also found in analyses stratified by specific colon location (proximal/distal) and gender, but not in analyses restricted to rectal cancer (Table 2). When the association between risk of other cancers and adherence to healthy dietary patterns was examined prospectively, lung and breast cancer risk was significantly decreased (RR = 0.78, 95%CI 0.65–0.93, and RR = 0.90, 95%CI 0.84–0.96, respectively), with little evidence of heterogeneity and no evidence of publication bias (see Figure S1, Figure S3, and Figure S4 in the Supporting Information online). However, the analysis of lung cancer risk was based on only 2 studies. Analyses of breast cancer risk stratified by menopausal status and hormone receptor status showed a significantly decreased

AQ6

AQ9

20

25

AQ3

risk in postmenopausal women (RR = 0.93; 95%CI 0.87–1.00) and in receptor-negative women (RR = 0.79; 95%CI 0.65–0.95). For no other cancer was there evidence of an association when prospective studies were analyzed, but analysis was based on a low number of studies for most cancers; only pancreatic cancer (n = 7 studies) and prostate cancer (n = 5 studies) were represented by more than a few studies.

Among the case-control studies, inverse associations were confirmed for colorectal and breast cancers, with effect estimates being considerably stronger than those for prospective studies (Table 2). An association was not confirmed for lung cancer, but only 2 studies were reported. Some other cancers, such as gastric (see Figure S5 in the Supporting Information online), pancreatic (see Figure S6 in the Supporting Information online), and ovarian (see Figure S7 in the Supporting Information online), were significantly related to healthy dietary patterns in meta-analyses of case-control studies (Table 2). Adherence to healthy dietary patterns was also associated with a significantly reduced risk of oral/pharyngeal, laryngeal (see Figure S8 in the Supporting Information online), and esophageal cancers (see Figure S9 in the Supporting Information online), though evidence of heterogeneity was high (Table 2). No significant results were found for thyroid, urinary tract, endometrial (see Figure S10 in the Supporting Information online), or prostate cancer (see Figure S11 in the Supporting Information online).

Unhealthy dietary patterns and cancer risk

The results of the analysis of high adherence to unhealthy dietary patterns and cancer risk are shown in Table 3. In prospective studies, unhealthy dietary patterns were associated with a higher risk of colorectal cancer (RR = 1.13; 95%CI 1.03–1.23; $I^2 = 31\%$), with no evidence of publication bias (see Figure S12 and Figure S13 in the Supporting Information online). Subanalyses by cancer site showed significant results for colon cancer (proximal and distal) in prospective studies (Table 3). Pooled analysis of prospective studies on colon cancer showed that higher adherence to unhealthy dietary patterns was associated with an 18% increased risk of cancer (RR = 1.18; 95%CI 1.06–1.31), with evidence of moderate heterogeneity ($I^2 = 31\%$). Heterogeneity was reduced when colon sites were analyzed separately (Table 3). For both colorectal and colon cancer, analysis by gender showed a higher association of unhealthy dietary patterns with increased cancer risk only among women (with negligible heterogeneity between prospective studies), while no significant results were found for rectal cancer when men and women were analyzed separately. For colorectal cancer, all

results of prospective studies were concordant with results of case-control studies, though the magnitude of the association was stronger for the latter (Table 3). Results for ovarian cancer were significant when prospective studies were analyzed, but this was based on only 2 studies (see Figure S14 in the Supporting Information online). Analyses of case-control studies on ovarian cancer, again based on only 2 studies, were compatible, but the association was not significant.

For breast cancer (see Figure S15 in the Supporting Information online), pancreatic cancer (see Figure S16 in the Supporting Information online), lung cancer (see Figure S17 in the Supporting Information online), and prostatic cancer (see Figure S18 in the Supporting Information online), adherence to unhealthy dietary patterns was associated with an increased risk in case-control studies, but results from prospective studies were null, and the point estimates were incompatible with those from case-control studies. For gastric cancer (see Figure S19 in the Supporting Information online) and endometrial cancer (see Figure S20 in the Supporting Information online), a positive association was observed in case-control studies, but the number of prospective studies was not adequate to confirm or refute these findings. Results from pooled case-control analyses on laryngeal and esophageal squamous cell carcinoma (see Figure S21 in the Supporting Information online) were also significant; no prospective data were available (Table 3). No significant results were found for liver, thyroid, urinary tract, or oral/pharyngeal cancers (see Figure S21 in the Supporting Information online), though only oral/pharyngeal cancers were represented by more than a few studies.

Subgroup analyses

When grouping unhealthy dietary patterns by the key component characterizing them, only dietary patterns related to alcohol consumption were associated with increased cancer risk in prospective studies, whereas inconclusive results were found for all other dietary patterns identified (see Table S4 in the Supporting Information online). Notably, most of analyses were conducted on a very limited number of prospective studies, with the exception of the animal/meat and Western dietary patterns. By contrast, irrespective of the number of case-control studies analyzed, all analyses showed a significantly higher cancer risk for high adherence to any of the unhealthy dietary patterns identified (see Table S4 in the Supporting Information online). However, when prospective studies that showed an association between unhealthy dietary patterns and individual cancer sites were analyzed, the animal dietary

AQ6

Table 3 Summary of results on the association between high adherence to unhealthy dietary patterns and risk of cancer, by type of cancer and study design

Type of cancer	Study design					
	Case-control			Prospective		
	No. of datasets	RR (95%CI)	I ² (%)	No. of datasets	RR (95%CI)	I ² (%)
Oral/pharyngeal	8	1.17 (0.88–1.57)	83	NA	NA	NA
Laryngeal	5	1.64 (1.40–1.92)	0	NA	NA	NA
Esophageal squamous cell carcinoma	19	1.78 (1.10–2.87)	96	NA	NA	NA
Esophageal squamous cell carcinoma, M	6	1.66 (1.09–2.52)	84	NA	NA	NA
Esophageal squamous cell carcinoma, F	3	1.36 (0.55–3.35)	89	NA	NA	NA
Esophageal adenocarcinoma	8	1.42 (0.95–2.13)	80	NA	NA	NA
Stomach	21	1.41 (1.15–1.72)	73	8	1.18 (0.85–1.64)	78
Stomach, gastric cardia	4	1.16 (0.74–1.83)	73	NA	NA	NA
Stomach, M	4	1.36 (1.05–1.76)	37	4	1.13 (0.68–1.86)	87
Stomach, F	4	1.22 (0.73–2.02)	65	4	1.26 (0.80–1.97)	66
Colorectal	15	1.54 (1.31–1.82)	51	18	1.13 (1.03–1.23)	31
Colorectal, M	4	1.34 (1.04–1.72)	0	5	1.08 (0.91–1.27)	26
Colorectal, F	3	1.40 (0.93–2.12)	37	10	1.26 (1.10–1.43)	19
Colon	15	1.24 (1.00–1.54)	82	18	1.18 (1.06–1.31)	31
Colon, M	2	2.03 (1.33–3.10)	0	7	1.11 (0.97–1.26)	0
Colon, F	2	1.58 (1.04–2.42)	0	9	1.36 (1.15–1.62)	36
Colon, proximal	3	1.58 (0.89–2.82)	66	10	1.18 (1.00–1.41)	15
Colon, distal	3	1.64 (0.82–3.28)	75	10	1.28 (1.09–1.50)	0
Rectal	11	1.23 (1.02–1.48)	51	15	1.06 (0.90–1.25)	27
Rectal, M	2	1.03 (0.66–1.60)	24	5	1.04 (0.72–1.50)	56
Rectal, F	2	0.77 (0.45–1.30)	0	8	1.01 (0.78–1.31)	25
Pancreatic	8	1.47 (1.17–1.85)	28	3	0.81 (0.59–1.12)	0
Pancreatic, M	3	1.63 (1.16–2.28)	12	1	0.89 (0.47–1.69)	NA
Pancreatic, F	3	1.05 (0.75–1.48)	0	2	0.79 (0.54–1.14)	0
Endometrial	8	1.19 (0.99–1.43)	44	3	1.10 (0.94–1.29)	0
Ovarian	4	1.27 (0.70–2.32)	93	2	1.46 (1.02–2.08)	0
Lung	7	1.24 (0.99–1.56)	54	12	1.11 (0.91–1.35)	66
Liver	NA	NA	NA	1	0.96 (0.50–1.84)	NA
Thyroid	1	0.96 (0.50–1.84)	NA	NA	NA	NA
Urinary tract	8	1.01 (0.71–1.44)	77	NA	NA	NA
Breast	29	1.33 (1.15–1.53)	88	13	1.00 (0.93–1.08)	40
Breast, premenopausal	8	1.17 (1.00–1.38)	45	6	0.98 (0.85–1.13)	2
Breast, postmenopausal	9	1.15 (0.96–1.39)	75	8	1.07 (0.98–1.17)	24
Breast, receptor positive	3	1.14 (0.80–1.62)	67	6	1.11 (0.97–1.27)	64
Breast, receptor negative	3	0.78 (0.56–1.10)	29	6	0.94 (0.80–1.10)	0
Prostate	16	1.44 (1.21–1.71)	62	4	0.87 (0.71–1.07)	59
Prostate, low grade	4	1.29 (0.73–2.28)	58	2	0.98 (0.82–1.17)	43
Prostate, high grade	4	1.28 (0.78–2.07)	52	2	1.06 (0.83–1.35)	0

Abbreviations: F, female; M, male; NA, not applicable; RR, relative risk.

pattern was significantly associated with an increased risk of colorectal cancer (RR = 1.15; 95%CI 1.00–1.32).

In another subgroup analysis, the relation between adherence to healthy or unhealthy dietary patterns and cancer risk was evaluated by country (see Table S5 in the Supporting Information online). Regardless of the cancer site, adherence to healthy dietary patterns was associated with decreased cancer risk of roughly 5% to 10% in case-control and prospective studies conducted in both North America and Europe (for prospective studies, RR = 0.94, 95%CI 0.89–0.99, and RR = 0.89, 95%CI 0.83–0.96, respectively), with no evidence of heterogeneity or publication bias. The meta-analysis of studies involving adherence to unhealthy dietary patterns showed significantly increased risk of cancer in

Europe and North America in case-control studies, but findings among prospective studies were significant only for those conducted in North America (RR = 1.07; 95%CI 1.00–1.14).

Association between dietary patterns and lifestyle factors

A total of 34 studies^{9,12,13,16,20,23,30–35,38,43,46,49–51,53,54,59,62,63,70,73,75,80,82,87,91,93,95,97,99} provided data for the analysis of the associations between dietary patterns and lifestyle risk factors. Although all analyses were affected by high heterogeneity, the difference in BMI between the highest and the lowest categories of exposure was not significant across healthy dietary patterns, while

AQ6 Table 4 Mean difference in body mass index (BMI) between the highest and the lowest categories of exposure to healthy and unhealthy dietary patterns, in total and by study design, gender, and geographical area

	Healthy patterns		Unhealthy patterns	
	No. of datasets	Mean difference in BMI ^a (95%CI)	No. of datasets	Mean difference in BMI ^a (95%CI)
Total, all studies	38	0.21 (−0.14, 0.57)	56	0.29 (0.10–0.48)
Study design				
Case-control	23	0.31 (−0.08, 0.71)	38	0.24 (0.00–0.48)
Prospective	15	0.07 (−0.54, 0.68)	18	0.38 (0.03–0.73)
Gender				
Male	11	0.03 (−0.40, 0.46)	16	0.16 (−0.32, 0.65)
Female	17	0.17 (−0.50, 0.85)	26	0.51 (0.21–0.80)
Geographical area				
North America	10	−0.13 (−1.05, 0.79)	5	1.2 (0.75–1.65)
Europe	7	0.28 (−0.41, 0.98)	3	0.6 (−0.32, 1.52)
Asia	15	0.31 (−0.14, 0.76)	10	−0.1 (−0.41, 0.21)

^aAll analyses showed significant heterogeneity.

it was greater with higher adherence to unhealthy dietary patterns (RR = 0.29; 95%CI 0.10–0.48; Table 4).

The results were consistent both in case-control and in prospective studies, and most likely in women (Table 4).

Interestingly, analyses by geographical region, despite being limited to a relatively low number of datasets, revealed that the mean difference in BMI between high and low adherence to unhealthy dietary patterns was significant (and particularly higher) only when considering studies conducted in North America (mean difference = 1.2; 95%CI 0.75–1.65; Table 4).

Results for other lifestyle factors are presented in Table 5. High exposure to healthy dietary patterns was significantly associated with high physical activity (OR = 1.53; 95%CI 1.22–1.91) and high education level (OR = 1.39; 95%CI 1.06–1.82), while no significant associations between unhealthy dietary patterns and lifestyle factors were found. Results for healthy dietary patterns and high physical activity were relatively stable over each subgroup analysis (no differences between types of study design or between males and females), despite the lack of significant results for the limited number of European studies (Table 5). In contrast, results for high education level were significant only among prospective studies involving males outside of North America (Table 5). Finally, both healthy and unhealthy dietary patterns were associated with smoking status (for current smoker: OR = 0.48, 95%CI 0.36–0.65; and OR = 1.35, 95%CI 1.07–1.69, respectively) (Table 5).

DISCUSSION

Despite the subjective nomenclature applied in each study included in this meta-analysis, dietary patterns based on healthy and unhealthy food choices were identified, and cancer risk was examined by cancer site,

pattern type, and geographical area of the study. The average consumption of selected food groups in healthy dietary patterns was roughly in line with general dietary guidelines, ie, 2–2.5 cups of fruit and vegetables per day and 12.5 oz of meat per week.¹⁰² Alcohol consumption was equivalent to 1 serving of wine or beer per day. When considering data from prospective studies, most of the significant results were limited to the association between high adherence to a healthy dietary pattern and decreased risk of colon, breast, and lung cancer and the association between high adherence to an unhealthy dietary pattern and increased risk of colon cancer. The risk of cancer at several other sites was also associated with dietary pattern, but evidence was mainly driven by case-control studies. However, an overall trend toward increased and decreased cancer risk associated with unhealthy and healthy dietary patterns, respectively, was found, suggesting that diet-related choices could significantly affect the risk of cancer. The present analysis examined the relation between dietary patterns and lifestyle behaviors, using, for the first time, a quantitative synthesis of the literature. The results show that unhealthy dietary patterns (including red and processed meat, sugary drinks and salty snacks, starchy foods, and refined carbohydrates) are associated with a higher BMI, while healthy dietary patterns are related to healthier lifestyle factors and background (including education, physical activity, and smoking habit).

Mechanisms linking dietary patterns and risk of cancer may rely on 2 main aspects: (1) the specific content of the diet, and (2) the association between dietary pattern and altered energy imbalance, body weight, and consequent homeostasis alterations. Diets rich in fruit, vegetables, legumes, and whole grains may have beneficial effects stemming from the balanced ratio of fatty acids, the high fiber content, and the substantial amounts of antioxidant compounds, which inhibit multiple cancer-related

Table 5 Association of lifestyle factors at baseline/controls between the highest vs the lowest category (reference) of exposure to healthy and unhealthy dietary patterns, in total and by study design, gender, and geographical area

AQ6

Lifestyle factor	Healthy patterns		Unhealthy patterns	
	No. of datasets	OR (95%CI)	No. of datasets	OR (95%CI)
High physical activity				
Total	25	1.53 (1.22–1.91)	34	0.9 (0.78–1.04)
Study design				
Case-control	13	1.53 (1.18–1.98)	20	0.85 (0.72–1.01)
Prospective	12	1.53 (1.12–2.10)	14	0.97 (0.79–1.20)
Gender				
Male	9	1.6 (1.36–1.88)	12	0.92 (0.66–1.28)
Female	10	1.69 (1.10–2.58)	16	0.86 (0.71–1.04)
Geographical area				
North America	7	1.76 (1.05–2.94)	6	0.69 (0.43–1.11)
Europe	4	1.15 (0.68–1.96)	9	1.06 (0.85–1.32)
Asia	10	1.56 (1.12–2.17)	12	0.92 (0.74–1.15)
High education level				
Total	21	1.39 (1.06–1.82)	28	1.05 (0.85–1.31)
Study design				
Case-control	7	1.3 (0.97–1.75)	13	1.35 (0.74–2.47)
Prospective	14	1.45 (1.05–2.01)	15	0.86 (0.66–1.14)
Gender				
Male	6	1.48 (1.02–2.17)	8	0.61 (0.44–0.84)
Female	10	1.26 (0.80–1.97)	11	1.1 (0.85–1.43)
Geographical area				
North America	9	1.36 (0.83–2.22)	9	0.85 (0.53–1.37)
Europe	1	1.28 (1.24–1.32)	3	1.03 (0.68–1.56)
Asia	8	1.64 (1.27–2.12)	9	1.18 (0.74–1.88)
Smoking (current)				
Total	19	0.48 (0.36–0.65)	53	1.35 (1.07–1.69)
Study design				
Case-control	19	0.49 (0.34–0.70)	31	1.29 (1.00–1.67)
Prospective	11	0.42 (0.27–0.65)	22	1.4 (0.99–1.98)
Gender				
Male	15	0.59 (0.39–0.89)	15	1.46 (1.12–1.90)
Female	13	0.4 (0.24–0.68)	22	1.2 (0.82–1.77)
Geographical area				
North America	5	0.73 (0.56–0.95)	12	1.4 (0.79–2.49)
Europe	14	0.54 (0.36–0.79)	8	1.39 (0.77–2.51)
Asia	10	0.47 (0.28–0.77)	17	1.21 (0.82–1.77)

Abbreviations: OR, odds ratio.

AQ8

^aAll analyses showed significant heterogeneity.

biological pathways, including carcinogen bioactivation, cell signaling, cell cycle regulation, angiogenesis, and inflammation.¹⁰³ With regard to colon cancer in particular, a dietary pattern rich in fiber may improve fecal bulking, increase satiety, viscosity, and short-chain fatty acid production, enhance fermentation of metabolites, increase stool bulk (thus reducing transit time and contact of carcinogens with the colonic mucosa), and act as a probiotic.¹⁰⁴

In contrast, unhealthy dietary patterns are characterized by, but not limited to, red and processed meat, sugary drinks and salty snacks, starchy foods, and refined carbohydrates. Red and, especially, processed meats are rich in salt, *N*-nitroso compounds, heterocyclic amines, heme iron, and, following cooking at high temperature, polycyclic aromatic hydrocarbons, which have been considered responsible for the carcinogenic

effects of meat consumption.¹⁰⁵ Alcohol-related dietary patterns are characterized by a high intake of spirits and fortified wines, demonstrated to increase the risk of cancer.¹⁰⁶ High alcohol consumption may lead to higher estrogen concentrations resulting from decreased metabolic clearance, thereby potentially increasing the risk of breast cancer.¹⁰⁷ It can also increase the permeability of membranes to carcinogens and inhibit the detoxification of carcinogens.¹⁰⁷ By contrast, diets containing antioxidant vitamins could have protective effects via anti-inflammatory action on estrogen metabolism and reduction of cell proliferation.¹⁰⁸ Finally, a diet rich in antioxidants may counter chronic inflammation caused by unhealthy lifestyle factors (ie, smoking), which may explain the association between healthy dietary patterns and decreased risk of lung cancer, especially in smokers.¹⁰⁹

In this study, there was also an indication of a possible relation between adherence to healthy or unhealthy dietary patterns and risk of esophageal, laryngeal, gastric, ovarian, and pancreatic cancers, although evidence was limited to case-control studies. Moreover, despite no evidence of a benefit of healthy dietary patterns against prostate cancer risk, a significantly increased risk associated with unhealthy dietary patterns was found when examining case-control studies. Overall, the mechanisms responsible for an association between dietary patterns and the types of cancer mentioned above in this paragraph may be similar to those previously reported, including the potential direct effects of certain foods on specific types of cancer (eg, the effect of salted foods on upper aerodigestive tract cancers)¹¹⁰ and the effects of hormonal alterations related to age (ie, menopause) or weight status.¹¹¹ For prostate cancer risk specifically, it is noteworthy that the effects of diet on cancer risk are more likely to be observed with advanced rather than total prostate cancers.¹¹¹ However, additional prospective studies are needed to verify whether the associations in case-control studies found in this meta-analysis can be confirmed.

Besides the qualitative aspects of dietary patterns, it is hypothesized that unhealthy dietary patterns may lead to increased body weight, leading to increased risk of obesity-related cancer. In support of such hypothesis, the results showed that unhealthy dietary patterns were associated with higher BMI in a summary meta-analysis of study populations, in line with the hypothesis that individuals who adhere to an unhealthy dietary pattern may have a higher cancer risk through the mediating effect of increased body weight and obesity. A diet rich in saturated fats and refined sugars may lead to increased body fat accumulation and impaired glucose and insulin regulation, which in turn alters physiological hormonal homeostasis and ultimately increases cancer risk.¹¹² Indeed, refined sugars may affect both the glycemic index (which indicates the absorption of carbohydrates and, hence, measures insulin demand) and the glycemic load (which measures both the quality and the quantity of carbohydrate consumed), which have been related to cancer risk because of the potential role of insulin and insulin-like growth factor in cancer promotion.¹¹³ Adipocytes and visceral adipose tissue produce leptin and adiponectin, 2 hormones with the capacity to regulate immune function, inflammatory cytokines, angiogenesis, insulin resistance, and other biological processes that have been associated with some cancers.^{7,114} Adipose tissue is also responsible for a low-grade, sub-clinical state of chronic inflammation, characterized by increased circulating inflammatory cytokines (interleukin 1 β , interleukin 6, tumor necrosis factor- α , monocyte

chemoattractant protein 1, and C-reactive protein, all of which, in turn, are associated with intracellular transcription factors involved in cell proliferation, apoptosis, inflammation, metastasis, and angiogenesis.¹¹⁵ Other mechanisms potentially linking increased BMI to cancer risk are related to vascular perturbations (ie, mediated by circulating levels of vascular endothelial growth factor) and epithelial-to-mesenchymal transition.¹¹⁵

In contrast to unhealthy dietary patterns, healthy diets were not associated with weight status, suggesting that other factors may play a role in their potential contribution to the development of cancer. Healthy dietary patterns have been associated with higher odds of being more physically active and more highly educated and lower odds of being a smoker. Leisure-time physical activity is associated with an overall lower risk for many types of cancer.¹¹⁶ Smoking status is a known risk factor for several cancers, including lung and colorectal, which were associated with healthy dietary patterns.¹¹⁷ It can be hypothesized that more-educated individuals are more conscious about their health, may be more aware of nutrition-related health issues, and, ultimately, may adopt an overall healthier lifestyle, including not only healthy food choices but also involvement in physical activities and abstinence from smoking.

The main advantage of analyzing *a posteriori*-derived dietary patterns is that dietary habits are evaluated globally.¹¹⁸ Nevertheless, the methods used involve important limitations. Factor analysis typically explains only a small proportion of the variation in diets, and the dietary pattern identified may not fully capture an ideal healthful or unhealthful diet. This is probably less of an issue with healthy dietary patterns because the main characteristics of a healthy diet are more universally recognized and more easily reproduced and compared across a population. In contrast, unhealthy dietary patterns may differ widely from one other, and individual components may have a more harmful effect on specific cancer sites (eg, an unhealthy dietary pattern characterized by processed meat may strongly affect colon cancer risk, or a diet rich in salted foods may be associated with a higher risk of gastric cancer but not other cancers). An attempt was made to test whether each dietary pattern characterized by a main component was associated with increased cancer risk. Subgroup analyses of these patterns resulted in significant findings, despite being affected by substantial heterogeneity. However, none of the patterns, with the exception of the alcohol-related dietary pattern, showed significant results when analysis was limited to prospective studies only. The limited number of studies included in each subgroup analysis besides the animal/meat and Western dietary patterns precluded more

definitive results. When individual cancer sites were analyzed, animal dietary patterns were significantly associated with increased colorectal cancer risk. The current scientific literature shows that heavy alcohol drinking is among the main causes of increased cancer risk, accounting for up to a 5% increased risk of digestive tract and breast cancers.¹¹⁹ Increasing evidence suggests that red and processed meat may directly increase the risk of colorectal cancer.¹²⁰

Regardless of the type of unhealthy dietary pattern adopted within a population, dietary pattern is strongly related to cultural habits and country-specific factors (eg, food availability). Thus, geographical area as a potential source of heterogeneity within the same group of dietary patterns was also assessed. Neither healthy nor unhealthy dietary patterns were unequivocally associated with cancer risk. In fact, the former showed a stronger association with cancer risk in studies conducted in North America and Europe, while the latter showed increased cancer risk only in North American studies, as most of the other evidence was supported only by case-control studies. Possible explanations for the geographical differences may be variation in the amount and proportion of individual unhealthy foods consumed, different food quality between geographical areas, and background characteristics that may interact with dietary habits. Of note, unhealthy dietary patterns were associated with higher BMI and higher cancer risk mainly in studies conducted in North America, which reinforces the hypothesis that BMI may mediate the effect of unhealthy dietary choices. The effect of confounding factors on individual food patterns should be further investigated in future studies, ideally to better characterize the differences between North American and European unhealthy dietary behaviors and to test whether 1 or more discriminant factors exist.

The methodology used to assess *a posteriori*-derived dietary patterns is also affected by other limitations, mainly those related to the arbitrariness of the approach. First, the estimation of dietary patterns depends on the number of items included in the food frequency questionnaires used for data collection.¹²¹ Second, methods to estimate dietary patterns may lack robustness and be poorly reproducible when evaluated in different populations or when based on different numbers of dietary components.¹¹⁸ Third, methodological homogeneity across studies with potential differences in dietary patterns was assumed. Similar subjective names for dietary patterns (eg, Western, animal-based, fat-rich, etc) may have included different components (ie, red meat vs processed meat vs fatty foods), and each component could have been weighted differently in a pattern or, by contrast, could have been represented in more than 1 pattern identified,

thus limiting the comparability of the results. However, besides the effort to group dietary patterns into qualitative groups on the basis of both arbitrary judgment and the nomenclature used by original authors of studies included, a significant level of heterogeneity across studies was found (especially among case-control studies) and impossible to deal with, making generalizability of results rather weak. Other limitations are related to the general study design of some of the studies included in the meta-analyses and comprise recall bias, selection bias, sample size, length of follow-up, and region studied (developed vs nondeveloped countries). Finally, the number of studies and datasets included for extraction, calculation, or imputation of data on food group consumption, energy intake, and lifestyle factors was often too limited to provide representative estimates.

CONCLUSION

The findings obtained from this study may have important public health implications. The evidence presented here suggests that a healthy diet has potential to modulate cancer risk, especially the risk of colon, breast, and lung cancers. In contrast, unhealthy dietary patterns showed a trend of association with increased cancer risk. Healthy dietary patterns have been found to be part of overall healthier lifestyle choices, which may explain in part their association with decreased cancer risk. Unhealthy dietary patterns were associated with a higher BMI, suggesting the potential mediating effect of obesity on cancer risk. Further investigation is needed to confirm the results of this meta-analysis, especially as related to those cancer sites for which evidence was based mainly on case-control studies. Moreover, further study is required to understand the geographical and interindividual variations in diet and lifestyle exposures and their relative contributions to cancer risk.

Acknowledgments

Author contributions. G.G. conceived the study design, performed the study search, performed the analysis, and wrote the manuscript; E.L.G. provided insights on methodology, data interpretation, and manuscript drafting; F.B. and J.G. performed the study search and analysis and edited the tables and figures; S.R., D.D.R., S.S., and F.G. provided critical review/revisions.

Funding/support. No external funds supported this work.

Declaration of interest. The authors have no relevant interests to declare.

Supporting Information

The following Supporting Information is available through the online version of this article at the publisher's website.

5 *Table S1* Main characteristics of studies included in the meta-analysis

Table S2 Number of studies, cases, and controls/exposed individuals included in the meta-analyses

10 *Table S3* Dietary patterns identified in the included studies

Table S4 Subgroup analyses by specific type of unhealthy dietary pattern identified and risk of cancer

15 *Table S5* Subgroup analyses by geographical area for healthy and unhealthy dietary patterns identified and risk of cancer

Figure S1 Funnel plots of studies evaluating the association between healthy dietary patterns and cancer risk

20 *Figure S2* Forest plot of studies evaluating pooled risk ratios (RRs) of colorectal cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

30 *Figure S3* Forest plot of studies evaluating pooled risk ratios (RRs) of lung cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

35 *Figure S4* Forest plot of studies evaluating pooled risk ratios (RRs) of breast cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

40 *Figure S5* Forest plot of studies evaluating pooled risk ratios (RRs) of gastric cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

45 *Figure S6* Forest plot of studies evaluating pooled risk ratios (RRs) of ovarian cancer by adoption of

healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

55 *Figure S7* Forest plot of studies evaluating pooled risk ratios (RRs) of ovarian cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

60 *Figure S8* Forest plot of studies evaluating pooled risk ratios (RRs) of oral/pharyngeal and laryngeal cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

65 *Figure S9* Forest plot of studies evaluating pooled risk ratios (RRs) of esophageal cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

70 *Figure S10* Forest plot of studies evaluating pooled risk ratios (RRs) of endometrial cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

75 *Figure S11* Forest plot of studies evaluating pooled risk ratios (RRs) of prostate cancer by adoption of healthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

80 *Figure S12* Funnel plots of studies evaluating the association between unhealthy dietary patterns and cancer risk

85 *Figure S13* Forest plot of studies evaluating pooled risk ratios (RRs) of colorectal cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds

represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

5 *Figures S14* Forest plot of studies evaluating pooled risk ratios (RRs) of ovarian cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

10 *Figures S15* Forest plot of studies evaluating pooled risk ratios (RRs) of gastric cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

15 *Figure S16* Forest plot of studies evaluating pooled risk ratios (RRs) of pancreatic cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

20 *Figure S17* Forest plot of studies evaluating pooled risk ratios (RRs) of lung cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

25 *Figure S18* Forest plot of studies evaluating pooled risk ratios (RRs) of prostate cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

30 *Figure S19* Forest plot of studies evaluating pooled risk ratios (RRs) of gastric cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

35 *Figure S20* Forest plot of studies evaluating pooled risk ratios (RRs) of endometrial cancer by adoption of unhealthy dietary patterns. Squares

represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

40 *Figure S21* Forest plot of studies evaluating pooled risk ratios (RRs) of esophageal cancer by adoption of unhealthy dietary patterns. Squares represent risk estimates of each study; horizontal line represents 95% confidence intervals (CI); diamonds represent pooled estimates and 95% CIs of risk assessed for the highest vs the lowest category of adherence to the dietary pattern

REFERENCES

1. Fitzmaurice C, Dicker D, Pain A, et al. Global Burden of Disease Cancer Collaboration. The global burden of cancer 2013. *JAMA Oncol.* 2015;1: 505–527. 70
2. Liu X, Wang X, Lin S, et al. Dietary patterns and oesophageal squamous cell carcinoma: a systematic review and meta-analysis. *Br J Cancer.* 2014;110:2785–2795.
3. Bertuccio P, Rosato V, Andreano A, et al. Dietary patterns and gastric cancer risk: a systematic review and meta-analysis. *Ann Oncol.* 2013;24:1450–1458. 75
4. Magalhaes B, Peleteiro B, Lunet N. Dietary patterns and colorectal cancer: systematic review and meta-analysis. *Eur J Cancer Prev.* 2012;21:15–23.
5. Albuquerque RC, Baltar VT, Marchioni DM. Breast cancer and dietary patterns: a systematic review. *Nutr Rev.* 2014;72:1–17.
6. Stroup DF, Berlin JA, Morton SC, et al., for the Meta-analysis of Observational Studies in Epidemiology (MOOSE) Group. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA.* 2000;283:2008–2012. 80
7. Joint WHO/FAO Expert Consultation. Strength of evidence. In: *Diet, Nutrition and the Prevention of Chronic Diseases.* Geneva, Switzerland: World Health Organization. Technical Report Series 916. http://www.who.int/nutrition/topics/5_population_nutrient/en/. Published April 2003. Accessed April 11, 2017. 85
8. Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions.* Version 5.1.0. London, UK: The Cochrane Collaboration; 2011. **AQ17**
9. De Stefani E, Deneo-Pellegrini H, Boffetta P, et al. Dietary patterns and risk of cancer: a factor analysis in Uruguay. *Int J Cancer.* 2009;124:1391–1397. 90
10. Edefonti V, Hashibe M, Ambrogi F, et al. Nutrient-based dietary patterns and the risk of head and neck cancer: a pooled analysis in the International Head and Neck Cancer Epidemiology consortium. *Ann Oncol.* 2012;23:1869–1880.
11. Marchioni DM, Fisberg RM, Francisco de Gois Filho J, et al. Dietary patterns and risk of oral cancer: a case-control study in Sao Paulo, Brazil. *Rev Saude Publica.* 2007;41:19–26. 95
12. Bahmanyar S, Ye W. Dietary patterns and risk of squamous-cell carcinoma and adenocarcinoma of the esophagus and adenocarcinoma of the gastric cardia: a population-based case-control study in Sweden. *Nutr Cancer.* 2006;54:171–178. 100
13. Hajizadeh B, Rashidkhani B, Rad AH, et al. Dietary patterns and risk of oesophageal squamous cell carcinoma: a case-control study. *Public Health Nutr.* 2010;13:1107–1112.
14. Navarro Silvera SA, Mayne ST, Risch HA, et al. Principal component analysis of dietary and lifestyle patterns in relation to risk of subtypes of esophageal and gastric cancer. *Ann Epidemiol.* 2011;21:543–550. 105
15. Bravi F, Edefonti V, Randi G, et al. Dietary patterns and the risk of esophageal cancer. *Ann Oncol.* 2012;23:765–770.
16. Ibiebele TI, Hughes MC, Whiteman DC, et al. Dietary patterns and risk of oesophageal cancers: a population-based case-control study. *Br J Nutr.* 2012;107:1207–1216. 110
17. Sewram V, Sitas F, O'Connell D, et al. Diet and esophageal cancer risk in the Eastern Cape Province of South Africa. *Nutr Cancer.* 2014;66:791–799.
18. Liu X, Wang X, Lin S, et al. Dietary patterns and the risk of esophageal squamous cell carcinoma: a population-based case-control study in a rural population. *Clin Nutr.* 2017;36:260–266. **AQ11**
19. Palli D, Russo A, Decarli A. Dietary patterns, nutrient intake and gastric cancer in a high-risk area of Italy. *Cancer Causes Control.* 2001;12:163–172.
20. Kim MK, Sasaki S, Sasazuki S, et al. Prospective study of three major dietary patterns and risk of gastric cancer in Japan. *Int J Cancer.* 2004;110:435–442.
21. Campbell PT, Sloan M, Kreiger N. Dietary patterns and risk of incident gastric adenocarcinoma. *Am J Epidemiol.* 2008;167:295–304. 120
22. Bertuccio P, Edefonti V, Bravi F, et al. Nutrient dietary patterns and gastric cancer risk in Italy. *Cancer Epidemiol Biomarkers Prev.* 2009;18:2882–2886.

23. Pham TM, Fujino Y, Kikuchi S, et al. Dietary patterns and risk of stomach cancer mortality: the Japan Collaborative Cohort Study. *Ann Epidemiol*. 2010;20:356–363.
24. Denova-Gutierrez E, Hernandez-Ramirez RU, Lopez-Carrillo L. Dietary patterns and gastric cancer risk in Mexico. *Nutr Cancer*. 2014;66:369–376.
25. Slattery ML, Potter JD, Ma KN, et al. Western diet, family history of colorectal cancer, *NA72*, *GSTM-1* and risk of colon cancer. *Cancer Causes Control*. 2000;11:1–8.
26. Terry P, Hu FB, Hansen H, et al. Prospective study of major dietary patterns and colorectal cancer risk in women. *Am J Epidemiol*. 2001;154:1143–1149.
27. Fung T, Hu FB, Fuchs C, et al. Major dietary patterns and the risk of colorectal cancer in women. *Arch Intern Med*. 2003;163:309–314.
28. Dixon LB, Balder HF, Virtanen MJ, et al. Dietary patterns associated with colon and rectal cancer: results from the Dietary Patterns and Cancer (DIETSCAN) Project. *Am J Clin Nutr*. 2004;80:1003–1011.
29. Wu K, Hu FB, Fuchs C, et al. Dietary patterns and risk of colon cancer and adenoma in a cohort of men (United States). *Cancer Causes Control*. 2004;15:853–862.
30. Kim MK, Sasaki S, Otani T, et al. Dietary patterns and subsequent colorectal cancer risk by subsite: a prospective cohort study. *Int J Cancer*. 2005;115:790–798.
31. Nkondjock A, Ghadirian P. Associated nutritional risk of breast and colon cancers: a population-based case-control study in Montreal, Canada. *Cancer Lett*. 2005;223:85–91.
32. Kesse E, Clavel-Chapelon F, Boutron-Ruault MC. Dietary patterns and risk of colorectal tumors: a cohort of French women of the national education system (E3N). *Am J Epidemiol*. 2006;164:1085–1093.
33. Butler LM, Wang R, Koh WP, et al. Prospective study of dietary patterns and colorectal cancer among Singapore Chinese. *Br J Cancer*. 2008;99:1511–1516.
34. Flood A, Rastogi T, Wirfalt E, et al. Dietary patterns as identified by factor analysis and colorectal cancer among middle-aged Americans. *Am J Clin Nutr*. 2008;88:176–184.
35. Satia JA, Tseng M, Galanko JA, et al. Dietary patterns and colon cancer risk in Whites and African Americans in the North Carolina Colon Cancer Study. *Nutr Cancer*. 2009;61:179–193.
36. Williams CD, Satia JA, Adair LS, et al. Dietary patterns, food groups, and rectal cancer risk in Whites and African-Americans. *Cancer Epidemiol Biomarkers Prev*. 2009;18:1552–1561.
37. Bravi F, Edefonti V, Bosetti C, et al. Nutrient dietary patterns and the risk of colorectal cancer: a case-control study from Italy. *Cancer Causes Control*. 2010;21:1911–1918.
38. Kurotani K, Budhathoki S, Joshi AM, et al. Dietary patterns and colorectal cancer in a Japanese population: the Fukuoka Colorectal Cancer Study. *Br J Nutr*. 2010;104:1703–1711.
39. Miller PE, Lazarus P, Lesko SM, et al. Diet index-based and empirically derived dietary patterns are associated with colorectal cancer risk. *J Nutr*. 2010;140:1267–1273.
40. Stefani ED, Deneo-Pellegrini H, Ronco AL, et al. Dietary patterns and risk of colorectal cancer: a factor analysis in Uruguay. *Asian Pacific J Cancer Prev*. 2011;12:753–759.
41. Pou SA, Diaz Mdel P, Osella AR. Applying multilevel model to the relationship of dietary patterns and colorectal cancer: an ongoing case-control study in Cordoba, Argentina. *Eur J Nutr*. 2012;51:755–764.
42. Safari A, Shariff ZM, Kandiah M, et al. Dietary patterns and risk of colorectal cancer in Tehran Province: a case-control study. *BMC Public Health*. 2013;13:222. doi:10.1186/1471-2458-13-222.
43. Kumagai Y, Chou WT, Tomata Y, et al. Dietary patterns and colorectal cancer risk in Japan: the Ohsaki Cohort Study. *Cancer Causes Control*. 2014;25:727–736.
44. Azizi H, Asadollahi K, Davtalab Esmaeili E, et al. Iranian dietary patterns and risk of colorectal cancer. *Health Promot Perspect*. 2015;5:72–80.
45. Chen Z, Wang PP, Woodrow J, et al. Dietary patterns and colorectal cancer: results from a Canadian population-based study. *Nutr J*. 2015;14:8. doi:10.1186/1475-2891-14-8.
46. Michaud DS, Skinner HG, Wu K, et al. Dietary patterns and pancreatic cancer risk in men and women. *J Natl Cancer Inst*. 2005;97:518–524.
47. Nkondjock A, Krewski D, Johnson KC, et al. Dietary patterns and risk of pancreatic cancer. *Int J Cancer*. 2005;114:817–823.
48. Inoue-Choi M, Flood A, Robien K, et al. Nutrients, food groups, dietary patterns, and risk of pancreatic cancer in postmenopausal women. *Cancer Epidemiol Biomarkers Prev*. 2011;20:711–714.
49. Bosetti C, Bravi F, Turati F, et al. Nutrient-based dietary patterns and pancreatic cancer risk. *Ann Epidemiol*. 2013;23:124–128.
50. Chan JM, Gong Z, Holly EA, et al. Dietary patterns and risk of pancreatic cancer in a large population-based case-control study in the San Francisco Bay Area. *Nutr Cancer*. 2013;65:157–164.
51. Zhang W, Xiang YB, Li HL, et al. Vegetable-based dietary pattern and liver cancer risk: results from the Shanghai Women's and Men's Health Studies. *Cancer Sci*. 2013;104:1353–1361.
52. Ronco AL, De Stefani E, Boffetta P, et al. Food patterns and risk of breast cancer: a factor analysis study in Uruguay. *Int J Cancer*. 2006;119:1672–1678.
53. Cui X, Dai Q, Tseng M, et al. Dietary patterns and breast cancer risk in the Shanghai Breast Cancer Study. *Cancer Epidemiol Biomarkers Prev*. 2007;16:1443–1448.
54. Hirose K, Matsuo K, Iwata H, et al. Dietary patterns and the risk of breast cancer in Japanese women. *Cancer Sci*. 2007;98:1431–1438.
55. Edefonti V, Decarli A, La Vecchia C, et al. Nutrient dietary patterns and the risk of breast and ovarian cancers. *Int J Cancer*. 2008;122:609–613.
56. Marchioni DML DLF, Fisberg RM. Dietary patterns and risk of breast cancer: a case-control study in the Northeast of Brazil. *Nutrire Revista de Sociedade Brasileira de Alimentação Nutrição*. 2008;33:31–42.
57. Murtaugh MA, Sweeney C, Giuliano AR, et al. Diet patterns and breast cancer risk in Hispanic and non-Hispanic white women: the Four-Corners Breast Cancer Study. *Am J Clin Nutr*. 2008;87:978–984.
58. Wu AH, Yu MC, Tseng CC, et al. Dietary patterns and breast cancer risk in Asian American women. *Am J Clin Nutr*. 2009;89:1145–1154.
59. Cho YA, Kim J, Shin A, et al. Dietary patterns and breast cancer risk in Korean women. *Nutr Cancer*. 2010;62:1161–1169.
60. Ronco AL, de Stefani E, Aune D, et al. Nutrient patterns and risk of breast cancer in Uruguay. *Asian Pacific J Cancer Prev*. 2010;11:519–524.
61. Ronco AL, De Stefani E, Deneo-Pellegrini H, et al. Dietary patterns and risk of ductal carcinoma of the breast: a factor analysis in Uruguay. *Asian Pacific J Cancer Prev*. 2010;11:1187–1193.
62. Buck K, Vrieling A, Flesch-Janys D, et al. Dietary patterns and the risk of postmenopausal breast cancer in a German case-control study. *Cancer Causes Control*. 2011;22:273–282.
63. Zhang CX, Ho SC, Fu JH, et al. Dietary patterns and breast cancer risk among Chinese women. *Cancer Causes Control*. 2011;22:115–124.
64. Bessaoud F, Tretarre B, Daures JP, et al. Identification of dietary patterns using two statistical approaches and their association with breast cancer risk: a case-control study in southern France. *Ann Epidemiol*. 2012;22:499–510.
65. Demetriou CA, Hadjisavvas A, Loizidou MA, et al. The Mediterranean dietary pattern and breast cancer risk in Greek-Cypriot women: a case-control study. *BMC Cancer*. 2012;12:113. doi:10.1186/1471-2407-12-113.
66. Jordan I, Hebestreit A, Swai B, et al. Dietary patterns and breast cancer risk among women in northern Tanzania: a case-control study. *Eur J Nutr*. 2013;52:905–915.
67. Tumas N, Niclis C, Aballay LR, et al. Traditional dietary pattern of South America is linked to breast cancer: an ongoing case-control study in Argentina. *Eur J Nutr*. 2014;53:557–566.
68. Pot GK, Stephen AM, Dahm CC, et al. Dietary patterns derived with multiple methods from food diaries and breast cancer risk in the UK Dietary Cohort Consortium. *Eur J Clin Nutr*. 2014;68:1353–1358.
69. Sieri S, Krogh V, Pala V, et al. Dietary patterns and risk of breast cancer in the ORDET cohort. *Cancer Epidemiol Biomarkers Prev*. 2004;13:567–572.
70. Adebamowo CA, Hu FB, Cho E, et al. Dietary patterns and the risk of breast cancer. *Ann Epidemiol*. 2005;15:789–795.
71. Fung TT, Hu FB, Holmes MD, et al. Dietary patterns and the risk of postmenopausal breast cancer. *Int J Cancer*. 2005;116:116–121.
72. Mannisto S, Dixon LB, Balder HF, et al. Dietary patterns and breast cancer risk: results from three cohort studies in the DIETSCAN project. *Cancer Causes Control*. 2005;16:725–733.
73. Velie EM, Schairer C, Flood A, et al. Empirically derived dietary patterns and risk of postmenopausal breast cancer in a large prospective cohort study. *Am J Clin Nutr*. 2005;82:1308–1319.
74. Agurs-Collins T, Rosenberg L, Makambi K, et al. Dietary patterns and breast cancer risk in women participating in the Black Women's Health Study. *Am J Clin Nutr*. 2009;90:621–628.
75. Cottet V, Touvier M, Fournier A, et al. Postmenopausal breast cancer risk and dietary patterns in the E3N-EPIC prospective cohort study. *Am J Epidemiol*. 2009;170:1257–1267.
76. Butler LM, Wu AH, Wang R, et al. A vegetable-fruit-soy dietary pattern protects against breast cancer among postmenopausal Singapore Chinese women. *Am J Clin Nutr*. 2010;91:1013–1019.
77. Baglietto L, Krishnan K, Severi G, et al. Dietary patterns and risk of breast cancer. *Br J Cancer*. 2011;104:524–531.
78. Link LB, Canchola AJ, Bernstein L, et al. Dietary patterns and breast cancer risk in the California Teachers Study cohort. *Am J Clin Nutr*. 2013;98:1524–1532.
79. Catsburg C, Kim RS, Kirsh VA, et al. Dietary patterns and breast cancer risk: a study in 2 cohorts. *Am J Clin Nutr*. 2015;101:817–823.
80. Karimi Z, Jessri M, Houshiar-Rad A, et al. Dietary patterns and breast cancer risk among women. *Public Health Nutr*. 2014;17:1098–1106.
81. Biel RK, Friedenreich CM, Cszizmadi I, et al. Case-control study of dietary patterns and endometrial cancer risk. *Nutr Cancer*. 2011;63:673–686.
82. Bravi F, Bertuccio P, Turati F, et al. Nutrient-based dietary patterns and endometrial cancer risk: an Italian case-control study. *Cancer Epidemiol*. 2015;39:66–72.
83. Canchola AJ, Lacey JV Jr, Bernstein L, et al. Dietary patterns and endometrial cancer risk in the California Teachers Study cohort. *Cancer Causes Control*. 2015;26:627–634.

84. Dalvi TB, Canchola AJ, Horn-Ross PL. Dietary patterns, Mediterranean diet, and endometrial cancer risk. *Cancer Causes Control*. 2007;18:957–966.
85. McCann SE, Marshall JR, Brasure JR, et al. Analysis of patterns of food intake in nutritional epidemiology: food classification in principal components analysis and the subsequent impact on estimates for endometrial cancer. *Public Health Nutr*. 2001;4:989–997.
86. Chang ET, Lee VS, Canchola AJ, et al. Dietary patterns and risk of ovarian cancer in the California Teachers Study cohort. *Nutr Cancer*. 2008;60:285–291.
87. Kolahtooz F, Ibiebele TI, van der Pols JC, et al. Dietary patterns and ovarian cancer risk. *Am J Clin Nutr*. 2009;89:297–304.
88. Balder HF, Goldbohm RA, van den Brandt PA. Dietary patterns associated with male lung cancer risk in the Netherlands Cohort Study. *Cancer Epidemiol Biomarkers Prev*. 2005;14:483–490.
89. Gnagnarella P, Maisonneuve P, Bellomi M, et al. Nutrient intake and nutrient patterns and risk of lung cancer among heavy smokers: results from the COSMOS screening study with annual low-dose CT. *Eur J Epidemiol*. 2013;28:503–511.
90. Gorlova OY, Weng SF, Hernandez L, et al. Dietary patterns affect lung cancer risk in never smokers. *Nutr Cancer*. 2011;63:842–849.
91. Clero E, Doyon F, Chungue V, et al. Dietary patterns, goitrogenic food, and thyroid cancer: a case-control study in French Polynesia. *Nutr Cancer*. 2012;64:929–936.
92. Pou SA, Niclis C, Eynard AR, et al. Dietary patterns and risk of urinary tract tumors: a multilevel analysis of individuals in rural and urban contexts. *Eur J Nutr*. 2014;53:1247–1253.
93. Ambrosini GL, Fritschi L, de Klerk NH, et al. Dietary patterns identified using factor analysis and prostate cancer risk: a case control study in Western Australia. *Ann Epidemiol*. 2008;18:364–370.
94. Askari F, Parizi MK, Jessri M, et al. Dietary patterns in relation to prostate cancer in Iranian men: a case-control study. *Asian Pacific J Cancer Prev*. 2014;15:2159–2163.
95. Jackson M, Tulloch-Reid M, Walker S, et al. Dietary patterns as predictors of prostate cancer in Jamaican men. *Nutr Cancer*. 2013;65:367–374.
96. Muller DC, Severi G, Baglietto L, et al. Dietary patterns and prostate cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2009;18:3126–3129.
97. Niclis C, Roman MD, Osella AR, et al. Traditional dietary pattern increases risk of prostate cancer in Argentina: results of a multilevel modeling and bias analysis from a case-control study. *J Cancer Epidemiol*. 2015;2015:179562. doi:10.1155/2015/179562.
98. Rosato V, Edefonti V, Bravi F, et al. Nutrient-based dietary patterns and prostate cancer risk: a case-control study from Italy. *Cancer Causes Control*. 2014;25:525–532.
99. Tseng M, Breslow RA, DeVellis RF, et al. Dietary patterns and prostate cancer risk in the National Health and Nutrition Examination Survey Epidemiological Follow-up Study cohort. *Cancer Epidemiol Biomarkers Prev*. 2004;13:71–77.
100. Walker M, Aronson KJ, King W, et al. Dietary patterns and risk of prostate cancer in Ontario, Canada. *Int J Cancer*. 2005;116:592–598.
101. Wu K, Hu FB, Willett WC, Giovannucci E. Dietary patterns and risk of prostate cancer in U.S. men. *Cancer Epidemiol Biomarkers Prev*. 2006;15:167–171.
102. Millen BE, Abrams S, Adams-Campbell L, et al. The 2015 Dietary Guidelines Advisory Committee Scientific Report: Development and Major Conclusions. *Adv Nutr*. 2016;7:438–444.
103. Giacosa A, Barale R, Bavaresco L, et al. Cancer prevention in Europe: the Mediterranean diet as a protective choice. *Eur J Cancer Prev*. 2013;22:90–95.
104. Grosso G, Buscemi S, Galvano F, et al. Mediterranean diet and cancer: epidemiological evidence and mechanism of selected aspects. *BMC Surg*. 2013;13(suppl 2):S14. doi:10.1186/1471-2482-13-S2-S14.
105. Abid Z, Cross AJ, Sinha R. Meat, dairy, and cancer. *Am J Clin Nutr*. 2014;100(suppl 1):386S–393S.
106. Bagnardi V, Rota M, Botteri E, et al. Alcohol consumption and site-specific cancer risk: a comprehensive dose-response meta-analysis. *Br J Cancer*. 2015;112:580–593.
107. Shield KD, Soerjomataram I, Rehm J. Alcohol use and breast cancer: a critical review. *Alcohol Clin Exp Res*. 2016;40:1166–1181.
108. Baena Ruiz R, Salinas Hernández P. Cancer chemoprevention by dietary phytochemicals: epidemiological evidence. *Maturitas*. 2016;94:13–19.
109. Rondanelli M, Faliva MA, Peroni G, et al. Focus on pivotal role of dietary intake (diet and supplement) and blood levels of tocopherols and tocotrienols in obtaining successful aging. *Int J Mol Sci*. 2015;16:23227–23249.
110. Giovannucci EL. Are most cancers caused by specific risk factors acting on tissues with high underlying stem cell divisions? *J Natl Cancer Inst*. 2015;108:djv343. doi:10.1093/jnci/djv343.
111. Giovannucci E, Michaud D. The role of obesity and related metabolic disturbances in cancers of the colon, prostate, and pancreas. *Gastroenterology*. 2007;132:2208–2225.
112. Renehan AG, Tyson M, Egger M, et al. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*. 2008;371:569–578.
113. Choi Y, Giovannucci E, Lee JE. Glycaemic index and glycaemic load in relation to risk of diabetes-related cancers: a meta-analysis. *Br J Nutr*. 2012;108:1934–1947.
114. Wei T, Ye P, Peng X, et al. Circulating adiponectin levels in various malignancies: an updated meta-analysis of 107 studies. *Oncotarget*. 2016;7:48761–48691.
115. Hursting SD. Obesity, energy balance, and cancer: a mechanistic perspective. *Cancer Treat Res*. 2014;159:21–33.
116. Moore SC, Lee IM, Weiderpass E, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med*. 2016;176:816–825.
117. Gandini S, Botteri E, Iodice S, et al. Tobacco smoking and cancer: a meta-analysis. *Int J Cancer*. 2008;122:155–164.
118. Edefonti V, Randi G, La Vecchia C, et al. Dietary patterns and breast cancer: a review with focus on methodological issues. *Nutr Rev*. 2009;67:297–314.
119. Giacosa A, Adam-Blondon AF, Baer-Sinnott S, et al. Alcohol and wine in relation to cancer and other diseases. *Eur J Cancer Prev*. 2012;21:103–108.
120. Bouvard V, Loomis D, Guyton KZ, et al. Carcinogenicity of consumption of red and processed meat. *Lancet Oncol*. 2015;16:1599–1600.
121. Moeller SM, Reedy J, Millen AE, et al. Dietary patterns: challenges and opportunities in dietary patterns research. *J Am Diet Assoc*. 2007;107:1233–1239.