

University of East London Institutional Repository: <http://roar.uel.ac.uk>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

**Author(s):** Jones, Edward., Adcock, Ian M., Ahmed, Bushra Y., Punchard, Neville A.

**Article Title:** Modulation of LPS stimulated NF-kappaB mediated Nitric Oxide production by PKC $\epsilon$  and JAK2 in RAW macrophages

**Year of publication:** 2007

**Citation:** Jones, E. et al (2007) 'Modulation of LPS stimulated NF-kappaB mediated Nitric Oxide production by PKC $\epsilon$  and JAK2 in RAW macrophages.' Journal of Inflammation 4:23

**Link to published version:**

<http://www.journal-inflammation.com/content/4/1/23>

**DOI:** 10.1186/1476-9255-4-23

**Publisher statement:**

<http://www.biomedcentral.com/info/about/license>

Research

Open Access

## Modulation of LPS stimulated NF- $\kappa$ B mediated Nitric Oxide production by PKC $\epsilon$ and JAK2 in RAW macrophages

Edward Jones<sup>1</sup>, Ian M Adcock<sup>2</sup>, Bushra Y Ahmed<sup>1</sup> and Neville A Punchard<sup>\*3</sup>

Address: <sup>1</sup>Division of Science, University of Luton, Luton. UK, <sup>2</sup>Airway Diseases, NHLI, Imperial College London, London. UK and <sup>3</sup>School of Health & Biosciences, University of East London, London. UK

Email: Edward Jones - edward\_r\_jones@hotmail.co.uk; Ian M Adcock - ian.adcock@imperial.ac.uk; Bushra Y Ahmed - Bushra.ahmed@beds.ac.uk; Neville A Punchard\* - n.punchard@uel.ac.uk

\* Corresponding author

Published: 24 November 2007

Received: 16 April 2007

*Journal of Inflammation* 2007, 4:23 doi:10.1186/1476-9255-4-23

Accepted: 24 November 2007

This article is available from: <http://www.journal-inflammation.com/content/4/1/23>

© 2007 Jones et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Nuclear factor kappa B (NF- $\kappa$ B) has been shown to play an important role in regulating the expression of many genes involved in cell survival, immunity and in the inflammatory processes. NF- $\kappa$ B activation upregulates inducible nitric oxide synthase leading to enhanced nitric oxide production during an inflammatory response. NF- $\kappa$ B activation is regulated by distinct kinase pathways independent of inhibitor of  $\kappa$ B kinase (IKK). Here, we examine the role of protein kinase C isoforms and janus activated kinase 2 (JAK2) activation in NF- $\kappa$ B activation and LPS-stimulated NO production.

**Methods:** Murine RAW 264.7 macrophages were treated with lipopolysaccharide (LPS), Phorbol 12-myristate 13-acetate (PMA) and a combination of LPS and PMA in the presence or absence of various inhibitors of PKC isoforms and JAK2. Nuclear translocation of the NF- $\kappa$ B p65 subunit, was assessed by Western blot analysis whilst NO levels were assessed by Greiss assay.

**Results:** LPS-stimulated NO production was attenuated by PMA whilst PMA alone did not affect NO release. These effects were associated with changes in p65 nuclear translocation. The PKC $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  and  $\zeta$  inhibitor Gö 6983 (Go) had no effect on LPS-induced NO release. In contrast, Bisindolymaleimide I (Bis), a PKC  $\alpha$ ,  $\beta$ <sub>I</sub>,  $\beta$ <sub>II</sub>,  $\gamma$ ,  $\delta$  and  $\epsilon$  isoform inhibitors completely inhibited LPS-stimulated NO production without affecting p65 nuclear translocation. Furthermore, a partial inhibitory effect on LPS-induced NO release was seen with the JAK2 inhibitor AG-490 and the p38 MAPK inhibitor SB 203850.

**Conclusion:** The results further define the role of NF- $\kappa$ B in LPS stimulated NO production in RAW macrophages. The data support a function for PKC $\epsilon$ , JAK2 and p38 MAPK in NF- $\kappa$ B activation following p65 nuclear import.

## Introduction

Increasing emphasis is currently placed on the role of the innate immune system in inflammatory responses, in particular those involving macrophages. As in other cells, the transcription factor, NF- $\kappa$ B plays a pivotal role in changes in gene expression during such inflammatory responses. A range of inflammatory stimuli, including endotoxin [1,2] and cytokines [3], produce activation and nuclear translocation of NF $\kappa$ B following rapid degradation and release of I $\kappa$ B.

One of the genes upregulated by NF- $\kappa$ B during an inflammatory response is the inducible nitric oxide synthase (NOS2), that produces nitric oxide (NO), a highly reactive free radical with important second messenger functions involving the mediation of inflammatory events [4]. Increased expression of NOS2 and concomitant NO levels have been reported in several inflammatory diseases, such as Crohn's disease [5], asthma [6] and rheumatoid arthritis [7]. The NOS2 gene promoter contains twenty two putative transcription factor binding elements [8], however, so far only the NF- $\kappa$ B responsive  $\kappa$ B element [9] and an interferon- $\gamma$ -activated site (GAS) [10] have been shown to enhance NOS2 expression.

Macrophages are the primary producers of NO *in vivo*

[1]. Lipopolysaccharide (LPS) stimulates NO production in macrophages. The induction of NOS2 protein expression in response to stimulation with LPS involves the Janus kinase (JAK) family of protein kinases [11]. Furthermore, both protein kinase C (PKC) and Janus Kinase2 (JAK2) [12-14] have been implicated in NF- $\kappa$ B activation. However, although nine isoforms of PKC have been identified in macrophages [15] it is unknown which of these are involved in NF- $\kappa$ B activation.

PKC activation has been identified as an early response in LPS-stimulated macrophages [16] and is essential for the up-regulation of NO production [16,17] However, the function of PKC isoforms involved in upregulation of NO production remains to be determined. Thus PMA, a direct activator of the PKC family of kinases, was used to investigate the role of PKC in LPS-stimulated NO production and NF- $\kappa$ B activation in RAW cells. PMA has been shown to induce a PKC mediated proteasomal-independent pathway of NF- $\kappa$ B nuclear translocation in human intestinal epithelial cells [18].

The present study uses pharmacological tools to indicate a role for PKC $\epsilon$  in LPS-stimulated NF- $\kappa$ B-mediated NO release in RAW macrophages. We also implicate a role for JAK2 and p38 MAPK on these effects.

## Methods

### Cell Culture

RAW 264.7 cells (ECACC, Salisbury, UK) were maintained in 25 cm<sup>2</sup> flasks in DMEM medium supplemented with 2 mM L-glutamine and 10% v/v FCS, without antibiotics, at 37°C in a humidified atmosphere of 95% air and 5% CO<sub>2</sub>. For Western blotting, cells were grown in 25 cm<sup>2</sup> flasks, whilst for the measurement of NO the cells were grown to 95% confluence in 96-well plates and stimulation carried out within these plates. Cells were stimulated by replacing the culture medium with medium containing LPS, LPS with phorbol-12-myristate-13-acetate (PMA) or PMA alone in the presence or absence of various inhibitors.

Inhibitors used were: the PKC inhibitors Gö 6983 (Go) and Bisindolymaleimide I (Bis); the JAK2 inhibitor AG-490; the p38 MAP kinase inhibitor SB 203580 (Calbiochem, Nottingham, UK). Bis shows high selectivity for PKC  $\alpha$ ,  $\beta$ <sub>I</sub>,  $\beta$ <sub>II</sub>,  $\gamma$ ,  $\delta$  and  $\epsilon$  isoforms at 20  $\mu$ M [19] whilst Go inhibits PKC  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  and  $\zeta$  isoforms at 10  $\mu$ M [20]. AG-490 was used at 10 mM, a concentration previously shown to inhibit JAK2 [21], and SB 204580 at 10 mM, a concentration previously shown to inhibit the p68 MAP kinase family [22]. For the purpose of specific inhibition of PKC translocation, the following MALY-TAT linked peptides (kindly supplied by Dr M. Lindsay, AstraZeneca, Charnwood, UK) were used: MALY01 (TAT-RFARKGAL-RQKNHEVK), MALY10 (TAT-EAVSLKPT), MALY II (TAT-LSETKPAVO) at concentrations previously shown to inhibit translocation of PKC isoforms [23]. For Western blotting cells were incubated for 0, 1, 2, 3 or 5 hours, whilst for the NO assay, cells were incubated for 24 hours.

### Assessment of NF- $\kappa$ B-p65 nuclear translocation by Western blot analysis

RAW 264.7 cells were harvested in ice cold PBS after stimulation with LPS from 0 to 5 h. Cells were then lysed in 70  $\mu$ l of buffer A (10 mM HEPES pH 7.9, 1.5 mM MgCl<sub>2</sub>, 10 mM KCl, 0.25% v/v noident P-40, 0.5 mM dithiothreitol (DTT), 0.5 mM phenylmethylsulfonyl fluoride (PMSF) in de-ionised water (dH<sub>2</sub>O) for 20 min on ice, to yield the cytoplasmic cellular fraction, as described previously [24]. The samples were microfuged at 12,000 g for 15 sec to pellet the unlysed nuclei and the supernatant (cytoplasmic fraction) was collected. The nuclei were lysed in 15  $\mu$ l of buffer B (20 mM HEPES pH 7.9, 1.5 mM MgCl<sub>2</sub>, 0.42 M NaCl, 0.5 mM DTT, 25% v/v Glycerol, 0.5 mM PMSF in diH<sub>2</sub>O) for 20 min on ice, and microfuged or centrifuged at 12,000 g for 60 sec to pellet the cellular debris. The supernatant (nuclear fraction) was collected and 60  $\mu$ l of buffer C (20 mM HEPES pH 7.9, 50 mM KCl, 0.5 mM DTT, 0.2 mM EDTA, 0.5 mM PMSF) was added to it. At this stage the protein concentration of the samples was assessed by BioRad protein assay™ (Biorad, UK).

Samples (10 µg) were separated by 10% SDS-PAGE and proteins transferred to a nitrocellulose membrane (Amersham-Pharmacia, Amersham, UK) by electroblotting. Equal protein loading was confirmed by Ponceau S staining of the membrane. Non-specific protein binding was blocked by incubation of the membrane in PBS-T + 1% w/v milk overnight at 4°C. Membranes were then washed twice in PBS-T for 5 min before incubation for 1 hour at room temperature (RT) with rabbit anti-p65 antibody (1:4000, Santa Cruz, Wembley, UK). Membranes were then washed twice in phosphate buffered saline (pH 7.4) containing 0.05% v/v Tween 20 (PBS-T) and 1% w/v milk for 5 min followed by an hour incubation at room temperature (RT) with goat anti-rabbit HRP conjugate (1:4000, Dako, UK). All antibodies incubations were carried out in PBS-T containing 1% w/v milk. Membranes were washed three times for 5 min in PBS-T before incubation with ECL substrate (Amersham-Pharmacia, UK), followed by exposure to an autoradiographic film and subsequent semi-quantification of band intensity by densitometry (UVP Ltd, Cambridge, UK).

#### **Nitrite determination by Greiss assay**

NO levels were assessed by nitrite quantification as described previously [25]. Briefly, 90 µl of sample (cell culture medium) was incubated for 5 min in dark at RT with 90 µl of suphanilamide (1% w/v in 4 M HCl). 90 µl of naphthylethylenediamine (1% w/v in dH<sub>2</sub>O) was then added and a further 5 min incubation was carried out in dark at RT. Absorbance was read at 540 nm.

All reagents were purchased from Sigma (Poole, UK) unless otherwise stated above.

#### **Statistical analysis**

Data are reported as mean ± SEM. Statistical analysis was performed in Prism 5 (Graph Pad Software, Inc. San Diego, USA) using one way analysis of variance (ANOVA) followed by Tukey's Multiple Comparison Test (TMCT) when ANOVA indicated a statistical significance existed.

## **Results**

#### **The effect of LPS and PMA on RAW cell NO production**

Constitutive NO production by RAW 264.7 macrophages (Fig. 1) was at the lower limit of the sensitivity of the assay (3 µM). LPS induced a concentration-dependent increase in NO production with a maximum response at 50 µg/ml at 24 h (Figure 1b.). For subsequent experiments, LPS (1 µg/ml) was chosen for its ability to stimulate high levels of NO production whilst not possessing the significant ( $p < 0.001$ ) toxicity seen with 10 and 50 µg/ml (38% and 51% reduction in viability respectively).

LPS stimulation induced NO production with nitrite levels peaking at 33 µM ( $p < 0.001$ , Fig. 1), in agreement with

the results of others [26,27]. In contrast, PMA alone (0.5–500 ng/ml) had no effect on NO production but significantly attenuated LPS-induced NO production by ~50% (Fig. 1) even at concentrations (50 ng/ml) previously shown to activate NF-κB.

#### **The effect of LPS on NF-κB activation**

LPS (1 µg/ml) induced a significant 4-fold induction of p65 nuclear translocation which was maintained for up to 5 hours ( $p < 0.05$ , Fig. 2). PMA alone also significantly induced NF-κB activation (data not shown), however, combined effect of PMA and LPS-stimulated NF-κB showed an 8-fold increase within 30 minutes PMA reduced the duration of LPS-stimulated p65 nuclear translocation from > 5 to less than 2 hours (Fig. 3).


#### **The role of PKC in LPS stimulated RAW cell NO production**

Effects on PKC isoforms degradation have been reported following PMA treatment at the concentration of PMA used in this study (50 ng/ml) [28]. The induction of NOS2 activity upon activation of PKC by LPS stimulation has also been demonstrated previously using a non-selective inhibitor of all PKC isoforms [16]. However, the role of specific PKC isoforms has remained unclear. The MALY peptides have been reported to mimic the PKC variable regions 1 and 2 (V1-2). These regions are necessary for binding PKC to the receptors for activated C kinase (RACK) and thereby prevent nuclear-cytoplasmic translocation of specific PKC isoforms [29]. The addition of a TAT sequence (GGGGYGRKKRRQRRR-GGGG) to the MALY peptides ensures that they are transported into the nucleus, an important facet for some PKC enzymes. Pre-treatment with these translocation inhibitor-peptides had no effect on NO production in these cells (Table 1). In addition, the PKC inhibitor Gö 6983 (Go, 10 µM) had no effect on LPS-stimulated NO production (Fig. 4).

In contrast, bisindolymaleimide I (Bis, 20 µM) completely inhibited LPS (1 µg/ml) -stimulated NO production (Fig. 4). This was not due to an inhibitory effect on peak NF-κB p65 nuclear translocation as Bis (20 µM) had no effect on LPS-stimulated nuclear translocation at 3 hr (Fig. 5). This time point was chosen as it represented the time at which NF-κB p65 nuclear translocation was at its peak (Fig. 2). The results also contrast with the effect of PMA which returned p65 nuclear translocation to baseline within 2 hr.

#### **The effect of AG-490 on LPS induced RAW cell NO production**

AG-490, a potent inhibitor of JAK2, caused a concentration-dependent inhibition of LPS-stimulated NO production and this was significant at 10 µM (Fig. 6). JAK2 has been shown to be activated in response to a wide variety

**Figure 1**

**a. The effect of PMA on LPS stimulated NO production in RAW 264.7 cells.** RAW cells were stimulated for 24 hr with either vehicle alone (control), 1  $\mu\text{g/ml}$  LPS, 50 ng/ml PMA or 1  $\mu\text{g/ml}$  LPS with 50 ng/ml PMA. NO levels were assessed by Greiss assay. LPS alone significantly increased NO production whereas PMA alone had no effect. PMA significantly inhibited LPS-stimulated NO production. **Figure 1b** (inset) displays the concentration-response curve for LPS-stimulated NO production over 24 hrs. Results are expressed as mean  $\pm$  SEM; <sup>+++</sup> $p < 0.001$  vs control, <sup>\*\*\*</sup> $p < 0.001$  vs LPS-stimulated;  $n = 6$  for the effects of PMA and  $n = 9$  for the concentration-response curve.

of stimuli [30] and AG-490 to block induced NOS2 expression in IL-1 $\beta$ /TNF $\alpha$ /IFN $\gamma$ -stimulated human epithelial-like colon carcinoma DLD-1 cells [31] and in IFN $\gamma$ /LPS stimulated RAW cells [32]. This present finding extends the previous studies and further indicates the involvement of JAK2 in NF- $\kappa$ B-induced LPS-stimulated NO production and NF- $\kappa$ B activation in RAW cells.


#### **The effect of SB 203580-induced inhibition of p38 on LPS induced RAW cell NO production**

p38 MAP kinase has previously been implicated in NF- $\kappa$ B activation [33] and in IFN $\gamma$ /LPS stimulated NOS2 expression in RAW 264.7 cell, although this has not been reported for LPS-stimulated cells alone [34]. SB203580 inhibited LPS-stimulated NO production in a concentration-dependent manner with an  $\text{IC}_{50}$  of  $\sim 3 \mu\text{M}$  indicating selectivity of this effect. Maximal inhibition ( $\sim 35\%$ ) was seen at 10  $\mu\text{M}$  (Fig. 7).

#### **Discussion**

This study investigated the effect of distinct kinase pathways on their ability to modulate NF- $\kappa$ B activation and thereby modify LPS stimulated NO production in RAW macrophages. Previous findings have been extended, with NF- $\kappa$ B-mediated NO production stimulated by LPS is shown to be multifactorial in nature, involving the coordinated activation of PKC $\epsilon$ , JAK2 and p38 MAPK. The differential effects of the PKC inhibitors Go and Bis suggested that PKC $\epsilon$  was involved in NO release.


Under unstimulated conditions, p65 is restricted to the cytoplasm by a set of inhibitory proteins and, upon stimulation, translocated to the nucleus. This stimulation can be modulated by phosphorylation of p65 at serine residues. The degree of activation by NF- $\kappa$ B is thus likely to result from a combination of p65 nuclear translocation and post-translational modifications of p65 [35].



**Figure 2**  
**NF-κB activation in LPS stimulated RAW 264.7 cells.** RAW cells were stimulated with LPS (1 μg/ml) for between 0–5 hours. Cells were harvested and nuclear NF-κB-p65 levels were assessed by Western blotting. (a) Representative Western blot of p65 expression. Equal amounts of nuclear proteins are loaded onto each lane. (b) Graphical representation of % increase in p65 nuclear localisation shown in (a) above. Data is presented as mean ± sem, \**p* < 0.05, *n* = 6 independent measurements.

It is evident from many studies that LPS-stimulated NO release from RAW macrophages is NF-κB dependent [36]. In the present study, although PMA enhanced the amount of p65 nuclear translocation, it also decreased the period over which LPS was able to maintain NF-κB nuclear translocation and this may be linked to the reduction in NO release.

Inhibition of PKCε by Bis had no effect on nuclear translocation in our study confirming a previous report in LPS-stimulated blood monocytes [37]. This data, in conjunction with the data showing a lack of effect of a TAT-linked MALY inhibitor, indicates that PKC is not involved in the nuclear translocation or DNA-binding of NF-κB. Thus, PKCε probably acts on nuclear NF-κB to either affect its nuclear retention or more likely to affect p65 transcriptional activity through a posttranslational modification event leading to differential recruitment or activation of transcriptional co-activators. Indeed, a PKC phosphorylation site exists on the p65 subunit and such phosphorylation is known to increase the transactivation potential of NF-κB without affecting its DNA binding or nuclear translocation [38,39]. Furthermore, the data presented here suggests that the effect of PMA on LPS-stimulated NO release is not through a PKC-mediated effect but that PMA induces additional pathways that regulate LPS-induced



**Figure 3**  
**NF-κB activation in LPS and PMA stimulated RAW 264.7 cells.** RAW cells were stimulated with LPS (1 μg/ml) and PMA (50 ng/ml) for between 0–120 minutes. Cells were harvested and nuclear NF-κB-p65 levels were assessed by Western blotting. (a) Representative Western blot of p65 expression. Equal amounts of nuclear proteins were loaded onto each lane. (b) Graphical representation of % increase in p65 nuclear localisation shown in (a) above. Data is presented as mean of 2 independent experiments.

NF-κB activation and NO production. Thus, altered PKC activity may also impinge upon the NF-κB functional response either by affecting co-factor or histone phosphorylation [40].

A role for JAK2 in LPS and IFN stimulated NO production in RAW cells has been described previously [41]. However, JAK2 was hypothesised to work solely through STAT1 activation and be activated by IFNγ. There is currently growing evidence for cross talk between the JAK2 and the NF-κB signalling pathway [42] and also the JNK pathway indirectly through an effect on PI3K [43,44]. JAK2 has been demonstrated to phosphorylate IκB thereby facilitating NF-κB activation [14]. AG-490 has also been reported to inhibit LPS stimulated NF-κB activation and subsequent NOS2 induction in a skin dendritic cell line [45].

As with JAK2, the results from the present study suggest that the p38 MAPK protein (MAP) kinase is also involved in NF-κB activation [29,33] and therefore LPS-induced NO production. Previous reports have shown equivocal data as to the role of p38 MAPK in these events [46,47]. Although p38 MAPK does not appear to be involved in NO release induced by other agents in RAW cells [48]. The IC<sub>50</sub> of inhibition of SB203580 indicates relative selectiv-

**Table 1:**



	Nitrite ( $\mu\text{M}$ ); mean $\pm$ SEM						
	Control	Concentration of peptide alone		1 $\mu\text{g/ml}$ LPS with stated concentration of peptides below			
		10 $\mu\text{M}$	100 $\mu\text{M}$	0 $\mu\text{M}$	1 $\mu\text{M}$	10 $\mu\text{M}$	100 $\mu\text{M}$
MALY OI	0.70 $\pm$ 0.2	2.00 $\pm$ 1.6	0.77 $\pm$ 0.3	44.6 $\pm$ 2.7	46.1 $\pm$ 4.2	47.2 $\pm$ 5.1	51.6 $\pm$ 2.0
MALY IO	0.70 $\pm$ 0.2	0.87 $\pm$ 0.4	0.85 $\pm$ 0.3	44.6 $\pm$ 3.3	47.3 $\pm$ 0.8	40.3 $\pm$ 1.9	46.4 $\pm$ 0.9
MALY II	0.70 $\pm$ 0.2	0.64 $\pm$ 0.3	0.66 $\pm$ 0.3	44.6 $\pm$ 2.8	46.2 $\pm$ 0.9	43.4 $\pm$ 2.9	45.9 $\pm$ 0.5

RAW cells were stimulated with medium alone (control), LPS (1  $\mu\text{g/ml}$ ) or various concentrations of the MALY TAT-linked PKC translocation inhibitor peptides (MALYOI = cPKC inhibitor, MALYIO = PKC specific, MALYII = scrambled PKC inhibitor peptide), with or without LPS for 24 hr. NO production was assessed by Greiss assay. Results represent mean  $\pm$  SEM (n= 3) and are representative of two separate experiments.

ity and further analysis of LPS-induced p38 MAPK activation or the use of more selective inhibitors may provide additional evidence for its role effects in these cells and for p38 inhibitors potential in the treatment of LPS activated disease.


In conclusion, the findings of the present study demonstrate the role of NF- $\kappa$ B in LPS stimulated NO production in RAW cells and indicate the importance of cross-talk

with other kinase pathways, namely PKC $\epsilon$ . Furthermore, the present findings further define the involvement of PKC $\epsilon$  and JAK2 in inducing NO production, probably through their effects on NF- $\kappa$ B induced NOS2 expression. Figure 8 provides a pictorial summary of these findings. However, these conclusions have been drawn on the basis of use of well characterised inhibitors, rather than actual measurement of the activity of the target proteins which




**Figure 5**  
**Effect of bisindolylemaleimide (Bis) on LPS stimulated NF- $\kappa$ B activation in RAW 264.7 cells.** RAW cells were stimulated with LPS (1  $\mu\text{g/ml}$ ) in the presence or absence of Bis (20  $\mu\text{M}$ ) for 3 hr. Cells were harvested and nuclear NF- $\kappa$ B-p65 levels were assessed by Western blotting. (a) Representative Western blot of p65 expression. Equal amounts of nuclear proteins were loaded onto each lane. (b) Graphical representation of % increase in p65 nuclear localisation is shown in (a) above. Data is presented as mean  $\pm$  SEM, n = 3 independent measurements.


**Effect of PKC inhibitors on LPS induced NO production in RAW 264.7 cells.** Cells were treated with vehicle (control) or LPS (1  $\mu\text{g/ml}$ ) in the presence or absence of PKC inhibitors Go 6978 (Go, 10  $\mu\text{M}$ ) or bisindolylemaleimide (Bis, 20  $\mu\text{M}$ ) for 24 hours. The culture medium was then harvested and assayed for nitrite content by Greiss assay. The data show that only Bis was able to inhibit LPS-stimulated RAW cell NO production. Results are expressed mean  $\pm$  SEM, \*\*\*p < 0.001, n = 9.



**Figure 6**  
**Effect of JAK2 inhibitor on LPS-stimulated NO production in RAW 264.7 cells.** Concentration dependent effect of the JAK2 inhibitor AG-490 (0–10 µM) on LPS (1 µg/ml)-stimulated NO production measured at 24 hr. Cells were stimulated and the culture medium harvested and assayed for nitrite content by Greiss assay. The data show that AG-490 was able to inhibit LPS-stimulated NO production. Results for the effects of the inhibitor are expressed mean ± SEM, \*\*p < 0.01 vs no inhibitor, n = 6.



**Figure 7**  
**Effect of SB203580 on LPS stimulated NO production.** Concentration dependent effect of the p38 MAPK inhibitor SB203580 (0–10 µM) on LPS (1 µg/ml)-stimulated NO production measured at 24 hours. Cells were stimulated and the culture medium harvested and assayed for nitrite content by Greiss assay. The data show that SB203580 was able to inhibit LPS-stimulated NO production. Results for the effects of the inhibitor are expressed mean ± SEM, \*\*\*p < 0.001 vs no inhibitor, n = 6.



**Figure 8**  
 Diagram showing a pictorial representation of the conclusions from all experiments (? = areas of uncertainty).



would provide further confirmation of this regulatory network. The work presented here further illustrates the complex network of signalling pathways involved in modulation of NF- $\kappa$ B-mediated gene transcription.

### Competing interests

The author(s) declare that they have no competing interests.

### Authors' contributions

EJ carried out the cell culture experiments, molecular studies and analysis and presentation of the results and the initial interpretation. IA provided the training and expertise for the molecular studies, jointly conceived the study, participated in its design and coordination and significantly contributed to the drafting of the manuscript. BA contributed expertise and critical knowledge of the molecular studies and redrafted and formatted the early draft of the manuscript. NP initiated the project, jointly conceived the study, raised the initial funding, provided training in cell culture techniques, supervised the work of EJ and produced the final drafts of the manuscript. All authors read and approved the final manuscript.

### Acknowledgements

This work was funded by the University of Bedfordshire and GlaxoSmithKline (UK). The MALY-TAT linked peptides were kindly supplied by Dr M. Lindsay, AstraZeneca, Charnwood, UK.

### References

- Zhang G, Ghosh S: **Toll-like receptor-mediated NF-kappaB activation: a phylogenetically conserved paradigm in innate immunity.** *J Clin Invest* 2001, **107**:13-9.
- Collins T, Read MA, Neish AS, Whitley MZ, Thanos D, Maniatis T: **Review: Transcriptional regulation of endothelial cell adhesion molecules: NF-kappaB and cytokine-inducible enhancers.** *FASEB* 1995, **9**:899-909.
- Jobin C, Hellerbrand C, Licato LL, Brenner DA, Sartor RB: **Mediation by NF-kappaB of cytokine induced expression of intercellular adhesion molecule 1 (ICAM-1) in an intestinal epithelial cell line, a process blocked by proteasome inhibitors.** *Gut* 1998, **42**:779-787.
- McCafferty DM: **Peroxyinitrite and inflammatory bowel disease.** *Gut* 2000, **46**:436-439.
- Boughton-Smith NK, Evans SM, Hawkey CJ, Cole AT, Balsitis M, Whittle BJ, Moncada S: **Nitric oxide synthase activity in ulcerative colitis and Crohn's disease.** *Lancet* 1993, **342**:338-340.
- Kharitinov SA, Barnes JP: **Effects of corticosteroids on noninvasive biomarkers of inflammation in asthma and chronic obstructive pulmonary disease.** *Proc Am Thorac Soc* 2004, **1**:191-199.
- Yasuda T, Kakinuma T, Julovi SM, Hiramatsu T, Akiyoshi M, Nakamura T: **COOH-terminal heparin-binding fibronectin fragment induces nitric oxide production in rheumatoid cartilage through CD44.** *Rheumatology* 2004, **43**:1116-1120.
- Xie Q-W, Kashiwabara Y, Nathan C: **Role of transcription factor NF-kappaB/Rel in induction of nitric oxide synthase.** *J Biol Chem* 1994, **269**:4705-4708.
- Kim Y-M, Lee S-B, Yi K-Y, Palk S-G: **Upstream NF-kappaB site is required for the maximal expression of mouse inducible nitric oxide synthase gene in interferon-g plus lipopolysaccharide-induced RAW 264.7 macrophages.** *Biochem Biophys Res Comm* 1997, **236**:655-660.
- MacMicking J, Xie Q-W, Nathan C: **Nitric oxide and macrophage function.** *Annu Rev Immunol* 1997, **15**:323-350.
- Ilhe JN: **Cytokine receptor signalling.** *Nature* 1995, **377**:591-594.
- Steffan NM, Bren GD, Frantz B, Tocci MJ, O'Neill EA, Paya CV: **Regulation of I $\kappa$ B phosphorylation by PKC and Ca<sup>2+</sup> dependent signal transduction pathways.** *J Immunol* 1995, **155**:4685-4691.
- Cataisson C, Pearson AJ, Torgerson S, Nedospasov SA, Yuspa ST: **Protein kinase C $\alpha$ -mediated chemotaxis of neutrophils requires NF-kB activity but is independent of TNF $\alpha$  signalling in mouse skin in vivo.** *J Immunol* 2005, **174**:1686-1692.
- Digicaylioglu M, Lipton SA: **Erythropoietin-mediated neuroprotection involves cross-talk between JAK2 and NF $\kappa$ B signalling cascades.** *Nature* 2001, **412**:641-647.
- Lin W-W, Chen BC: **Distinct isoforms mediate the activation of cPLA2 and adenylyl cyclase by phorbol ester in RAW 264.7 macrophages.** *Brit J Pharmacol* 1998, **125**:1601-1609.
- Paul A, Pendreigh RH, Plevin R: **Protein kinase C and tyrosine kinase pathways regulate lipopolysaccharide-induced nitric oxide synthase activity in RAW 264.7 macrophages.** *Brit J Pharmacol* 1995, **114**:482-488.
- Severn S, Wakelam MJO, Liew FY: **The role of PKC in the induction of nitric oxide synthase by murine macrophages.** *Biochem Biophys Res Comm* 1993, **188**:199-1002.
- Sun Y, Oberley L: **Redox regulation of transcriptional activators.** *Free Radic Biol Med* 1996, **21**:335-348.
- Ku WC, Cheng AJ, Wang TC: **Inhibition of telomerase activity by PKC inhibitors in human nasopharyngeal cancer cells in culture.** *Biochem Biophys Res Comm* 1997, **241**:730-6.
- Gschwendt M, Dieterich S, Rennecke J, Kittstein W, Mueller HJ, Johannes FJ: **Inhibition of protein kinase C mu by various inhibitors. Differentiation from protein kinase C isoenzymes.** *FEBS Letts* 1996, **392**:77-80.
- Levitzi A: **Tyrphostins - potential antiproliferative agents and novel molecular tools.** *Biochem Pharmacol* 1990, **40**:913-918.
- Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Heys JR, Landvatter SV, et al.: **A protein kinase involved in the regulation of inflammatory cytokine biosynthesis.** *Nature* 1994, **372**:739-746.
- Yedovitzky M, Mochly-Rosen D, Johnson JA, Gray MO, Ron D, Abramovitch E, Cerasi E, Neshor R: **Translocation inhibitors define specificity of protein kinase C isoenzymes in pancreatic beta-cells.** *J Biol Chem* 1997, **272**:1417-1420.
- Osborn L, Kunel S, Nabel GJ: **Tumour necrosis factor- $\alpha$  and interleukin-1 stimulate the human immunodeficiency virus enhancer by activation of NF-kappaB.** *Proc Natl Acad Sci* 1989, **86**:2336-2340.
- Tsai SH, Lin-Shiau SY, Lin JK: **Suppression of nitric oxide synthase and the down-regulation of the activation of NF-kappaB in macrophages by resveratrol.** *J Pharmacol* 1999, **126**:673-680.
- Paul A, Bryant C, Lawson MF, Chilvers ER, Plevin R: **Dissociation of lipopolysaccharide-mediated induction of nitric oxide synthase and inhibition of DNA synthesis in RAW 264.7 macrophages and rat aortic smooth muscle cells.** *Brit J Pharmacol* 1997, **120**:1439-1444.
- Patel R, Attur MG, Dave MN, Kumar S, Lee JC, Abramson SB, Amin AR: **Regulation of nitric oxide and prostaglandin E<sub>2</sub> production by CSAIDS (SB203580) in murine macrophages and bovine chondrocytes stimulated with LPS.** *Inflamm Res* 1999, **48**:337-43.
- Paul A, Doherty K, Plevin R: **Differential regulation by protein kinase C isoforms of nitric oxide synthase induction in RAW 264.7 macrophages and rat aortic smooth muscle cells.** *Brit J Pharmacol* 1997, **120**:940-946.
- Mayne GC, Murray AW: **Evidence that protein kinase C mediates phorbol ester inhibition of calphostin C- and tumor necrosis factor - induced apoptosis in U937 histiocytic lymphoma cells.** *J Biol Chem* 1998, **273**:24115-24121.
- Parganas E, Wang D, Stravopodis D, Topham DJ, Jean-Christopher M, Teglund S, Nanin EF, Bodner S, Colamonici OR, Deursen JM, Grosveld G, Ihle JN: **Jak2 is essential for signalling through a variety of cytokine receptors.** *Cell* 1998, **93**:385-395.
- Kleinert H, Euchenhofer C, Fritz G, Ihrig-Biedert I, Forstermann U: **Involvement of protein kinases in the induction of NO synthase II in human DLD-1 cells.** *Brit J Pharmacol* 1997, **123**:1716-1722.
- Marrero MB, Venema VJ, He H: **Inhibition by the Jak/Stat pathway of IFN $\gamma$ - and LPS-stimulated nitric oxide synthase induc-**

- tion in vascular smooth muscle cells. *Biochem Biophys Res Comm* 1998, **252**:508-512.
33. Berghe WV, Plaisance S, Boone E, Bosscher KD, Schmitz ML, Fiers W, Haegeman G: **p38 and extracellular signal-regulated kinase mitogen activate protein kinase pathways are required for nuclear factor-B p65 transactivation mediated by tumor necrosis factor.** *J Biol Chem* 1998, **273**:3285-3290.
  34. Dorn GWII, Tepe NM, Wu G, Yatani A, Liggett SB: **Mechanisms of impaired beta-adrenergic receptor signaling in Gq-mediated cardiac hypertrophy and ventricular dysfunction.** *Mol Pharmacol* 2000, **57**:278-287.
  35. Caramori G, Adcock IM, Ito K: **Anti-inflammatory inhibitors of IkkappaB kinase in asthma and COPD.** *Curr Opin Investig Drugs* 2004, **5**:1141-7.
  36. Bae IK, Min HY, Han AR, Seo EK, Lee SK: **Suppression of lipopolysaccharide-induced expression of inducible nitric oxide synthase by brazilin in RAW 264.7 macrophage cells.** *Eur J Pharmacol* 2005, **513**:237-42.
  37. Shames BD, Selzman CH, Puliido EJ, Meng X, Meldrum DR, McIntyre RC, Harken AH, Banerjee A: **LPS-induced NF-kappaB activation and TNF- release in human monocytes are protein tyrosine kinase dependent and protein kinase C independent.** *J Surg Res* 1999, **83**:69-74.
  38. Wang D, Westerheide SD, Hanson JL, Baldwin AS Jr: **Tumor necrosis factor alpha-induced phosphorylation of RelA/p65 on Ser529 is controlled by casein kinase II.** *J Biol Chem* 2000, **275**:32592-32597.
  39. Zhong H, SuYang H, Erdjument-Bromage H, Tempst P, Ghosh S: **The transcriptional activity of NF-kappaB is regulated by the IkkappaB-associated PKAc subunit through a cyclic AMP-independent mechanism.** *Cell* 1997, **89**:413-24.
  40. Edmunds JW, Mahadevan LC: **MAP kinases as structural adaptors and enzymatic activators in transcription complexes.** *J Cell Sci* 2004, **117**:3715-23.
  41. Delgado M, Ganea D: **Inhibition of IFN-gamma-induced janus kinase-1-STAT1 activation in macrophages by vasoactive intestinal peptide and pituitary adenylate cyclase-activating polypeptide.** *J Immunol* 2000, **165**:3051-7.
  42. Digicaylioglu M, Lipton S: **Erthropoietin-mediated neuroprotection involves cross-talk between Jak2 and NF-kB signalling cascades.** *Nature* 2001, **412**:641-647.
  43. Okugawa S, Ota Y, Kitazawa T, Nakayama K, Yanagimoto S, Tsukada K, Kawada M, Kimura S: **Janus kinase 2 is involved in lipopolysaccharide-induced activation of macrophages.** *Am J Physiol Cell Physiol* 2003, **285**:C399-408.
  44. Alblas J, Honing H, de Lavalette CR, Brown MH, Dijkstra CD, van den Berg TK: **Signal regulatory protein alpha ligation induces macrophage nitric oxide production through JAK/STAT- and hosphatidylinositol 3-kinase/Rac1/NAPDH oxidase/H2O2-dependent pathways.** *Mol Cell Biol* 2005, **25**:7181-92.
  45. Cruz MT, Duarte CB, Goncalo M, Carvalho AP, Lopes MC: **LPS induction of IkkappaB-alpha degradation and iNOS expression in a skin dendritic cell line is prevented by the Janus kinase 2 inhibitor, Tyrophostin B42.** *Nitric Oxide* 2001, **5**:53-61.
  46. Paul A, Cuenda A, Bryant CE, Murray J, Chilvers ER, Cohen P, Gould GW, Plevin R: **Involvement of mitogen-activated protein kinase homologues in the regulation of lipopolysaccharide-mediated induction of cyclo-oxygenase-2 but not nitric oxide synthase in RAW 264.7 macrophages.** *Cell Signal* 1999, **11**:491-7.
  47. Ajizian SJ, English BK, Meals EA: **Specific inhibitors of p38 and extracellular signal-regulated kinase mitogen-activated protein kinase pathways block inducible nitric oxide synthase and tumor necrosis factor accumulation in murine macrophages stimulated with lipopolysaccharide and interferon-gamma.** *J Infect Dis* 1999, **179**:939-44.
  48. Poteser M, Wakabayashi I: **Serum albumin induces iNOS expression and NO production in RAW 267.4 macrophages.** *Br J Pharmacol* 2004, **143**:143-51.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

