



University of Pennsylvania  
ScholarlyCommons

Management Papers

Wharton Faculty Research

2015

# Cluster-Robust Variance Estimation for Dyadic Data

Peter M. Aronow  
*Yale University*

Cyrus Samii  
*New York University*

Valentina A. Assenova  
*University of Pennsylvania*

Follow this and additional works at: [https://repository.upenn.edu/mgmt\\_papers](https://repository.upenn.edu/mgmt_papers)

 Part of the [Management Sciences and Quantitative Methods Commons](#)

## Recommended Citation

Aronow, P. M., Samii, C., & Assenova, V. A. (2015). Cluster-Robust Variance Estimation for Dyadic Data. *Political Analysis*, 23 (4), 564-577. <http://dx.doi.org/10.1093/pan/mpv018>

At the time of publication, author Valentina A. Assenova was affiliated with Yale University. Currently, she is a faculty member at the University of Pennsylvania.

This paper is posted at ScholarlyCommons. [https://repository.upenn.edu/mgmt\\_papers/170](https://repository.upenn.edu/mgmt_papers/170)  
For more information, please contact [repository@pobox.upenn.edu](mailto:repository@pobox.upenn.edu).

---

# Cluster-Robust Variance Estimation for Dyadic Data

## **Abstract**

Dyadic data are common in the social sciences, although inference for such settings involves accounting for a complex clustering structure. Many analyses in the social sciences fail to account for the fact that multiple dyads share a member, and that errors are thus likely correlated across these dyads. We propose a non-parametric, sandwich-type robust variance estimator for linear regression to account for such clustering in dyadic data. We enumerate conditions for estimator consistency. We also extend our results to repeated and weighted observations, including directed dyads and longitudinal data, and provide an implementation for generalized linear models such as logistic regression. We examine empirical performance with simulations and an application to interstate disputes.

## **Keywords**

cluster robust variance estimation, dyadic data, agnostic regression

## **Disciplines**

Management Sciences and Quantitative Methods

## **Comments**

At the time of publication, author Valentina A. Assenova was affiliated with Yale University. Currently, she is a faculty member at the University of Pennsylvania.

# Cluster-Robust Variance Estimation for Dyadic Data

Peter M. Aronow, Cyrus Samii and Valentina A. Assenova\*

July 24, 2015

---

\*Peter M. Aronow is Assistant Professor, Department of Political Science, Yale University, 77 Prospect St., New Haven, CT 06520 (Email: peter.aronow@yale.edu). Cyrus Samii (contact author) is Assistant Professor, Department of Politics, New York University, 19 West 4th St., New York, NY 10012 (Email: cds2083@nyu.edu). Valentina A. Assenova is Doctoral Student, School of Management (Organizations and Management), Yale University, 165 Whitney Avenue, New Haven, CT 06520 (Email: valentina.assenova@yale.edu). The authors thank Neal Beck, Allison Carnegie, Dean Eckles, Donald Lee, Winston Lin, Kelly Rader, Olav Sorenson, the *Political Analysis* editors and two reviewers for helpful comments. We thank Jonathan Baron and Lauren Pinson for research assistance. Replication materials are available on the *Political Analysis* Dataverse (<https://dataverse.harvard.edu/dataverse/pan>).

# Cluster-Robust Variance Estimation for Dyadic Data

## Abstract

Dyadic data are common in the social sciences, although inference for such settings involves accounting for a complex clustering structure. Many analyses in the social sciences fail to account for the fact that multiple dyads share a member, and that errors are thus likely correlated across these dyads. We propose a nonparametric, sandwich-type robust variance estimator for linear regression to account for such clustering in dyadic data. We enumerate conditions for estimator consistency. We also extend our results to repeated and weighted observations, including directed dyads and longitudinal data, and provide an implementation for generalized linear models such as logistic regression. We examine empirical performance with simulations and an application to interstate disputes.

Keywords: cluster robust variance estimation, dyadic data, agnostic regression

Word count: 5,022 (including main text, appendix for print, captions, and references).

# 1 Introduction

Dyadic data are central in social science applications ranging from international relations to “speed dating.”<sup>1</sup> A challenge with dyadic data is to account for the complex dependency structure that arises due to the connections between dyad members. For instance, in a study of international conflict, a change in leadership in one country may affect relations with all countries with which that country is paired in the data, thereby inducing a correlation between these dyadic observations. This generates a cluster of dependent observations associated with that country. As leadership changes occur in multiple countries, the correlations emanating from each of these countries will overlap into a web of interwoven clusters. We refer to such interwoven dependency in dyadic data as “dyadic clustering.” By ignoring the dyadic clustering, the analysis would take the dyad-level changes emanating from a single leadership change as independently informative events, rather than a single, clustered event. An analysis that only accounts for correlations in repeated observations of dyads (whether by clustering standard errors or using dyad fixed effects) would fail to account for such inter-dyad correlation.

Statistical analysis of dyads typically estimates how dyad-level outcomes (e.g., whether there is open conflict between countries or the decision for one person to ask another on a date) relate to characteristics of the individual units as well as to the dyad as a whole (e.g., measures of proximity between units). The usual approach is to regress the dyad-level outcome on unit- and dyad-level predictors. Due to dyadic clustering, the observations contributing to such an analysis are not independent. Failure to account for dyadic clustering may result in significance tests or confidence intervals with sizes that are far too small relative to the true distribution of the parameters of interest.

We establish sufficient conditions for the consistency of a non-parametric sandwich estimator

---

<sup>1</sup>For example, in the past five years the *American Political Science Review*, *American Journal of Political Science*, and *International Organization* have together published 62 papers that include dyadic analyses.

(Huber, 1967; White, 1980a) for the variance of regression coefficients under dyadic clustering. Cluster-robust sandwich estimators are common for addressing dependent data (Angrist and Pischke, 2009, Ch. 8; Liang and Zeger, 1986). Cameron, Gelbach and Miller (2011) provide a sandwich estimator for “multi-way” clustering, accounting, for example, for clustering between people by geographic location and age category. We extend these methods to dyadic clustering, accounting for the fact that dyadic clustering does not decompose neatly into a few crosscutting and disjoint groups of units; rather, each *unit* is the basis of its own cluster that intersects with other units’ clusters. Fafchamps and Gubert (2007, Eq. 2.5) propose a sandwich estimator for dyadic clustering that is very similar to what we propose below. Their derivation is constructed through analogy to the results of Conley (1999). However, neither paper establishes conditions for consistency under dyadic clustering. We establish such consistency conditions. We also provide extensions to the longitudinal case and generalized linear models such as logistic regression. We evaluate performance with simulations and a reanalysis of a classic study on interstate disputes (Russett and Oneal, 2001). The appendix generalizes to weighted data and generalized linear models, and the supporting information provides another illustration with a speed dating experiment (Fisman et al., 2006).

Current statistical approaches to handling dyadic clustering include the use of parametric restrictions in mixed-effects models (Gelman and Hill, 2007; Hoff, 2005; Kenny, Kashy and Cook, 2006), the spatial error-lag model (Beck, Gleditsch and Beardsley, 2006), or permutation inference for testing against sharp null hypotheses (Erikson, Pinto and Rader, 2014). These approaches have important limitations that our approach overcomes. First, mixed effects models and spatial lag models impose a parametric structure to address the clustering problem. This makes them sensitive to misspecification of the conditional mean (that is, deviations between the data and the linearity assumption). Our approach is robust in that it provides asymptotically valid standard errors even under such misspecification. This is valuable in itself to the extent that models are typically approximations (Buja et al., 2014; Hubbard et al., 2014), although readers should not take

this to mean that they are free from the obligation to fit as good an approximation as possible. It is also valuable in providing a reliable benchmark to use in evaluating model specification (King and Roberts, 2014; White, 1980a, 1981). Second, Erikson, Pinto and Rader (2014)’s solution of non-parametric randomization inference does not provide a procedure for obtaining valid confidence intervals, while our procedure does. Third, all three of these alternatives require considerable computational power, possibly exceeding available computational resources even for data analyses that are common in international relations or network analysis. We demonstrate below that a mixed-effects or spatial error-lag approach is infeasible in a typical international relations example. In contrast, our proposed estimator is easy to compute. The variance estimation methods we develop are a natural complement to non- and semi-parametric approaches to regression with dyadic data (Green, Kim and Yoon, 2001).

## 2 Setting

We work within the “agnostic” (Angrist and Imbens, 2002; Lin, 2013) or “assumption lean” (Buja et al., 2014) regression framework developed by Angrist and Pischke (2009), Goldberger (1991), and White (1980b, 1982).<sup>2</sup> We begin with a cross-section of undirected dyads and derive the basic convergence results for this case. Below we extend these results to repeated dyads, which covers directed dyads and longitudinal data. Proofs are in the appendix.

Begin with a large population from which we take a random (i.i.d.) sample of units, with the sample members indexed by  $i = 1, \dots, N$  and grouped into  $D = \binom{N}{2} = N(N-1)/2$  dyads. Pairs of unit indices within each dyad map to dyadic indices,  $1, \dots, D$ , through the index function  $d(i, j) =$

---

<sup>2</sup>Our results would hold under a regression model approach along the lines of Greene (2008) or Davidson and MacKinnon (2004). The value of using the “agnostic regression” approach is that the derivation is robust to violations of the assumption that the regression specification is correct for the conditional expectation. Nevertheless, this should not be taken as an endorsement for ignoring misspecification or assuming that linear approximations with robust standard errors can cure misspecification problems (King and Roberts, 2014). This paper leaves aside issues of causal identification and focuses only on estimation and statistical inference. Nonetheless, the issues discussed here would be relevant for studies that attempt to estimate causal effects with dyadic data.

$d(j, i)$ , with an inverse correspondence  $s(d(i, j)) = \{i, j\}$ , and we assume no  $d(i, i)$  type dyads. Consider a linear regression of  $Y_{d(i, j)}$  on a  $k$ -length column vector of regressors  $X_{d(i, j)}$  (which includes the constant):

$$Y_{d(i, j)} = X'_{d(i, j)}\beta + \varepsilon_{d(i, j)},$$

where  $\beta$  is the slope that we would obtain if we could fit this model to the entire population, allowing for possible non-linearity in the true relationship between  $E[Y_{d(i, j)}|X_{d(i, j)}]$  and  $X_{d(i, j)}$ , and  $\varepsilon_{d(i, j)}$  is the corresponding population residual.<sup>3</sup> To lighten notation, for the remainder of the discussion we use  $d = d(i, j)$ .

Define the sample data as  $\mathbf{X} = (X_1 \dots X_D)'$  and  $Y = (Y_1 \dots Y_D)'$ . The ordinary least squares (OLS) estimator is

$$\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'Y,$$

with residual  $e_d = Y_d - X'_d\hat{\beta}$ . Since the values of  $Y_d$  and  $X_d$  are determined by the characteristics of the units  $i$  and  $j$ ,  $\varepsilon_d$  and  $\varepsilon_{d'}$  for dyads containing either unit  $i$  or  $j$  are allowed to be correlated by construction. However, by random sampling of units,  $\text{Cov}(\varepsilon_d, \varepsilon_{d'}) = 0$  for all dyads that do not share a member. The number of pairwise dependencies between  $i$  and  $j$  for which  $\text{Cov}(\varepsilon_d, \varepsilon_{d'}) \neq 0$  is  $O(N^3)$ . Let the support for  $X_d$  and  $Y_d$  be bounded in a finite interval and assume the usual rank conditions on  $\mathbf{X}$  (Chamberlain, 1982; White, 1984).

**Lemma 1.** *Suppose data and  $\hat{\beta}$  as defined above. As  $N \rightarrow \infty$ , the asymptotic distribution of  $\sqrt{N}(\hat{\beta} - \beta)$  has mean zero and variance*

$$V = \frac{N}{D^2} E[X_d X'_d]^{-1} \text{Var} \left[ \sum_{d=1}^D X_d \varepsilon_d \right] E[X_d X'_d]^{-1}, \quad (1)$$

---

<sup>3</sup>If  $E[Y_{d(i, j)}|X_{d(i, j)}]$  is nonlinear in  $X_{d(i, j)}$ , the population residual will itself vary in  $X_{d(i, j)}$ , which undermines inference that assumes that  $\varepsilon_{d(i, j)}$  is independent of  $X_{d(i, j)}$ . The agnostic regression approach avoids such an assumption.



with,

$$\text{Var} \left[ \sum_{d=1}^D X_d \varepsilon_d \right] = \sum_{d=1}^D \left( \underbrace{\text{E} \{ X_d X_d' \text{Var} [\varepsilon_d | \mathbf{X}] \}}_A + \underbrace{\sum_{d' \in \mathcal{S}(d)} \text{E} \{ X_d X_{d'}' \text{Cov} [\varepsilon_d, \varepsilon_{d'} | \mathbf{X}] \}}_B \right), \quad (2)$$

where  $\mathcal{S}(d) = \{d' \neq d\} \cap \{d' : s(d) \cap s(d') \neq \emptyset\}$ , the set of dyads other than  $d$  that share a member from  $d$ .

$A$  is the dyad-specific contribution to the variance, and  $B$  is the contribution due to inclusion of common units in multiple dyads. Note that these features of the distribution of  $\sqrt{N}(\hat{\beta} - \beta)$  establish the consistency of  $\hat{\beta}$  as well, which is not surprising given standard results for the consistency of OLS coefficients on dependent data (White, 1984).

### 3 Identification and Estimation

Given data as defined above, we examine the properties of a plug-in variance estimator that is analogous to the sandwich estimators defined for heteroskedastic or clustered data (Arellano, 1987; Cameron, Gelbach and Miller, 2011; Liang and Zeger, 1986; White, 1980a). We establish sufficient conditions for consistency of the plug-in variance estimator. We consider the cross-sectional case and repeated observations case. The appendix also contains extensions to weighted observations and generalized linear models.

**Proposition 1.** *Define the variance estimator*

$$\hat{V} = (\mathbf{X}'\mathbf{X})^{-1} \sum_{d=1}^D \left( X_d X_d' e_d^2 + \sum_{d' \in \mathcal{S}(d)} X_d X_{d'}' e_d e_{d'} \right) (\mathbf{X}'\mathbf{X})^{-1}. \quad (3)$$

*Under the conditions of Lemma 1 and if  $X_d$  and  $e_d$  have bounded support with non-zero second*

moments and cross-moments, then as  $N \rightarrow \infty$ ,

$$N\hat{V} - V \xrightarrow{p} 0.$$

The proposition indicates that  $\hat{V}$  provides a consistent estimator to characterize the true asymptotic sampling distribution of the regression coefficients,  $\hat{\beta}$ . Standard error estimates for  $\hat{\beta}$  are obtained from the square roots of the diagonals of  $\hat{V}$ . The assumption of bounded support for  $X_d$  and  $e_d$  merely rules out situations that are unlikely to arise in real world data anyway (e.g., where the mass of the data pushes out toward infinity as  $N$  grows).

Repeated dyad observations are common in applied settings. For example, the data may include multiple observations for dyads over time. Dyadic panels are very common in studies of international relations. Or, if the dyadic information is directional, then the data will contain two observations for each dyad, with one observation capturing outcomes that go in the  $i$  to  $j$  direction, and the other capturing outcomes that go in the  $j$  to  $i$  direction. This is conceptually distinct from repeated observations over time. But if there are dyadic dependences for both senders and receivers of a directed dyad, then the dependence structure for a pair of directed  $i$ - $j$  dyads will be the same as if we had repeated observations of an undirected  $i$ - $j$  dyad. The results above translate straightforwardly to the repeated dyads setting. Formally, suppose that for each dyad  $d$  observations are indexed by  $t = 1, \dots, T(d)$ , where the  $T(d)$  values are fixed and finite. Let  $(Y_{dt}, X'_{dt})$  represent the data for observation  $t$  from dyad  $d$ ,

$$Y_d = \begin{pmatrix} Y_{d1} \\ \vdots \\ Y_{dT(d)} \end{pmatrix} \quad \text{and} \quad X_d = \begin{pmatrix} X'_{d1} \\ \vdots \\ X'_{dT(d)} \end{pmatrix},$$

and  $Y = (Y'_1 \dots Y'_d \dots Y'_D)'$  and  $\mathbf{X} = (X'_1 \dots X'_d \dots X'_D)'$ , that is, the stacked  $Y_d$  vectors and  $X_d$  matrices.

Let  $\beta_r$  and  $\hat{\beta}_r$  denote, respectively, the population slope and OLS estimator as defined above but now applied to the repeated dyads data.

**Corollary 1.** *For the repeated dyads case, assume the same conditions as in Proposition 1 and consider the following variance estimator*

$$\hat{V}_r = (\mathbf{X}'\mathbf{X})^{-1} \sum_{d=1}^D \left( X_d' e_d e_d' X_d + \sum_{d' \in \mathcal{S}(d)} X_d' e_d e_{d'}' X_{d'} \right) (\mathbf{X}'\mathbf{X})^{-1}.$$

Then as  $N \rightarrow \infty$ ,

$$N\text{Var}[\hat{\beta}_r - \beta_r] - V_r \xrightarrow{p} 0,$$

and

$$N\hat{V}_r - V_r \xrightarrow{p} 0,$$

where

$$V_r = \frac{N}{\left[ \sum_{d=1}^D T(d) \right]^2} \mathbb{E} [X_d X_d']^{-1} \mathbb{E} \left( \mathbb{E} [X_d' \text{Cov} [\varepsilon_d | \mathbf{X}] X_d] + \sum_{d' \in \mathcal{S}(d)} \mathbb{E} [X_d' \text{Cov} [\varepsilon_d, \varepsilon_{d'} | \mathbf{X}] X_{d'}] \right) \mathbb{E} [X_d X_d']^{-1}.$$

Some remarks are in order when it comes to the repeated observations case. First, our analysis of the repeated observations case takes the  $T(d)$  values to be fixed and finite and demonstrates consistency as  $N$  grows. In a cross-national study, this would mean that one should pay most attention to the number of countries, rather than the amount of time, that are in the data. Under fixed  $N$  and growing  $T(d)$  further assumptions would be required for consistency, such as serial correlation of fixed order or, more generally, strong mixing over time. Second, with repeated observations, a common strategy for identifying effects is to use dyad-specific fixed effects (Green, Kim and Yoon, 2001). With fixed and finite  $T(d)$ , this does not introduce any new complications. As in Arellano (1987) for the case of serial correlation with fixed effects, the results for dyadic clustering translate directly to centered data, and dyadic fixed-effects regression amounts to centering the data on

dyad-specific means. With fixed  $N$  and growing  $T(d)$ , however, additional assumptions are needed for consistency (Hansen, 2007; Stock and Watson, 2008).

For efficient computation, we follow Cameron, Gelbach and Miller (2011) to perform a “multi-way decomposition” of the dyadic clustering structure:

**Proposition 2.** *We have the following algebraic equivalence,*

$$\hat{V}_r = \sum_{i=1}^N \hat{V}_{C,i} - \hat{V}_D - (N-2)\hat{V}_0,$$

where

$$\hat{V}_{C,i} = (\mathbf{X}'\mathbf{X})^{-1}\hat{\Sigma}_{C,i}(\mathbf{X}'\mathbf{X})^{-1}$$

$$\hat{V}_D = (\mathbf{X}'\mathbf{X})^{-1}\hat{\Sigma}_D(\mathbf{X}'\mathbf{X})^{-1}$$

$$\hat{V}_0 = (\mathbf{X}'\mathbf{X})^{-1}\hat{\Sigma}_0(\mathbf{X}'\mathbf{X})^{-1},$$

and

$$\hat{\Sigma}_{C,i} = \sum_{j \neq i} \sum_{k \neq i} X'_{d(i,j)} e_{d(i,j)} e'_{d(i,k)} X_{d(i,k)} + \sum_{j,k \neq i} \sum_{t=1}^{T(d(j,k))} X_{d(j,k)t} X'_{d(j,k)t} e_{d(j,k)t}^2,$$

$$\hat{\Sigma}_D = \sum_{d=1}^D X'_d e_d e'_d X_d, \quad \text{and} \quad \hat{\Sigma}_0 = \sum_{d=1}^D \sum_{t=1}^{T(d)} X_{dt} X'_{dt} e_{dt}^2.$$

$\hat{V}_{C,i}$  is the usual asymptotically consistent cluster-robust variance estimator (with no degrees-of-freedom adjustment) that clusters all dyads containing unit  $i$  and assumes all other observations to be independent.  $\hat{V}_D$  is the same cluster robust estimator but clustering all repeated dyad observations.  $\hat{V}_0$  is the usual asymptotically consistent heteroskedasticity robust (HC) variance estimator that assumes all observations (even within repeated dyad groupings) are independent. To understand this decomposition, note that dyadic clustering involves clustering on each of  $N$  units. But in summing the contributions from unit-specific clusters (the  $\hat{V}_{C,i}$ s), we double count the dyad

contributions ( $\hat{V}_D$ ) and add in the independent contributions ( $V_0$ )  $N$  times. We can correct for this by subtracting  $\hat{V}_D$ , which also removes the  $V_0$  component, and then subtracting  $N - 2$  of the  $V_0$  components. (The cross section cases simply sets  $\hat{V}_D = \hat{V}_0$ ). This decomposition shows that one can compute the dyadic cluster robust estimator using readily available robust standard error commands.<sup>4</sup>

Our dyadic cluster robust variance estimator allows one to perform valid inference under less restrictive dependence assumptions that are used to identify random effects or spatial error lag models. Our multi-way decomposition also shows that it is computationally much simpler than those alternatives. The latter point is relevant when one has many units or time periods in which case random effects and spatial error lag models, despite their restrictions, may be infeasible with current computational resources given the challenges of evaluating a likelihood with as many dependencies as emerge in a dyadic analysis.

## 4 Simulation Evidence

We use Monte Carlo simulations to evaluate the finite sample properties of the proposed estimators under the cross-sectional and repeated dyads settings. We suppose that population values obey the following,

$$Y_{d(i,j)t} = \beta_0 + \beta_1 |X_i - X_j| + \underbrace{\alpha_i + \alpha_j + v_{d(i,j)t}}_{\varepsilon_{d(i,j)t}}, \quad (4)$$

where  $Y_{d(i,j)t}$  is the  $t^{th}$  observed outcome for the dyad that includes units  $i$  and  $j$ ,  $X_i$ ,  $X_j$ ,  $\alpha_i$ ,  $\alpha_j$ , and  $v_{d(i,j)t}$  are independent draws from standard Normal distributions, and the compound error,  $\varepsilon_{d(i,j)t} = \alpha_i + \alpha_j + v_{d(i,j)t}$ , is unobserved. In the cross-sectional case, we only observe one outcome per dyad, so  $t = 1$  for all observations (that is, the  $t$  subscript is extraneous for the cross-sectional

---

<sup>4</sup>Code for implementation in R is available from the replication archive posted to the Dataverse page references in the acknowledgments note).

case). In the repeated observations case, we have two observations per dyad, so  $t = 1, 2$  for all dyads. We fix  $\beta_0 = 0$  and  $\beta_1 = 1$ . We use OLS to estimate  $\hat{\beta}_0$  and  $\hat{\beta}_1$ . In the supplemental information we show results for mixed effects models as well. The fact that  $X_i$  and  $\alpha_i$  are constant across dyads that include unit  $i$  (same for  $j$ ) implies non-zero intra-class correlation in both  $X$  and  $\varepsilon$  among sets of dependent dyads, in which case ignoring the dependence structure will tend to understate the variability in  $\hat{\beta}_0$  and  $\hat{\beta}_1$ . This is the dyadic version of Moulton (1986)’s problem.

Results from 500 simulation runs are shown in Figure 1. The  $x$ -axis shows the sample size, where we show results for samples with 20, 50, 100, and 150 units (implying 190, 1,225, 4,950, and 11,175 dyads, respectively). The  $y$ -axis is on the scale of the standard error of the coefficients. The black diamonds plot the empirical standard standard errors for each of the sample sizes (that is, the standard deviation of the simulation replicates of the coefficient estimates,  $\hat{\beta}_0$  and  $\hat{\beta}_1$ ). The black box plots show the distribution of our proposed standard error estimator. The gray box plots show, in the top figures, the distribution of the “HC2” heteroskedasticity robust variance estimator, which does not account for either dyadic or repeat observation clustering (White, 1980a; MacKinnon and White, 1985), and for the bottom figures, the “cluster robust” analog of White (1980a)’s estimator accounting for dependence in repeated dyad observations (Arellano, 1987; Liang and Zeger, 1986).

Convergence of our proposed standard errors based on  $\hat{V}$  and  $\hat{V}_r$  (black box plots) to the true standard error (black diamonds) is quick. The alternative estimators, which do not account for dyadic clustering, grossly understate the variability in the coefficient estimates. These results suggest that in finite samples from data generating processes that resemble our simulations, the proposed estimator is quite reliable for inference on OLS coefficients so long as the sample size is on the order of 50 to 100 units. Most applications in political science operate with samples of at least this size. Changes to the shape of the error distributions (e.g., uniform or bimodal) yield the same results.<sup>5</sup>

---

<sup>5</sup>The simulation code, which allows one to choose arbitrary data distributions, is available from the replication archive posted to the Dataverse page referenced in the acknowledgement note.




Figure 1: Convergence of the standard error estimator as  $N$  goes from 20 to 150 (implying number of dyads goes from 190 to 11,175) for a cross section of dyads (top) and repeated dyad observations ( $T=2$  for all dyads). Black box plots show the distribution of standard error estimates using the proposed dyadic cluster robust estimator. In the top two figures, gray box plots show standard error estimates from a heteroskedasticity robust estimator while in the bottom two figures, gray box plots show standard error estimates from a “naive” cluster-robust estimator that clusters by dyad over repeated observations (but does not cluster by units across dyads).

In the supporting information, we demonstrate comparisons between mixed effects models and our “robust” estimator.<sup>6</sup> Specifically, we compare OLS with dyadic cluster robust standard errors to a two-way random effects model that incorporates normal random effects for each of the  $N$  units (Gelman and Hill, 2007). The data generating process captured above (that is, expression 4 with normal data) corresponds precisely to the assumptions of a two-way normal random effects model, and so it is no surprise that we find this model to be more efficient than OLS and to produce consistent standard errors. However, as discussed above, such models are sensitive to misspecification. For example, in fitting a linear approximation model to mildly non-linear data, the dyadic cluster robust standard errors continue to be consistent for the coefficients of the linear approximation. The degree of misspecification in this case is not extreme (as illustrated in the appendix) and resembles the kind of approximations that is common in applied work (Buja et al., 2014). Nonetheless, the random effects standard errors are inconsistent and substantially anti-conservative. Such sensitivity to misspecification is one problem for the random effects model. Another problem arises when there is unobserved country-level heterogeneity that could confound estimates of the coefficients of interest. In such situations, our dyadic cluster robust estimator would be a natural complement to a fixed effects analysis, analogous to what Arellano (1987) proposes for non-dyadic repeated observation settings. The third problem arises in computational feasibility, an issue that arises in our application to which we now turn.

## **5 International Militarized Disputes Application**

Our application is based on the classic study by Russett and Oneal (2001) on the determinants of international militarized conflicts. They study how the likelihood of a militarized dispute between

---

<sup>6</sup> We sought also to include a comparison to spatial error lag models, but the models failed to converge with  $N \geq 100$ . We note that in the original Beck, Gleditsch and Beardsley (2006) paper, they limited their analysis of dyadic data to small cross sections. Our experience suggests that it may be infeasible to fit such models with substantially larger datasets.



states in a dyad relates to various dyad-level attributes, including an indicator of whether the two states are formal allies, the log of the ratio of an index of military capabilities, the lowest score in the dyad on a democracy index, the ratio of the value of trade and the larger GDP of the two states, an indicator of whether both states are members of a common international organization, an indicator of whether the states are geographically noncontiguous, the log of the geographic (Euclidean) distance between the two states' capitals, and an indicator of whether both states are "minor powers" in the international system. Dyadic clustering could arise in many ways with these data, for example if a country entered into an alliance, thereby changing the joint alliance indicators, or if the military capabilities of a country changed, thereby changing the power ratios.

We replicate Russett and Oneal's primary analysis as reported in their Table A5.1. They use annual data on 146 states in the international system paired into 1,178 dyads (out of 10,585 possible) and observed for as few as one and as many as ninety years between 1885 and 1991, for a total of 39,988 observations. In their original analysis, Russett and Oneal fit a GEE model assuming AR(1) errors within dyads over time (Zorn, 2001). Table 1 shows the results of our reanalysis. Columns (1) and (2) replicate the original published results. Columns (3)-(6) show coefficients and various standard error estimates for a simple (pooled) logistic regression. There is little difference in the coefficient estimates from the GEE-AR(1) model as compared to the simple logistic regression, so we focus on the standard error estimates. Column (4) contains estimates that account only for dyad-year heteroskedasticity, column (5) also accounts for arbitrary dependence over time for each dyad, and then column (6) also accounts for dyadic clustering. Accounting for the dyadic and repeated-observation clustering results in standard error estimates that are sometimes an order of magnitude larger than what we obtain when we ignore all clustering and also considerably larger than what one would estimate were one to account for repeated dyads clustering but ignore the inter-dyadic clustering. The latter result is also relevant when comparing the standard errors from the original GEE-AR(1) model, as they resemble the estimates in column (5). In their original analysis, Russett and Oneal found that all of the predictors had a statistically significant relation-

Table 1: Estimates and robust standard error estimates from logistic regression of an indicator for militarized conflict between dyads on dyad-level predictors, 1885-1991, based on Russett and Oneal (2001, Table A5.1).

Predictor of Militarized Conflict	(1)	(2)	(3)	(4)	(5)	(6)
	Original GEE estimates		Simple logistic regression estimates			
	Coef.	S.E.	Coef.	Het. Robust S.E.	Naive Cluster Robust S.E.	Dyadic Cluster Robust S.E.
Alliances	-0.539	0.159	-0.595	0.069	0.175	0.265
Power ratio	-0.318	0.043	-0.328	0.020	0.047	0.070
Lower democracy	-0.061	0.009	-0.064	0.005	0.010	0.015
Lower dependence	-52.924	13.405	-67.668	10.734	17.560	24.749
International organizations	-0.013	0.004	-0.011	0.002	0.005	0.008
Noncontiguity	-0.989	0.168	-1.048	0.074	0.181	0.185
Log distance	-0.376	0.065	-0.375	0.026	0.068	0.102
Only minor powers	-0.647	0.178	-0.618	0.078	0.188	0.344
Constant	-0.128	0.536	-0.181	0.211	0.562	0.840

*N* = 39,988 dyad-year observations for 1,178 dyads and 146 units.

ship to the likelihood of militarized conflict. But when one takes into account dyadic clustering, the coefficient for international organizations would fail to pass a conventional significance test (e.g.,  $t > 1.64$ ).

We attempted also to study how inferences might change with a random effects approach and a spatial error lag approach. The appropriate random effects model would be a three-way random effects model to account for each state in a dyad as well as the dyad as a whole over time. We attempted to fit three-way random effects logit models using the `lmer` package in R and the `melogit` function in Stata. We posted the jobs to a university cluster with 24 Intel Xeon cores and 48 gigabytes of RAM per node. In neither case did the random effects models converge. In fact, in neither case could we obtain estimates of any form after days on the cluster. For the spatial error lag model, we sought to implement both the spatial error lag model in the `spdep` package in R and the two-step estimator of Neumayer and Pluempfer (2010) in Stata. The latter take as an input dyadic dependence information that can be produced using the `spundir` package in Stata. The spatial error lag model in the `spdep` failed to produce estimates and `spundir` package failed

to resolve within reasonable time (a week) while running on the same cluster.<sup>7</sup> This brings to the foreground the issue of computational complexity, by which our cluster robust estimator is much less demanding.

## 6 Conclusion

We have established convergence properties for a non-parametric variance estimator for regression coefficients that accounts for dyadic clustering. The estimator applies no restrictions on the dependency structure beyond the dyadic clustering assumption. The estimator is robust to the regression model being misspecified for the conditional mean. Such robustness is important because regression analysis typically relies on linear (in coefficients) approximations of unknown conditional mean functions (Angrist and Pischke, 2009; Buja et al., 2014; Hubbard et al., 2014). Of course our analysis in no way excuses analysts from their responsibility to obtain as good an approximation as possible nor does a robust fix for standard errors also solve the problem of finding a good approximation. Given a reasonable approximation for the conditional mean, the methods we have developed here allow for more accurate asymptotic statistical significance tests and confidence intervals.

The estimator is consistent in the number of units that form the basis of dyadic pairs. Simulations show that the estimator approaches the true standard error with modestly-sized samples from a reasonable data generating process. Applications show that inferences can be seriously anti-conservative if one fails to account for the dyadic clustering. This estimator is a natural complement to the non-parametric and semi-parametric regression analyses that are increasingly common in the social sciences (Angrist and Pischke, 2009). Given that we can express the estimator

---

<sup>7</sup>See also footnote 6. The computational complexity is due to the fact that the model attempts to account for dependence across all countries in the dataset (rather than small subsets, as would be the case in a typical spatial analysis) and the fact that in this application the set of cross-section units changes over time and therefore does not allow for a stable adjacency matrix over time.

as the sum of simpler and easy-to-compute robust variance estimators, it could be applied to any estimator for which a cluster-robust option is available.

A cost of the robust approach is efficiency. Our simulations show that a two-way (for cross-sectional data) random effects model can be considerably more efficient and provide reliable inference when the conditional mean is correctly specified. Our robust estimator could provide the basis of a test for misspecification (King and Roberts, 2014; White, 1980*a*, 1981). If there is little evidence of misspecification, the random effects estimator would be a reasonable choice given its efficiency. However, problems of computational non-convergence may make the random effects estimator infeasible—we encountered this in our application when we tried to use a three-way random effects model for dyadic panel data.

Of course, accounting for dynamic dyadic clustering may fail to fully account for all relevant dependencies in the data. For example, units may exhibit higher-order network effects: for example a shock to unit *A* may affect unit *C* through connections that run via a third unit, *B*. In such cases, the methods developed here will likely yield standard errors that are too small, although they should still outperform methods that fail even to account for dyadic clustering.

## A Proof of Lemma 1

*Proof.* We have

$$\sqrt{N}(\hat{\beta} - \beta) = \left[ \frac{1}{D} \sum_{d=1}^D X_d X_d' \right]^{-1} \frac{\sqrt{N}}{D} \sum_{d=1}^D X_d \varepsilon_d.$$

Take  $[(1/D) \sum_{d=1}^D X_d X_d']^{-1}$  first. By continuity of the inverse, the finite support of  $X_d$  and  $\varepsilon_d$ ,  $\mathbf{X}$  full rank, and the fact that the sum of dependent terms in this sum is  $O(D^{3/2})$  (Lehmann, 1999, Eq. 2.8.4), we have that

$$\left[ \frac{1}{D} \sum_{d=1}^D X_d X_d' \right]^{-1} \xrightarrow{p} E[X_d X_d']^{-1}.$$

By Slutsky's theorem,  $\sqrt{N}(\hat{\beta} - \beta)$  has the same asymptotic distribution as  $E[X_d X_d']^{-1} \frac{\sqrt{N}}{D} \sum_{d=1}^D X_d \varepsilon_d$ , which has mean zero and variance,

$$V = \frac{N}{D^2} E[X_d X_d']^{-1} \text{Var} \left[ \sum_{d=1}^D X_d \varepsilon_d \right] E[X_d X_d']^{-1}. \quad (5)$$

Then,

$$\begin{aligned} \text{Var} \left[ \sum_{d=1}^D X_d \varepsilon_d \right] &= \sum_{d=1}^D \sum_{d'=1}^D \text{Cov} [X_d \varepsilon_d, X_{d'} \varepsilon_{d'}] \\ &= \sum_{d=1}^D \sum_{d'=1}^D E \{ X_d X_{d'}' \text{Cov} [\varepsilon_d, \varepsilon_{d'} | \mathbf{X}] \} \\ &= \sum_{d=1}^D \left( \underbrace{E \{ X_d X_d' \text{Var} [\varepsilon_d | \mathbf{X}] \}}_A + \underbrace{\sum_{d' \in \mathcal{S}(d)} E \{ X_d X_{d'}' \text{Cov} [\varepsilon_d, \varepsilon_{d'} | \mathbf{X}] \}}_B \right) \end{aligned} \quad (6)$$

where  $\mathcal{S}(d) = \{d' \neq d\} \cap \{d' : s(d) \cap s(d') \neq \emptyset\}$ , the set of dyads other than  $d$  that share a member from  $d$ . □

## B Proof of Proposition 1

*Proof.* By Chebychev's inequality, two conditions are sufficient for  $N\hat{V} - V \xrightarrow{P} 0$  as  $N \rightarrow \infty$ : (i)  $E[\hat{V} - V] \rightarrow 0$ , and (ii)  $\text{Var}[N\hat{V} - V] = \text{Var}[N\hat{V}] \rightarrow 0$  (Lehmann, 1999, Thm. 2.1.1).

For (i), consider the interior (“meat”) of  $\hat{V}$ :

$$\hat{\Sigma} = \sum_{d=1}^D \left( X_d X_d' e_d^2 + \sum_{d' \in \mathcal{S}(d)} X_d X_{d'}' e_d e_{d'} \right)$$

and the corresponding term for  $V$ :

$$\Sigma = \sum_{d=1}^D \left( E\{X_d X_d' \text{Var}[\varepsilon_d | \mathbf{X}]\} + \sum_{d' \in \mathcal{S}(d)} E\{X_d X_{d'}' \text{Cov}[\varepsilon_d, \varepsilon_{d'} | \mathbf{X}]\} \right).$$

We have

$$E[\hat{\Sigma}] = \sum_{d=1}^D \left( E\{X_d X_d' E[e_d^2 | \mathbf{X}]\} + \sum_{d' \in \mathcal{S}(d)} E\{X_d X_{d'}' E[e_d e_{d'} | \mathbf{X}]\} \right) \rightarrow \Sigma$$

by consistency of  $\hat{\beta}$ , an implication of Lemma 1. Lemma 1 also established that  $(\mathbf{X}'\mathbf{X}/D)^{-1} \xrightarrow{P} E[X_d X_d']^{-1}$ , while boundedness of  $X_d$  implies uniform boundedness of  $E[X_d X_d']^{-1}$ . Taking these elements together, uniform continuity implies  $E[\hat{V} - V] \rightarrow 0$  (cf. White, 1980a, Thm. 1).

For (ii), we follow the strategy of (Lehmann, 1999, section 2.8), which requires showing that growth of covariance contributions due to dependent units is bounded in  $N$ . Begin by rewriting  $N\hat{V}$  as

$$N\hat{V} = N(\mathbf{X}'\mathbf{X}/D)^{-1} \left( D^{-2} \sum_d \sum_{d'} I_{d,d'} X_d X_{d'}' e_d e_{d'} \right) (\mathbf{X}'\mathbf{X}/D)^{-1}$$

where  $I_{d,d'}$  is an indicator function denoting a pair of dyads  $d$  and  $d'$  that share a member, as defined above. We have established that  $(\mathbf{X}'\mathbf{X}/D)^{-1}$  converges in probability to a finite limit,  $E[X_d X_d']^{-1}$ ,

so we can treat this as  $O(1)$ . Given bounded support,  $\text{Var}[N\hat{V}]$  is of the same order as

$$\begin{aligned}\Omega &= N^2 D^{-4} \text{Var} \left[ \sum_d \sum_{d'} I_{d,d'} X_d X_{d'}' e_d e_{d'} \right] \\ &= N^2 D^{-4} \sum_d \sum_{d'} \sum_{d''} \sum_{d'''} \text{Cov}(I_{d,d'} X_d X_{d'}' e_d e_{d'}, I_{d'',d'''} X_{d''} X_{d'''}' e_{d''} e_{d'''}).\end{aligned}$$

To show  $\text{Var}[N\hat{V}] \rightarrow 0$ , it is sufficient to establish that  $\Omega$  is at most  $O(N^{-1})$ . Consider quadruples of dyads indexed by  $d, d', d'', d'''$  such that  $d \neq d' \neq d'' \neq d'''$ . Suppose there are  $Q$  quadruples and define  $p_Q(A) = |A|/Q$ . The covariance terms take positive values when  $I_{d,d'} = I_{d'',d'''} = 1$ , which occurs at rate  $O(N^{-2})$  (that is, when we have pairs of terms that are actually summed into  $\hat{V}$ ), and we have dependence across  $(d, d')$  and  $(d'', d''')$ . Then, the six distinct ways that such dependence can occur and the associated proportion of quadruples for which these occur for a sample of size  $N$  are as follows:

$$\begin{aligned}p_Q(\{d \neq d''\} \cap \{d'' : s(d) \cap s(d'') \neq \emptyset\}) &= O(N^{-1}), \\ p_Q(\{d \neq d'''\} \cap \{d''' : s(d) \cap s(d''') \neq \emptyset\}) &= O(N^{-1}), \\ p_Q(\{d \neq d'' \neq d'''\} \cap \{d'', d''' : s(d) \cap s(d'') \neq \emptyset \cap s(d) \cap s(d''') \neq \emptyset\}) &= O(N^{-2}), \\ p_Q(\{d' \neq d''\} \cap \{d'' : s(d') \cap s(d'') \neq \emptyset\}) &= O(N^{-1}), \\ p_Q(\{d' \neq d'''\} \cap \{d''' : s(d') \cap s(d''') \neq \emptyset\}) &= O(N^{-1}), \text{ and} \\ p_Q(\{d' \neq d'' \neq d'''\} \cap \{d'', d''' : s(d') \cap s(d'') \neq \emptyset \cap s(d') \cap s(d''') \neq \emptyset\}) &= O(N^{-2}).\end{aligned}$$

Therefore the proportion of quadruples yielding a positive covariance is the proportion of cases contributing non-zero values to the sum  $\Omega$ . This proportion is bounded by the proportion of cases where the indicator functions are both one and the order of the proportion of dependent quadruples from across the six cases characterized above, that is  $O(N^{-2}N^{-1}) = O(N^{-3})$ . Then  $\text{Var}[N\hat{V}]$  is at most  $O(N^2 D^{-4} D^4 N^{-3}) = O(N^{-1}) \rightarrow 0$ .

□

## C Proof of Proposition 2

*Proof.* Consider the interior (“meat”) of  $\hat{V}_r$ :

$$\hat{\Sigma}_r = \sum_{d=1}^D \left( X'_d e_d e'_d X_d + \sum_{d' \in \mathcal{S}(d)} X'_d e_d e'_{d'} X_{d'} \right) = \sum_{d=1}^D X'_d e_d e'_d X_d + \sum_{d=1}^D \sum_{d' \in \mathcal{S}(d)} X'_d e_d e'_{d'} X_{d'}.$$

Taking the second component,

$$\sum_{d=1}^D \sum_{d' \in \mathcal{S}(d)} X'_d e_d e'_{d'} X_{d'} = \sum_{i=1}^N C_i, \text{ with } C_i = \sum_{j \neq i} \sum_{k \neq i, j} X'_{d(i,j)} e_{d(i,j)} e'_{d(i,k)} X_{d(i,k)}.$$

Define

$$D_i = \sum_{j \neq i} X'_{d(i,j)} e_{d(i,j)} e'_{d(i,j)} X_{d(i,j)} \quad \text{and} \quad E_i = \sum_{j, k \neq i} \sum_{t=1}^{T(d(j,k))} X_{d(j,k)t} X'_{d(j,k)t} e_{d(j,k)t}^2.$$

Then working with  $\hat{\Sigma}_{C,i}$ ,  $\hat{\Sigma}_D$ , and  $\hat{\Sigma}_0$  as defined in the proposition,

$$\hat{\Sigma}_{C,i} = C_i + D_i + E_i, \quad 2\hat{\Sigma}_D = \sum_{i=1}^N D_i, \quad \text{and} \quad (N-2)\hat{\Sigma}_0 = \sum_{i=1}^N E_i.$$

Then

$$\begin{aligned} \hat{\Sigma}_r &= \hat{\Sigma}_D + \sum_{i=1}^N C_i \\ &= \hat{\Sigma}_D + \sum_{i=1}^N [C_i + D_i + E_i - D_i - E_i] \\ &= \sum_{i=1}^N \hat{\Sigma}_{C,i} - \hat{\Sigma}_D - (N-2)\hat{\Sigma}_0, \end{aligned}$$



and by linear distributivity for  $(\mathbf{X}'\mathbf{X})^{-1}$

$$\hat{V}_r = \sum_{i=1}^N \hat{V}_{C,i} - \hat{V}_D - (N-2)\hat{V}_0,$$

with  $\hat{V}_{C,i}$ ,  $\hat{V}_D$ , and  $\hat{V}_0$  as defined in the proposition. □

## D Weighted Observations

Weighting is a common way to adjust for unequal probability sampling of dyadic interactions, among other applications. The extension to the weighted case is straightforward. Assume weighted directed dyad observations with weights finite and fixed; denote the weight for dyad  $d$  as  $w_d$ . Define the sample data as  $\mathbf{X} = (X'_1 \dots X'_D)'$  and  $Y = (Y_1 \dots Y_D)'$  and the matrix of weights as  $\mathbf{W} = \text{diag}\{w_1, \dots, w_D\}$ . Then the weighted least squares estimator is

$$\hat{\beta}_w = (\mathbf{X}'\mathbf{W}\mathbf{X})^{-1}\mathbf{X}'\mathbf{W}Y.$$

**Corollary 2.** *For the weighted dyads case, assume the same conditions as in Proposition 1 and consider the following variance estimator*

$$\hat{V}_w = (\mathbf{X}'\mathbf{W}\mathbf{X})^{-1} \sum_{d=1}^D \left( X_d X'_d w_d^2 e_d^2 + \sum_{d' \in \mathcal{S}(d)} X_d X'_{d'} w_d w_{d'} e_d e_{d'} \right) (\mathbf{X}'\mathbf{W}\mathbf{X})^{-1}. \quad (7)$$

Then as  $N \rightarrow \infty$ ,

$$N\text{Var}[\hat{\beta}_w - \beta_w] - V_w \xrightarrow{P} 0,$$

and

$$N\hat{V}_w - V_w \xrightarrow{P} 0,$$

where

$$V_w = \frac{N}{D^2} \mathbb{E} [w_d X_d X_d']^{-1} \mathbb{E} \left( \mathbb{E} [X_d X_d' w_d^2 \text{Var}[\varepsilon_d | \mathbf{X}]] + \sum_{d' \in \mathcal{S}(d)} \mathbb{E} [X_d X_{d'}' w_d w_{d'} \text{Cov}[\varepsilon_d, \varepsilon_{d'} | \mathbf{X}]] \right) \mathbb{E} [w_d X_d X_d']^{-1}.$$

## E Generalized Linear Models

An implementation for generalized linear models follows the usual  $M$ -estimation results (Stefanski and Boos, 2002; Wooldridge, 2010, Ch. 12). Given an estimating equation,  $\psi(D; \theta)$ , on data  $D$ , and with parameters  $\theta$  and parameter estimates  $\hat{\theta}$ , the sandwich approximation for the variance is  $\mathbf{A}^{-1} \mathbf{B} \{\mathbf{A}^{-1}\}'$ , where

$$\mathbf{A} = \mathbb{E} \left[ -\frac{\partial}{\partial \theta'} \psi(D; \hat{\theta}) \right] \quad \text{and} \quad \mathbf{B} = \mathbb{E} \psi(D; \hat{\theta}) \psi(D; \hat{\theta})'.$$

For logistic regression the estimating equation is,

$$\psi((\mathbf{X}_d, Y_d); \hat{\beta}) = \mathbf{X}_d (Y_d - p_d),$$

where  $p_d = \text{expit}(\mathbf{X}_d \hat{\beta})$ , the predicted probability for dyad  $d$ . The plug-in variance estimator for logistic regression coefficients is given by

$$\hat{V}_l = (\mathbf{X}' \mathbf{M} \mathbf{X})^{-1} \sum_{d=1}^D \left( X_d X_d' r_d^2 + \sum_{d' \in \mathcal{S}(d)} X_d X_{d'}' r_d r_{d'} \right) (\mathbf{X}' \mathbf{M} \mathbf{X})^{-1}$$

where

$$\mathbf{M} = \text{diag}(p_d(1 - p_d))_{D \times D} \quad \text{and} \quad r_d = Y_d - p_d.$$

The extensions to repeated and weighted observations follow analogously.

## References

- Angrist, Joshua D. and Guido W. Imbens. 2002. "Comment on 'Covariance adjustment in randomized experiments and observational studies' by Paul R. Rosenbaum." *Statistical Science* 17(3):304–307.
- Angrist, Joshua D. and Jorn-Steffen Pischke. 2009. *Mostly Harmless Econometrics: An Empiricist's Companion*. Princeton, NJ: Princeton University Press.
- Arellano, Manuel. 1987. "Computing Robust Standard Errors for Within-Group Estimators." *Oxford Bulletin of Economics and Statistics* 49(4):431–434.
- Beck, Nathaniel, Kristian Skrede Gleditsch and Kyle Beardsley. 2006. "Space is More than Geography: Using Spatial Econometrics in the Study of Political Economy." *International Studies Quarterly* 50:27–44.
- Buja, Andreas, Richard Berk, Lawrence Brown, Edward George, Emil Pitkin, Mikhail Traskin, Linda Zhao and Kai Zhang. 2014. "Models as Approximations: A Conspiracy of Random Predictors and Model Violations Against Classical Inference in Regression." Manuscript, The Wharton School, University of Pennsylvania, Philadelphia.
- Cameron, A. Colin, Jonah B. Gelbach and Douglas L. Miller. 2011. "Robust inference with multi-way clustering." *Journal of Business and Economic Statistics* 29(2):238–249.
- Chamberlain, Gary. 1982. "Multivariate Regression Models for Panel Data." *Journal of Econometrics* 18(1):5–46.
- Conley, Timothy G. 1999. "GMM estimation with cross sectional dependence." *Journal of Econometrics* 92:1–45.
- Davidson, Russell and James G. MacKinnon. 2004. *Econometric Theory and Methods*. Oxford: Oxford University Press.
- Erikson, R. S., P. M. Pinto and K. T. Rader. 2014. "Dyadic Analysis in International Relations: A Cautionary Tale." *Political Analysis* 22(4):457–463.
- Fafchamps, Marcel and Flore Gubert. 2007. "The Formation of Risk Sharing Networks." *Journal of Development Economics* 83:326–350.
- Fisman, Raymond, Sheena S Iyengar, Emik Kamenica and Itamar Simonson. 2006. "Gender differences in mate selection: Evidence from a speed dating experiment." *Quarterly Journal of Economics* 121:673–697.
- Gelman, Andrew and Jennifer Hill. 2007. *Data Analysis Using Regression and Multi-level/Hierarchical Models*. Cambridge: Cambridge University Press.

- Goldberger, Arthur S. 1991. *A Course in Econometrics*. Cambridge, MA: Harvard University Press.
- Green, Donald P., Soo Yeon Kim and David H. Yoon. 2001. "Dirty Pool." *International Organization* 55(2):441–468.
- Greene, William H. 2008. *Econometric Analysis*. 6th ed. Upper Saddle River, NJ: Pearson.
- Hansen, Christian B. 2007. "Asymptotic properties of a robust variance matrix estimator for panel data when  $T$  is large." *Journal of Econometrics* 141:597–620.
- Hoff, Peter D. 2005. "Bilinear Mixed-Effects Models for Dyadic Data." *Journal of the American Statistical Association* 100(469):286–295.
- Hubbard, Alan E., Jennifer Ahern, Nancy L. Fliescher, Mark Van Der Laan, Sheri A. Lippman, Nicholas Jewell, Tim Bruckner and William A. Satariano. 2010. "To GEE or Not to GEE: Comparing Population Average and Mixed Models for Estimating the Associations Between Neighborhood Risk Factors and Health." *Epidemiology* 21(4):467–474.
- Huber, Peter J. 1967. The behavior of maximum likelihood estimates under nonstandard conditions. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability*. Vol. 1 pp. 221–233.
- Kenny, David A., Deborah A. Kashy and William L. Cook. 2006. *Dyadic Data Analysis*. New York, NY: Guilford Press.
- King, Gary and Margaret E. Roberts. 2014. "How Robust Standard Errors Expose Methodological Problems They Do Not Fix, and What to Do About It." *Political Analysis* (online early view):1–12.
- Lehmann, Erich L. 1999. *Elements of Large Sample Theory*. New York: Springer-Verlag.
- Liang, Kung-Yee and Scott L. Zeger. 1986. "Longitudinal data analysis using generalized linear models." *Biometrika* 73(1):13–22.
- Lin, Winston. 2013. "Agnostic Notes on Regression Adjustments to Experimental Data: Reexamining Freedman's Critique." *Annals of Applied Statistics* 7(1):295–318.
- MacKinnon, James G. and Halbert White. 1985. "Some Heteroskedasticity-Consistent Covariance Matrix Estimators with Improved Finite Sample Properties." *Journal of Econometrics* 29(3):305–325.
- Moulton, Brent R. 1986. "Random group effects and the precision of regression estimates." *Journal of Econometrics* 32:385–397.
- Neumayer, Eric and Thomas Pluemper. 2010. "Spatial Effects in Dyadic Data." *International Organization* 64(1):145–165.

- Russett, Bruce M. and John R. Oneal. 2001. *Triangulating Peace: Democracy, Interdependence and International Organizations*. New York, NY: Norton.
- Stefanski, Leonard A. and Dennis D. Boos. 2002. "The Calculus of M-Estimation." *The American Statistician* 56(1):29–38.
- Stock, James H. and Mark W. Watson. 2008. "Heteroskedasticity-Robust Standard Errors for Fixed Effects Panel Data Regression." *Econometrica* 76(1):155–174.
- White, Halbert. 1980a. "A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity." *Econometrica* 48(4):817–838.
- White, Halbert. 1980b. "Using Least Squares to Approximate Unknown Regression Functions." *International Economic Review* 21(1):149–170.
- White, Halbert. 1981. "Consequences and Detection of Misspecified Nonlinear Regression Models." *Journal of the American Statistical Association* 76(374):419–433.
- White, Halbert. 1982. "Maximum Likelihood Estimation of Misspecified Models." *Econometrica* 50(1-25).
- White, Halbert. 1984. *Asymptotic Theory of Econometricians*. New York, NY: Academic Press.
- Wooldridge, Jeffrey M. 2010. *Econometric Analysis of Cross Section and Panel Data*. Cambridge, MA: MIT Press.
- Zorn, Christopher. 2001. "Generalized Estimating Equation Models for Correlated Data: A Review with Applications." *American Journal of Political Science* 45:470–490.

# Cluster Robust Variance Estimation for Dyadic Data

Supporting Information Not for Publication

## Additional Simulation Evidence

Figure 2 and 3 show additional simulation results where we make comparisons to mixed effects models—specifically, to a two-way random effects model that incorporates random effects for each of  $N$  units in the data. All random effects models were fit using the `lmer` package in R. In both sets of graphs, black box plots show the distribution of standard error estimates for OLS coefficients using the proposed dyadic cluster robust estimator. Gray box plots show standard error estimates for OLS coefficients from a heteroskedasticity robust estimator. Blue box plots show the distribution of standard error estimators from a two-way random effects model with a random effect for each of  $N$  units. Black diamonds show the true standard error of the OLS coefficients and the blue diamonds show the true standard error of the random effects coefficients.




Figure 2: Normal shocks case (same as main text).

Figure 2 uses the same data generating process as was described in the main text, and presents the same results for OLS with either the dyadic cluster robust standard error estimator or the heteroskedastic robust estimator. In addition, we have added random effects estimates. Given that the data generating process conforms exactly to the assumptions of a two-way random effects model, we obtain the straightforward result that the random effects estimates are both more efficient (the blue diamonds tend to be lower than the black ones) and the random effects standard errors are consistent.

Figure 3 illustrates the robustness of the dyadic cluster robust estimator relative to a random effects estimator. In this case, we induce mild misspecification by assuming that the true data




Figure 3: Misspecification case.




Figure 4: Illustrative scatter plot and linear approximation for the misspecification case. The blue line is the true conditional mean, and the red line is the linear approximation.



generating process is

$$Y_{d(i,j)t} = \beta_0 + \beta_1 |X_i - X_j| + \beta_2 (X_i - X_j)^2 + \underbrace{\alpha_i + \alpha_j + v_{d(i,j)t}}_{\varepsilon_{d(i,j)t}},$$

where all parameters are set as in the main text but we also have  $\beta_2 = 0.25$ . We nonetheless fit a model that includes only  $|X_i - X_j|$ , thereby mildly misspecifying the model such that we do not account for the nonlinearity induced by the  $\beta_2 (X_i - X_j)^2$  term. Figure 4 shows the scatter plot of  $Y_{d(i,j)t}$  against  $|X_i - X_j|$ , the linear approximation (in red) for one of the simulation replicates ( $N = 50$ ), and the true conditional mean (in blue). Such linear approximations are the norm in social science research (Buja et al., 2014). In this case, we see that, despite the mild misspecification, the dyadic cluster robust standard error correctly characterizes the standard error of the linear approximation coefficients (the black box plots still zero in on the black diamonds). The random effects model is still more efficient (the blue diamonds still tend to be below the black diamonds). But the random effects standard error estimator (as computed in the `lmer` package for R) is inconsistent and anti-conservative: the blue box plots converge to a limit that is substantially below the true standard error (the blue diamonds).

## Speed Dating Application

Our second application is based on Fisman et al. (2006)’s seminal study of the determinants of mate selection in a “speed-dating” experiment. We replicate their primary analysis for female participants (as reported in column 1 of their Table III) using data from 21 dating sessions conducted by the experimenters.<sup>8</sup> These data include 278 women paired into 3,457 female-male dyads. Fisman et al. (2006) regress a binary indicator for whether the female subject desired contact information for a male partner on the subject’s ratings of the partner’s ambition, attractiveness, and intelligence based on a 10-point Likert scale. The regression controls for female-subject fixed effects and weights observations contributed by each female subject by the inverse of the number of partners with which she was paired. Table 2 shows the results of our re-analysis.<sup>9</sup> Again, we see that accounting for the dyadic clustering (column 4) yields standard error estimates that are larger than what one would get if one ignored all clustering (column 2), although in this case there are no pronounced differences with what one gets when one only accounts for within-subject clustering (column 3). This is to be expected given that the amount of dyadic dependence in this dataset is limited: dyads were formed only *within* sessions that included between 5 and 22 male partners for each female subject. In addition, the female subjects are never paired with each other. The only clustering that occurs is therefore within female subjects and then for male partners that appear across multiple female subjects. The former is addressed in part by the female subject-specific fixed effects. The latter is limited by the within-session pairings.

Table 2: Estimates and robust standard error estimates from a linear fixed-effects regression of an indicator for mate selection between dyads on dyad-level predictors, based on Fisman et al. (2006, Table III, Column 1).

Predictor of Mate Selection	(1)	(2)	(3)	(4)
	Coef.	Het. Robust S.E.	Naive Cluster Robust S.E.	Dyadic Cluster Robust S.E.
Ambition	0.0192	0.0052	0.0057	0.0061
Attractiveness	0.1157	0.0041	0.0051	0.0054
Intelligence	0.0465	0.0062	0.0073	0.0074
<i>N</i> = 3,457 dyad observations for 278 subjects.				

---

<sup>8</sup>The dataset to which we had access is not identical that used in the original Fisman et al. (2006) paper. Rather, it is the version made available in association with Gelman and Hill (2007, pp. 322-323), available at <http://www.stat.columbia.edu/~gelman/arm/software>. We contacted the original study authors and they indicated that the dataset from the original paper was not available.

<sup>9</sup>For the reasons explained in footnote 8, our estimates differ slightly from those that appear in the original paper.