

appeared responsible for PMC breakdown. The sterile plants also differed from the fertiles in the enlargement of the inner middle layer of the anther wall and in the lack of development of the endothecium. The different anther morphology of the present male sterility source could be attributed to these latter characteristics.

The numerous cytological and morphological differences between this male sterility source and that of the translucent type suggests that these two cases are conditioned by different genes that have different physiological effects. Detailed studies on microsporogenesis and anther wall development have been submitted for publication elsewhere.

- I.S. Dundas, K.B. Saxena, D.E. Byth, and E.S. Wallis (Department of Agriculture, University of Queensland)

Male Sterility Associated With Obcordate Leaf Shape in Pigeonpea

Male sterility associated with obcordate leaf shape (Fig.1) was observed during 1980-81 in



Fig. 1. Male sterile pigeonpea plant with obcordate leaves.

one of the F₃ progenies derived from a cross between a plant with obcordate leaf from a germplasm line GP-400 and Hy-3C. Twelve plants of the F₃ in that progeny exhibited obcordate leaf shape and pollen sterility ranging from 60% to 100%. In the F₂ progeny of another cross involving GP-400 and GP-127, all the plants with obcordate leaf showed high pollen sterility, providing us with a basis for assuming that the obcordate leaf shape is associated with male sterility. Flower drop was very high in the early stages but pod setting (by open pollination) increased as the temperature rose with the advancement of the season.

The widely opened, free and thread-like keel petals of the flowers in these male sterile plants mean that the gynoecium is well exposed to pollinators which may encourage cross-pollination and pod setting in these plants.

All these male sterile plants have been crossed with known parents and further studies are in progress to determine the cause, type, and inheritance of this male sterility.

- S. Venkateswarlu, A.R. Reddy, R. Nandan, O.N. Singh, and R.M. Singh (Pulse Research Laboratory, Banaras Hindu University)

A Cytoplasmic-Genetic Male Sterile Line in Pigeonpea

The use of existing stable genetic male sterile lines in the hybrid program poses two problems. First, as genetic male sterile lines must be maintained in the heterozygous state, it is necessary to identify and collect seeds from male sterile plants in the segregating populations. Secondly, in the hybrid seed production block the prompt identification and removal of about 50% normal fertile plants from within the female rows involves an additional cost of labor, land, and vigilance. To circumvent these problems, a search was made for a cytoplasmic-genetic male sterile system among the derivatives of *Cajanus cajan* x *Atylosia* spp. crosses. This was done by following up the clue given in the literature which indicates that alien cytoplasm in conjunction with the nucleus of cultivated types often produces this kind of male sterility.

Our past experience with the *Cajanus* x *Atylosia* spp crosses showed that, for most of the *Atylosia* species studied, crosses would

be successful only when *Cajanus* is used as the female parent. Therefore, to obtain inter-genetic derivatives with *Atylosia* cytoplasm, backcrosses were made by pollinating *A. scarabaeoides* with pollen from derivatives of the cross *Cajanus cajan* (cv T-21) x *A. scarabaeoides*. Similarly the backcross *A. sericea* x [*Cajanus cajan* (cv T-21) x *A. sericea*] was made. The backcross (BC) F₁s of both these crosses were normal.

A few BCF₂ male sterile plants from the *A. scarabaeoides* x (T-21 x *A. scarabaeoides*) cross that did not set seeds when selfed were noticed. This male sterility is associated with two kinds of abnormalities. In one case, the anthers are modified into petaloid structures (Fig.2) and the male sterility is linked with female sterility. In the second case, the male sterility is associated with free stamens and heterostyly, and the anthers appear morphologically normal. However, these normal-looking anthers are devoid of any viable pollen, so that rubbing these anthers when mature on to the stigma does not result in seed being formed. Histological studies revealed early degeneration of pollen mother cells and tapetal cells, indicating that there is little probability of viable pollen being produced by this male sterile line under any climatic conditions.

When the BCF₂s of this second type of male sterile line received pollen from the three pigeonpea cultivars, Pant A-2, Baigani, and ICP-6997, the resulting F₁s all segregated for male sterility. However, when these F₁s

were again backcrossed with their respective pigeonpea parents as the male parent, they produced male sterile plants solely. Thus it appears that this male sterility is governed by *A. scarabaeoides* cytoplasm. However, the fact that the F₂ male sterile plants segregated when crossed for the first time with the pigeonpea cultivars, suggests that this male sterility is cytoplasmic-genetic. Therefore, *A. scarabaeoides* should have the restorer gene(s) that is eliminated in the process of backcrossing with the pigeonpea cultivars.

In order to verify the above assumption, the male sterile line was crossed with *A. scarabaeoides*. All the F₁ plants were found to be normal, indicating that the male sterility is indeed cytoplasmic-genetic. The male sterility is being maintained by backcrossing the F₁ male sterile to the pigeonpea cultivar, Pant A-2 (as the male parent). This male sterile line was crossed with a series of pigeonpea genotypes. In the process C-11, a standard pigeonpea cultivar, has been found to restore fertility. This observation further confirms the cytoplasmic-genetic nature of this sterility. Unfortunately, this male sterility appears to be associated with varying degrees of female sterility which in turn results in poor pod-setting on the male sterile plants. Attempts are being made to transfer the sterile cytoplasm into different nuclear backgrounds with the hope of recovering male sterile genotypes with good pod-setting qualities.

- L.J. Reddy and D.G. Faris (ICRISAT)




Fig. 2. From left to right: Illustrations of anthers of normal, male sterile, and petaloid male sterile plants derived from *A. scarabaeoides* x (T-21 x *A. scarabaeoides*). The upper pictures show the general appearance of the flower with the petals removed. The lower pictures illustrate individual anthers.

Genetic Control of Sparse Pollen Production in Pigeonpea

Normally, pigeonpea anthers produce an abundance of pollen grains, sufficient to affect self- as well as insect-aided cross-fertilization. This note reports the occurrence of a "sparse pollen production" condition which can interfere with normal fertilization and result in low pod set.

This phenomenon was observed in an F₂ population of cross MS-4A x QPL-1 grown at the University of Queensland, Redland Bay Farm, Brisbane, Australia. As expected, this population segregated for fertile and translucent male sterile plants (3:1 ratio respectively). However, some fertile plants appeared macroscopically to produce relatively little pollen