



## Original Research Article

<https://doi.org/10.20546/ijcmas.2017.606.393>**Effect of temperature on *Rhizoctonia bataticola* and dry root rot in chick pea**P. Srinivas<sup>1\*</sup>, S. Ramesh Babu<sup>1</sup>, Mamta Sharma<sup>2</sup>, P. Narayan Reddy<sup>1</sup> and B. Pushpavathi<sup>1</sup><sup>1</sup>College of Agriculture, PJTSAU, Hyderabad, India<sup>2</sup>Legumes Pathology, ICRISAT, Hyderabad, India*\*Corresponding author***A B S T R A C T****Keywords**Chickpea,  
*Rhizoctonia solani*,  
Temperature**Article Info****Accepted:**

20 May 2017

**Available Online:**

10 June 2017

Chickpea dry root rot caused by *Rhizoctonia bataticola* (Taub.) Butler is a soil borne fungal pathogen causing significant yield losses due to change in environmental conditions. Influence of seven temperatures regimes (15°C, 20°C, 25°C, 30°C, 35°C, 40°C and 45°C) were tested on growth of *R. bataticola* isolates representing Madhya Pradesh, Karnataka, Maharashtra, Andhra Pradesh and Telangana states of India. The maximum mycelial growth was observed at 35°C followed by 30 and 25°C in all the isolates. The optimum temperature for dry root rot severity rating was at 35°C (8.5) followed by 30°C (7.9) followed by 25°C (7.0). Among the isolates, Telangana isolate was virulent and caused maximum disease severity.

**Introduction**

Chickpea (*Cicer arietinum* L.) is the premier pulse crop grown in more than 50 countries originated in south west Asia and is cultivated from ancient times both in Asia and European countries. It is the world's second most important food legume next to common bean. In India, chickpea is grown in an area of 10.22million hectares with a production of 9.53 million tonnes and productivity of 967kg<sup>ha</sup><sup>-1</sup> (Ministry of Agriculture, Govt. of India, 2013-14). Chickpea usually grown after rainy season on conserved soil moisture during winter in the tropics; in spring in the temperate and Mediterranean regions. Chickpea cultivation is often subjected to significant yield losses due to insects and diseases ranging from 5-10% in temperate and 50-100% in tropical regions (Van Emden *et al.*, 1988). Among the diseases of chickpea,

dry root rot is emerging as the most destructive constraint to chickpea productivity and production. Dry root rot caused by *Rhizoctonia bataticola* (Taub.) Butler [Pycnidial stage: *Macrophomina phaseolina* (Tassi) Goid] is a soil borne necrotrophic fungal pathogen that has a global distribution, which can infect more than 284 plant species throughout the world including monocot and dicots (Farr *et al.*, 1995). The disease is more prevalent during high temperature and low soil moisture conditions (Pande and Sharma, 2010). The most favorable temperatures for development of the *Rhizoctonia bataticola* ranged between 25 and 35°C.

In chickpea field, the onset of the disease appears as scattered drying of the plants. Affected plants are usually straw coloured,

but in some cases the lower leaves and stems show brown discoloration. The tap root appears black, rotten and devoid of most of the lateral and fine roots. The dead root become quite brittle and shows shredding of bark. Dark minute sclerotial bodies can be seen on the roots exposed or inside the wood. When the dry stem of the collar region is split vertically, sparse mycelium or minute sclerotia can be seen in the pith (Nene *et al.*, 1991). The effect of temperature on the growth of *Rhizoctonia bataticola* has been investigated by several authors, but up to now no satisfactory answer has been given on the optimal temperature conditions required for this necrotrophic pathogen on chickpea and its isolates from different locations.

In this paper, we report the response of isolates of *Rhizoctonia bataticola* at different temperatures on the growth and the disease severity on susceptible cultivar.

### **Materials and Methods**

A roving survey was conducted to collect dry root rot effected chickpea plants in central and southern parts of India *viz.* Madhya Pradesh, Karnataka, Maharashtra, Andhra Pradesh and Telangana. The Isolates collected from these locations were studied for pathogenicity on chickpea. The isolates representing five states which are showing disease severity more than seven rating were selected for the temperature studies. The five isolates selected were Rb 1 (Madhya Pradesh), Rb 2 (Karnataka), Rb 3 (Maharashtra), Rb 4 (Andhra Pradesh) and Rb 5 (Telangana).

### **Effect of temperature on growth of *Rhizoctonia bataticola***

Effect of seven different temperatures *viz.* 15°C, 20°C, 25°C, 30°C, 35°C, 40°C and 45°C were studied on growth of *Rhizoctonia bataticola* on potato dextrose agar. Mycelial

discs of 5mm diameter were cut from the edge of 3 days old culture of five isolates grown in 25°C were transferred to the center of 90mm Petri dish and incubated at different temperatures with 12h photoperiod. Each treatment was replicated three times in a completely randomized design. The average diameter of the fungal colony was recorded at 48h, 72h and 96h after incubation.

### **Effect of temperature on disease development**

Effect of different temperatures *viz.* 15°C, 20°C, 25°C, 30°C, 35°C, 40°C and 45°C on severity of dry root rot on susceptible genotype BG 212 with same five representative isolates was studied by adopting paper towel technique. Inoculum was prepared from the seven days old culture of *Rhizoctonia bataticola* isolates grown on 100ml potato dextrose broth medium (PDB). The culture of each isolate was grinded in a blender by adding 50ml of sterile distilled water to each fungal mat. Seven day old seedlings of BG 212 grown in sterilized sand were uprooted, washed under running water and were inoculated by dipping in the inoculum of *Rhizoctonia bataticola* isolates for 2 min. Inoculation was done through root dip inoculation technique.

Seedlings inoculated with sterile deionized water served as control. Seven to ten inoculated seedlings were placed in paper towel with the shoot left outside, folded, moistened and placed in trays. Trays were transferred to incubators maintained at seven different temperatures with 12h photoperiod and regularly moistened with sterile deionized water for seven days. At seven days after inoculation, the data on disease severity was recorded using 1 to 9 disease severity rating scale (Table 1) developed by Nene *et al.*, (1991).

## Results and Discussion

Temperature was known to have profound effect on the growth of fungal organism. Present studies were taken up to know the optimum, minimum and maximum temperature requirements for the growth of pathogen. Five isolates (Rb 1, Rb 2, Rb 3, Rb 4 and Rb 5) representing 5 different states were selected to conduct the study. All the isolates were grown at seven different temperatures *viz.*, 15°C, 20°C, 25°C, 30°C, 35°C, 40°C and 45°C.

At 48h after inoculation, among the seven different temperatures tested, the maximum colony growth was observed in Rb 3 (13.0mm) and no growth was observed in Rb 5 at 15°C whereas the maximum colony growth was observed in Rb 5 (38.3mm) and the least in Rb 1 (27.7 mm) at 20°C. Maximum colony growth of 54.0, 77.0 and 15.0 mm in Rb 3 and the least of 44.0, 62.3 and 3.7 mm in Rb 1 was recorded at 25, 30 and 40°C respectively. In contrast to this, maximum colony growth was observed in Rb 4 (85.3mm) and the least in Rb 1 (71.3 mm) at 35°C. The difference between isolates at different temperatures was significant except in Rb 2 and Rb 5 at 30 °C.

At 72h after inoculation, the maximum colony growth of 25.3 and 67.0 mm in Rb 3 and the least growth of 17.3 and 60.3 mm was observed in Rb 2 at 15 and 20°C respectively while at 25°C, maximum colony growth was observed in Rb 2 (84.7mm) and the least in Rb 1 (67.0 mm). At 30°C, maximum colony growth was observed in Rb 2, Rb 4, Rb 3 (90 mm) whose difference is at par with each other and the least in Rb 1 (82.0mm). At 35°C, colony growth reached its maximum (90mm) in all the isolates which were at par with each other. At 40°C, maximum colony growth was observed in Rb 3 (20.7mm) and the least in Rb 1 (8.3mm). There was

significant difference between isolates at 15, 20, 25 and 40°C except in Rb 2 and Rb 4.

At 96h after inoculation, maximum colony growth of 35.3 and 85.3 mm was recorded in Rb 3 while the least was observed in Rb 4 (20.3mm) and Rb 2 (74.7mm) at 15 and 20°C respectively. At 25°C, 30°C and 35°C, all the isolates had covered the entire 90 mm Petri plate. The difference between the isolates at 25, 30 and 35°C was at par with each other. Maximum colony growth was observed in Rb 3 (24.0mm) and the least in Rb 1 (10.7mm) at 40°C. No growth was observed till 96h after inoculation in any isolate and medium also became dry at 45°C.

The maximum colony growth was observed 72h after inoculation in all five isolates at 35°C. After 96h of incubation, all the isolates had covered Petri plates at 25°C and 30°C. The sclerotial initiation was started after 48 hours at 30°C and 35°C. The sclerotial initiation started 72h after inoculation at 25°C and it was observed at 96h after incubation in 20°C. At 15°C, it was observed that the growth was very slow and sclerotial initiation was observed after 144 hours after inoculation.

The above results were supported by Khan *et al.*, (2012) as they also observed pathogen growth over a wide range of temperature from 10<sup>0</sup>C to 45<sup>0</sup>C, but the optimum temperature for its growth was found to be 30<sup>0</sup>C. The next best temperature for its growth was recorded 35<sup>0</sup>C. Statistically the growth of the pathogen gradually decreased both at below 30<sup>0</sup>C and above 35<sup>0</sup>C. Patel and Patel (1990) also reported 35°C to be optimum temperature for growth and sclerotial formation of *M. phaseolina* in sesame. The results of Cesondes *et al.*, (2012) also showed that the cultures were well grown at 20°C, their colony size on the 3rd day were 14 times larger than at 10 and 15°C.

**Table.2** Colony diameter of *Rhizoctonia bataticola* isolates at different temperatures under *in vitro* condition

48 HAI (mm)*						
Temperature (°C)	Rb 1	Rb 2	Rb 3	Rb 4	Rb 5	Mean
15	4.0	3.3	13.0	6.7	0.0	5.4
20	27.7	32.7	37.0	31.7	38.3	33.5
25	44.0	53.7	54.0	51.3	50.0	50.6
30	62.3	71.3	77.0	75.0	71.0	71.3
35	71.3	82.0	80.0	85.3	77.7	79.3
40	3.7	9.3	15.0	13.7	10.3	10.4
45	0.0	0.0	0.0	0.0	0.0	0.0
<b>Mean</b>	30.4	37.7	39.4	36.5	35.3	
72 HAI (mm)						
Temperature (°C)	Rb 1	Rb 2	Rb 3	Rb 4	Rb 5	Mean
15	20.0	17.3	25.3	17.7	18.3	19.7
20	64.0	60.3	67.0	61.0	61.7	62.8
25	67.0	84.7	82.0	80.0	71.7	77.1
30	82.0	90.0	90.0	90.0	88.3	88.1
35	90.0	90.0	90.0	90.0	90.0	90.0
40	8.3	14.7	20.7	18.3	13.0	15.0
45	0.0	0.0	0.0	0.0	0.0	0.0
<b>Mean</b>	47.3	52.3	53.6	50.0	49.0	
96 HAI (mm)						
Temperature (°C)	Rb 1	Rb 2	Rb 3	Rb 4	Rb 5	Mean
15	25.3	21.3	35.3	20.3	24.3	25.3
20	80.3	74.7	85.3	78.3	75.0	78.7
25	90.0	90.0	90.0	90.0	90.0	90.0
30	90.0	90.0	90.0	90.0	90.0	90.0
35	90.0	90.0	90.0	90.0	90.0	90.0
40	10.7	17.3	24.0	20.0	15.3	17.5
45	0.0	0.0	0.0	0.0	0.0	0.0
<b>Mean</b>	55.2	57.0	59.2	55.2	54.9	
Factors	CD					
	48 HAI		72 HAI		96 HAI	
Temperature	0.64		0.54		0.55	
Isolate	0.54		0.46		0.47	
Temperature x Isolate	1.42		1.21		1.23	

HAI – Hours after inoculation \* mean of three replications

**Table.3** Disease severity (1-9 rating) of *Rhizoctonia bataticola* isolates on BG 212 at different temperatures

Isolates	Temperatures (°C)							Mean
	15	20	25	30	35	40	45	
Rb 1	1.7	3.1	6.7	7.4	8.3	9.0 <sup>#</sup>	9.0 <sup>#</sup>	6.5
Rb 2	1.3	3.0	7.2	8.2	8.7	9.0 <sup>#</sup>	9.0 <sup>#</sup>	6.6
Rb 3	1.3	2.9	7.0	7.9	8.3	9.0 <sup>#</sup>	9.0 <sup>#</sup>	6.5
Rb 4	1.7	3.2	6.5	8.0	8.1	9.0 <sup>#</sup>	9.0 <sup>#</sup>	6.5
Rb 5	1.0	3.4	7.4	8.2	9.0	9.0 <sup>#</sup>	9.0 <sup>#</sup>	6.7
control	1.0	1.0	1.0	1.0	1.0	1.0 <sup>##</sup>	1.0 <sup>##</sup>	1.0
Mean	1.4	3.1	7.0	7.9	8.5	9.0	9.0	
Factors		C.D.						
Temperature (T)		0.2						
Isolates (I)		0.17						
T x I		0.45						

<sup>#</sup>Mortality of plants due to combined effect of physiological wilting and Pathogen

<sup>##</sup>Plants showed physiological wilting

\* mean of three replications

**Table.1** Rating scale used to record disease severity of dry root rot in chickpea

Rating	Observation
1	No infection on roots
>1 - ≤3	Very few small lesions on roots
>3 - ≤5	Lesions on roots clear but small, new roots free from infection
>5 - ≤7	Lesions on roots many, new roots generally free from lesions
>7 - 9	Roots infected and completely discoloured

**Effect of temperature on development of disease**

There was very significant relation between the temperature and the development of the disease on the chickpea. The disease severity on the BG 212 cultivar was observed after seven days of incubation at different temperatures. The average disease severity at different temperatures was given in the Table 3. Maximum disease severity rating was recorded in Rb 1 and Rb 4 (1.7) and no symptom was observed in Rb 5 (1.0) inoculated plants at 15°C. In contrast, maximum disease severity rating was

recorded in Rb 5 (3.4) and the least in Rb 3 (2.9) inoculated plants at 20°C. Maximum disease severity rating of 7.4 in Rb 5 inoculated plants and the least in Rb 4 (6.5) and Rb 1 (6.7) which was at par with each other in inoculated plants at 25°C. At 30°C, maximum disease severity was observed in Rb 5 and Rb 2 (8.2) and the least in Rb 1 (7.4). At 35°C, maximum disease severity was observed in Rb 5 (9.0) and least was observed in Rb 4 (8.1) inoculated plants. At 40 and 45°C, mortality of plants was due to combined effect of physiological wilting and *Rhizoctonia bataticola* isolates. In control, plants were shriveled at 40°C whereas at 45°C

there was complete physiological wilting. It was also supported by the scarce and absence of mycelial growth in Petriplates at 40 and 45°C respectively in all the isolates.

The optimum temperature for dry root rot development was 35°C as maximum disease severity rating of 8.5 was observed irrespective of the isolate. This was followed by 30°C (7.9) followed by 25°C (7.0). It was observed that 20°C and 15°C had helped in the development of the lesions but could not develop further. The disease severity was very low 1.4 rating at 15°C while it was 3.1 at 20°C.

Among Rb 1, Rb 2, Rb 3 and Rb 4 isolates, there was no significant difference while Rb 5 was virulent isolate compared to others. The plants had showed symptoms on the tap root leaving the lateral roots unaffected at 25°C. At 30, 35, 40, 45°C, there was complete blackening of the roots and the reisolation from the roots showed the presence of *Rhizoctonia bataticola*. The uninoculated plants did not show any symptoms except at 40 and 45°C which showed physiological wilting and death of plants by complete drying respectively.

Similar observations were recorded by Sharma and Pande (2013) as disease incidence of dry root rot was significantly affected by high temperature. Out of five temperature levels viz., 15°C, 20°C, 25°C, 30°C and 35°C tested, chickpea predisposed to dry root rot early and severity was more at 35°C. Singh and Mehrotra (1982) observed increased levels of seed exudates when incubated at 35°C than at 15 and 25°C which contributed to increased pre-emergence damping off in gram seedling by *R. bataticola*. This study clearly demonstrated high temperature (35°C) was predisposing chickpea to *R. bataticola* infection, colonization and development of disease.

From the results we can say there is variation among the isolates of *Rhizoctonia bataticola* and temperature had profound effect on the growth and disease severity on chickpea. The optimum growth of the fungal isolates was found in 35°C. The next best temperature was 30°C followed by 25°C, 20°C and 15°C. There was very meager growth in 40°C and no growth was observed at 45°C.

## References

- Cesondes I, Cseh A, Taller J, Poczai, P. 2012. Genetic diversity and effect of temperature and pH on the growth of *Macrophomina phaseolina* isolates from sunflower fields in Hungary. *Molecular Biology Reports*. 39:3259-69.
- Farr, D.F., Bills, G.F., Chamuris, G.P and Rossman A.Y. 1995. *Fungi on plants and plant products in the United States*. 2nd ed. St Paul, MN: APS Press.
- Khan, R.A., Bhat, T.A and Kumar, K. 2012. Management of chickpea (*Cicer arietinum* L.) dry root rot caused by *Rhizoctonia bataticola* (Taub.) Butler. *International Journal of Research in Pharmaceutical and Biomedical Sciences*. 3(4): 1539-1548.
- Ministry of Agriculture. 2013-14. <http://agricoop.nic.in>
- Nene, Y.L., Reddy, M. V., Haware, M. P., Ghanekar, A. M and Amin, K. S. 1991. Field diagnosis of chickpea diseases and their control. *ICRISAT Information Bulletin*. 28: 52.
- Pande, S. and Sharma, M. 2010. Climate Change: Potential Impact on Chickpea and Pigeonpea Diseases in the Rainfed Semi-Arid Tropics (SAT). In: *5th International Food Legumes Research Conference (IFLRC V) & 7th European Conference on Grain Legumes (AEP VII) April 26-30, 2010*. Antalya, Turkey.



- Patel, K.K and Patel. A.J. 1990. Meterological correlation of charcoal rot of sesamum. *Indian Journal of Mycology and Plant Pathology*. 20(1): 64-65.
- Sharma, M and Pande, S. 2013. Unraveling effects of temperature and soil moisture stress response on development of dry root rot [*Rhizoctonia bataticola* (Taub.)] Butler in chickpea. *American Journal of Plant Sciences*. 4: 584-589.
- Singh, P. J and Mehrotra, R.S. 1982. Influence of soil moisture and temperature on *Rhizoctonia bataticola* infection of gram. *Indian Phytopathology*. 35 (2): 327-329.
- Van Emden, H.F., Ball, S.L and Rao, M.R. 1988. Pest diseases and weed problems in pea, lentil and faba bean and chickpea. In *summer field R.J. (ed.), World crops: cool season food legumes. Dordrecht: Kluwer*. 519-534.

**How to cite this article:**

Srinivas P., S. Ramesh Babu, Mamta Sharma, P. Narayan Reddy and Pushpavathi B. 2017. Effect of temperature on *Rhizoctonia bataticola* and dry root rot in chick pea. *Int.J.Curr.Microbiol.App.Sci*. 6(6): 3349-3355. doi: <https://doi.org/10.20546/ijemas.2017.606.393>