

Identification of stable genotypes and genotype by environment interaction for grain yield in sorghum (*Sorghum bicolor* L. Moench)

Massaoudou Hamidou^{1*}, Oumarou Souleymane¹, Malick N. Ba², Eric Yirenkyi Danquah³, Issoufou Kapran¹, Vernon Gracen³ and Kwadwo Ofori³

¹Departement des cultures pluviales, Institut National de la Recherche Agronomique du Niger (INRAN), BP: 429 Niamey, Niger, ²International Crop Research Institute for the Semi-Arid Tropics (ICRISAT), BP 12404 Niamey, Niger and ³West Africa Centre for Crop Improvement, College of Basic and Applied Sciences, University of Ghana Legon, Accra, PMB: 30, Ghana

Received 9 June 2018; Accepted 5 October 2018

Abstract

Sorghum is a staple food crop in Niger and its production is constrained by sorghum midge and the use of low yielding, local sorghum varieties. To improve sorghum productivity, it is crucial to provide farmers with high yielding sorghum cultivars that are resistant to midge. We evaluated 282 genotypes in four environments of Niger Republic. Alpha (0.1) lattice with two replications was the experimental design. Genotype and genotype by environment (GGE) biplot analysis was used to study grain yield (GY) stability and G × E interactions. The results revealed that two distinct mega environments were present. Genotype L232 was the best genotype for GY in the first planting date at Konni and the first and second planting dates (PDs) at Maradi. Genotype L17 was the best for GY in the second PD at Konni. The second PD at Konni was the most discriminating environment while the first PD at Konni is suitable for selecting widely adapted genotypes for GY.

Keywords: GGE biplot, Niger, planting dates, yield stability

Introduction

Sorghum (*Sorghum bicolor* L. Moench) is the second cereal crop in Niger after pearl millet. It is used in human as well as animal nutrition. In Niger, sorghum production rarely meets the demand of the growing population. Grain sorghum yields are very low, about 0.28 t/ha, which is far below the genetic potential of the crop compared with countries like the USA (4.3 t/ha), Argentina (4.9 t/ha) and China (3.2 t/ha) (FAOSTAT, 2014). The low production is attributed to abiotic and biotic stresses such as sorghum midge *Stenodiplosis sorghicola*. According to Hamidou (2016), sorghum midge is a panicle insect found in most of sorghum growing environments in Niger where it causes high sorghum grain yield (GY) reduction. In Niger, delay or advance sorghum planting time is one of midge control

means used by farmers (Hamidou, 2016). However, one cannot predict the rainy season in the Sahelian countries like Niger. Hence, early or late environmental stress could drastically impact sorghum GY for early or late planted fields. According to Menezes *et al.* (2015), stress caused by environmental factors on final yield of sorghum depends on the growth stage of the crop at which the stress occurs and the genotype being cultivated. The effect of environmental conditions may vary depending on planting date (PD). According to Diawara (2012), PD has significant influence on GY in sorghum. Early or late planting increases the risk of poor crop establishment resulting in low yield with early planting and water or heat stress at reproductive stages with late planting. The use of indigenous varieties with low-yielding capacity also limits sorghum productivity in Niger. Farmers mostly rely on low-yielding landraces, so sorghum production fails to meet the demand of increasing population and food insecurity remains a major issue (Maman *et al.*, 2004). Variation in performance

*Corresponding author. E-mail: hmassaoud@yahoo.fr

caused by interaction of genotype and environment on farms is probably the main reason why traditional plant breeding has failed to fully benefit resource-poor farmers, especially in marginal and fragile environments (Ceccarelli and Grando, 2007). To stabilize sorghum production in Niger, identification of good yielding and stable varieties is necessary. Genotype by environment interaction can be evaluated using GGE biplots which graphically display the genotype by environment interaction in a two-dimensional space thereby giving a visual presentation of the relationships between genotypes and environments (Yan *et al.*, 2000). The objective of this study was to identify new varieties of sorghum with high and stable yield across environments.

Materials and methods

Experimental germplasm

Two hundred and eighty recombinant F5 inbred lines and two local checks were evaluated. The F5 lines were obtained by crossing a local sorghum variety (MDK) to an exotic sorghum midge resistant cultivar from ICRISAT (ICSV88032) and the progeny were advanced using single-seed descent from F1 to F5. The local variety has white grain with good qualities and is widely cultivated by sorghum farmers in Niger. However, this variety is photosensitive and highly susceptible to sorghum midge.

Experimental sites

The study was done during 2015 rainy season at the research stations of INRAN at Konni and Maradi. Both locations are sorghum midge hotspots in Niger. Konni has a latitude of 13°47'23" North and a longitude of 5°14'57" East and the average annual rainfall of 589.7 mm with average temperature of 29.3°C. Maradi has a latitude of 13°18'25" North and a longitude of 7°09'35" East and the average annual rainfall of 537.4 mm with average temperature of 20.5°C.

Experimental design

The experimental design was an alpha (0.1) lattice with two replications in two different PDs, giving four environments. However, blocking was not significant and the dates were reanalysed using randomized complete block design. Environments 1, 2, 3 and 4 were the first PD at Konni, the second PD at Konni, the first PD at Maradi and the second PD at Maradi, respectively. Two different PDs were used to simulate early or late beginning of the rainy season. The early planting was done upon the first rainfall, while the late planting was done 3 weeks later. Each genotype was

Table 1. Combined mean squares for GY measured in 282 sorghum genotypes evaluated in 2015 rainy season

Source of variation	df	GY ^a
Replications	1	1,234,336*
Entries	281	177,377**
PDs	1	3,548,702**
Sites	1	30,681,580**
Entries × PD	281	129,768**
Entries × sites	281	119,234**
PD × sites	1	7,723,780**
Entries × PD × sites	281	104,061**
Residual	1127	62,286

^aGY (t/ha).

***Significant at the 5 and 1% probability levels, respectively.

grown in a single row of 3 m; the intra- and inter-row spacing was 0.20 m × 0.80 m. The material was subjected to natural infestation of sorghum midge. In order to evaluate midge damage on the panicles, three panicles were covered at emergence using selfing bags. At harvesting, panicle and grain mass were recorded for the three covered and three non-covered panicles. The loss in GY in three non-covered panicles was expressed as a percentage of GYs in covered panicles.

Data collection and analysis

Data were collected on GY. The GY was measured in t/hectare adjusted to grain moisture content at 12%. GGE biplot analyses in GenStat version 18 were performed to identify high yielding and stable cultivars for GY across the four environments.

Results

Combined analysis of variance showed significant differences among genotypes for GY (Table 1). The GY varied from 3.67 t/ha to 0.14 t/ha in the first PD at Konni and from 5.51 t/ha to 0.08 t/ha in the second PD. At Maradi, the yield performance varied from 4.03 t/ha to 0.05 t/ha in the first PD and from 3.52 t/ha to 0.00 t/ha in the second PD (Table 2).

Two mega-environments were observed from the GGE biplot. Environments 1, 3 and 4, representing early planting at Konni, early planting at Maradi and late planting at Maradi, formed one mega-environment while environment 2, representing late planting at Konni, is a different mega-environment. Mega-environment 2 was more discriminating for GY. Genotypes L232, L17, L207 and L75 were superior genotypes in the evaluation (Fig. 1).

Table 2. GY performance of the top 20 and bottom five genotypes out of 282 sorghum genotypes evaluated in 2015 rainy season

Environment 1 ^a		Environment 2 ^b		Environment 3 ^c		Environment 4 ^d	
Top 20 genotypes	GY (t/ha)	Top 20 genotypes	GY (t/ha)	Top 20 genotypes	GY (t/ha)	Top 20 genotypes	GY (t/ha)
L232	3.67	L17	5.51	L232	4.03	L232	3.52
L219	3.64	L207	5.16	L19	3.49	L84	3.25
L202	3.60	L23	5.06	L140	3.11	L115	2.44
L268	3.56	L75	4.80	L17	2.68	L127	2.32
L17	3.39	L168	4.75	L49	2.44	L162	2.23
L227	3.30	L64	4.72	L250	2.40	L234	1.91
L72	3.23	L202	4.67	L54	2.36	L17	1.91
L118	3.18	L182	4.46	L182	2.36	L86	1.74
L123	2.90	L177	4.07	L43	2.35	L207	1.52
L158	2.85	L60	3.92	L281	2.20	L72	1.51
L116	2.80	L249	3.90	L174	2.19	L47	1.47
L68	2.72	L185	3.79	L128	2.10	L148	1.46
L258	2.68	L11	3.75	L200	2.05	L60	1.43
L59	2.61	L5	3.73	L123	2.03	L257	1.41
L215	2.60	L170	3.72	L219	2.01	L54	1.31
L233	2.55	L52	3.50	L41	1.99	L41	1.31
L106	2.49	L244	3.43	L277	1.93	L132	1.30
L112	2.49	L250	3.30	L210	1.91	L277	1.29
L107	2.46	L236	3.28	L257	1.91	L109	1.28
L271	2.45	L100	3.27	L248	1.87	L113	1.28
Check	1.21	Check	0.28	Check	1.10	Check	0.19
Bottom five genotypes	GY (t/ha)	Bottom five genotypes	GY (t/ha)	Bottom five genotypes (t/ha)	GY (t/ha)	Bottom five genotypes	GY (t/ha)
L211	0.25	L110	0.16	L271	0.19	L89	0.00
L120	0.22	L190	0.15	L246	0.18	L106	0.00
L2	0.21	L55	0.14	L268	0.14	L141	0.00
L88	0.20	L263	0.09	L89	0.07	L173	0.00
L176	0.14	L277	0.08	L192	0.05	L224	0.00
LSD at 5%^e	507.6		736.5		390.21		146.91

^aFirst PD at Konni.

^bSecond PD at Konni.

^cFirst PD at Maradi.

^dSecond PD at Maradi.

^eLeast significant difference.




Fig. 1. Which-won-where view of GGE biplot of GY of different 282 sorghum genotypes evaluated in 2015 rainy season.




Fig. 2. Scoring of 282 sorghum genotypes and the four different evaluation environments.

GGE biplot analysis helps in identifying stable genotypes associated with the best environments. Environments 1, 3 and 4, representing the early planting at Konni, early and late planting at Maradi, were located below the average environment coordinate (AEC). Environment 2, representing

the late planting at Konni, was located above the AEC (Fig. 2). Genotypes L232, L17, L182, L202, L168 and L64 were above the AEC and closer to average environment axis than other genotypes (Fig. 2) indicating they are the most stable genotypes.




Fig. 3. Scoring of four evaluation environments based on an ideal environment (a) and the 282 sorghum genotypes based on ideal genotypes (b).

Environment 1, representing the early planting at Konni, was the best environment for GY, followed by environment 3, representing the early planting at Maradi, environment 4, representing the late planting at Maradi and environment 2, representing the early planting at Konni (Fig. 3(a)).

The arrow pointing at the middle of the concentric circle locates the ideal genotypes. Genotypes closest to the arrow are the best ones. L232 was the best genotype followed by L17, L182, L202, L168 and L64 (Fig. 3(b)).

Discussion

According to the analysis of variances, germplasms were genetically diverse and considerable amount of variability existed in the materials used for this study. This indicates that selection among these germplasms can be successful. Hence the observed variability can be exploited by plant breeder to undertake further hybridization activities. GGE biplots provide information on the mean expression of a trait and its stability. They help to identify the best environments as well as the best genotypes. In this study, two distinct groups of environments for GY evaluation were observed. The first PD at Konni and first and second PDs at Maradi were similar but the second PD at Konni was different from the three other environments. However, the second PD at Konni should be primarily used in the evaluation of sorghum genotypes for GY since it was the most discriminating environment. In an earlier study,

Teodoro *et al.* (2016) reported two mega-environments in sorghum evaluated over five different environments; while De Figueiredo *et al.* (2015) documented several mega-environments for green mass yield and total soluble solids in sweet sorghum using GGE biplot analysis.

In terms of GY performance, genotype L232 in environments 1, 3 and 4 and genotype and genotype L17 in environment 2 were the leading genotypes. L232 had 3.67, 4.03 and 3.52 t/ha in environments 1, 3 and 4, respectively, while genotype L17 recorded 5.51 t/ha in environment 2. This makes the two genotypes to be identified as the best performing entries of this study. Al-Naggar *et al.* (2018) documented four genotypes based on GY performance and stability.

The second PD at Konni is more discriminating and more representative for GY evaluation. This could be due to the occurrence of higher sorghum midge pressure when there is delay in sorghum planting time in this sorghum midge hotspot. Therefore, this environment should be regarded as the best environment for selecting sorghum for yield under midge infestation. Evaluating sorghum genotypes in this environment could help minimize the cost for experimenting sorghum for high GY in several environments. On the other hands, genotype L17 was the most stable genotype in this environment. Overall, genotype L17 is the best variety identified here because it is high yielding and stable for late planting at Konni where midge pressure is expected to be high. In a similar study, Ezzat *et al.* (2010) studied yield stability in sorghum

genotypes and identified four genotypes as stable and high yielding.

The first PD at Konni is the best environment for selecting widely adapted genotypes for GY because of low midge pressure. In this environment, L232, L17, L182, L202, L168 and L64 were found to be the best performing genotypes. Based on actual yield and stability, genotypes L202 and L17 were the best performers in the first and second PDs at Konni and genotypes L232, L17 and L182 were the best in the first PD at Maradi where there is lower midge pressure. Genotypes L232 and L17 were the best in the second PD at Maradi where midge pressure was the greatest. Ezzat *et al.* (2010) reported four sorghum genotypes as promising cultivars based on GY and yield-stability analysis. Adugna (2007) recommended four sorghum genotypes out of 15 evaluated in eight environments for growing in Ethiopian dry lowland areas.

Conclusions

Stable genotypes adapted across environments for GY were identified. Environments suitable for evaluating sorghum germplasm for GY were also identified. Overall, lines L232 and L17 were found to be stable for GY under both non-midge and midge pressure conditions whereas the L182, L202 L168 and L64 performed well under non-midge pressure conditions. These lines are promising for high-grain yielding and adaptation in sorghum production areas. These genotypes can also be used as parental materials for GY and/or midge resistance in sorghum improvement programme.

Acknowledgements

The authors acknowledge Alliance for Green Revolution in Africa (AGRA) for funding this work. This work has also been undertaken as part of the CGIAR Research Program on Dryland Cereals.

References

- Adugna A (2007) Assessment of yield stability in sorghum. *African Crop Science Journal* 15: 167–171.
- Al-Naggar AMM, Abd El-Salam RM, Hovny MRA and Walaa YSY (2018) Genotypexenvironment interaction and stability of *Sorghum bicolor* lines for some agronomique and yield traits in Egypt. *Asian Journal of Agricultural and Horticultural Research* 1: 1–14.
- Ceccarelli S and Grando S (2007) Decentralized-participatory plant breeding: an example of demand driven research. *Euphytica* 155: 349–360.
- De Figueiredo UJ, Nunes JAR, Parrella RDC, Souza ED, da Silva AR, Emygdio BM, Machado JRA and Tardin FD (2015) Adaptability and stability of genotypes of sweet sorghum by GGE biplot and Toler methods. *Genetics and Molecular Research* 14: 11211–11221.
- Diawara B (2012) *Effect of planting date on growth, development and yield of grain sorghum hybrids*. M.Sc. Thesis, Kansas State University.
- Ezzat EM, Ali MA and Mahmoud AM (2010) Agronomic performance, genotypexenvironment interactions and stability analysis of grain sorghum (*Sorghum bicolor* L. Moench). *Asian Journal of Crop Science* 2: 250–260.
- FAOSTAT (2014) Statistical Yearbook. World Food and Agriculture. *FAO Food Agric. Organization UN Rome Italy*. Available at: <http://faostat.fao.org/site/291/default.aspx> (Accessed 19 March 2015).
- Hamidou M (2016) *Breeding sorghum (Sorghum bicolor L. Moench) for high grain yield and resistance to sorghum midge (Stenodiplosis sorghicola Coquillett) in Niger*. Ph.D. Thesis, University of Ghana.
- Maman N, Mason SC, Lyon DJ and Dhungana P (2004) Yield components of pearl millet and grain sorghum across environments in the Central Great Plains. *Crop Science* 44: 2138–2145.
- Menezes CB, Saldanha DC, Santos CV, Andrade LC, Júlio MM, Portugal AF and Tardin FD (2015) Evaluation of grain yield in sorghum hybrids under water stress. *Genetics and Molecular Research* 14: 12675–12683.
- Teodoro PE, Almeida FJ, Daher RF, Menezes CB, Cardoso MJ, Godinho VP and Tardin FD (2016) Identification of sorghum hybrids with high phenotypic stability using GGE biplot methodology. *Genetics and Molecular Research: GMR* 15: 234–239.
- Yan W, Hunt LA, Sheng Q and Szlavnic Z (2000) Cultivar evaluation and mega-environment investigation based on the GGE biplot. *Crop Science* 40: 597–605.