

BMJ Open Modelling the effect of the introduction of antenatal screening for group B *Streptococcus* (GBS) carriage in the UK

David Bevan,¹ Alicia White,² John Marshall,³ Catherine Peckham⁴

To cite: Bevan D, White A, Marshall J, *et al.* Modelling the effect of the introduction of antenatal screening for group B *Streptococcus* (GBS) carriage in the UK. *BMJ Open* 2019;**9**:e024324. doi:10.1136/bmjopen-2018-024324

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-024324>).

Received 21 May 2018

Revised 22 October 2018

Accepted 25 January 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Health and Human Services, Melbourne, Australia

²Bazian Ltd, London, UK

³UK National Screening Committee, London, UK

⁴Department of Paediatric Epidemiology Unit, University College London, London, UK

Correspondence to

John Marshall;
John.Marshall@phe.gov.uk

ABSTRACT

Objectives To estimate the potential impact of the addition of culture-based screening for group B streptococcus (GBS) carriage in pregnancy to a risk-based prevention policy in the UK. We aimed to establish agreement within a multidisciplinary group of key stakeholders on the model input parameters.

Design Deterministic model using a consensus approach for the selection of input parameters.

Setting and participants A theoretical annual cohort of 711 999 live births in the UK (excluding births by elective caesarean section).

Interventions Culture-based screening for GBS at 35–37 weeks of pregnancy added to the recommended risk-based prevention policy in place on the date of modelling.

Outcome measures Outcomes assessed included use of intrapartum antibiotic prophylaxis (IAP), early onset GBS (EOGBS), EOGBS mortality, severe EOGBS-related morbidity and maternal penicillin anaphylaxis.

Results With no prophylaxis strategy, the model estimated that there would be 421 cases of culture positive EOGBS in a year (0.59/1000 live births). In the risk-based prophylaxis scenario, 30 666 women were estimated to receive IAP and 70 cases of EOGBS were prevented. Addition of screening resulted in a further 96 260 women receiving IAP and the prevention of an additional 52 to 57 cases of EOGBS. This resulted in the prevention of three EOGBS deaths and four cases of severe disability. With screening, an additional 1675 to 1854 women receive IAP to prevent one EOGBS case and 24 065 to 32 087 receive IAP to prevent one EOGBS death.

Conclusions The evidence base available for a broad range of model input parameters was limited, leading to uncertainty in the estimates produced by the model. Where data was limited, the model input parameters were agreed with the multidisciplinary stakeholder group, the first time this has been done to our knowledge. The main impact of screening is likely to be on the large group of low-risk women where the clinical impact of EOGBS tends to be less severe. This model suggests that the reduction in mortality and severe disability due to EOGBS with antenatal GBS screening is likely to be very limited, with a high rate of over-detection and overuse of antibiotics.

INTRODUCTION

Group B *Streptococcus* (GBS) is a bacterium which can be commonly found in the digestive system and female reproductive tract. It

Strengths and limitations of this study

- Model structure and input parameters were agreed to by a multidisciplinary stakeholder group using a consensus building mechanism agreed to in advance.
- Deterministic one-way sensitivity analyses were carried out to assess the impact of changes to key model inputs.
- Systematic review to support each model input was not feasible; data sources were identified through broad systematic searches for recently published evidence. Higher quality evidence, such as systematic reviews, was used where available, as well as national sources including guidance and the UK National Screening Committee's reviews.
- The modelling approach used was pragmatic and deterministic and so it does not provide CIs for model outputs.
- The evidence base available for many input parameters was limited and this leads to uncertainty in the model's predictions.

can be transmitted from a pregnant carrier to her newborn, typically during vaginal delivery.¹

While the bacterium does not usually cause harm, in some cases it can cause early onset infection in the newborn's first week of life. This is termed early onset GBS (EOGBS) disease. In the UK, GBS is the most common cause of neonatal sepsis and meningitis.^{2,3}

Comprehensive surveillance established an overall rate of EOGBS at 0.48 per 1000 live births in the UK and Ireland in 2000–2001.⁴ Since then, routine laboratory surveillance has shown a fluctuation in incidence of culture positive EOGBS with slight increases between 2000 and 2010.⁵ Preliminary national surveillance data from 2014 to 2015 suggests that the overall incidence of EOGBS was 0.57 per 1000 live births in the UK and Ireland.⁶ Clindamycin had been used as the main alternative to penicillin prophylaxis in women with penicillin allergy; however, since 2000 there has been a marked increase

in resistance to clindamycin in patients of all ages.⁵ As a result, the most recent guidance from the Royal College of Obstetricians and Gynaecologists (RCOG), published in 2017, has recommended that clindamycin should no longer be used for this purpose.⁷ Vancomycin is now recommended by the RCOG as the antibiotic of choice for women with severe allergy to penicillin.

In the UK, the mortality rate in term newborns with EOGBS is estimated to be between 6% and 10.6%,^{4 8} with a similar proportion left with severe morbidity.^{8 9} In pre-term newborns with EOGBS, the mortality rate is reported to be higher.¹⁰

Antenatal screening for GBS

Pregnant women can be screened for GBS carriage in late pregnancy. Screening involves the collection of specimens using vaginal and rectal swabs which are processed using selective culture media.^{11 12} The purpose of screening is to identify a group of women who are eligible for intravenous intrapartum antibiotic prophylaxis (IAP) as a means of preventing EOGBS disease.¹¹ The mainstay of IAP is benzylpenicillin.

In the UK, currently there is agreement between the guidance issued by the UK National Screening Committee (UK NSC), the National Institute for Health and Care Excellence (NICE) and the RCOG that routine screening for GBS carriage should not be offered.^{6 7 13}

A number of maternal risk factors for EOGBS have been identified. These include having a baby with GBS in a previous pregnancy, incidentally detected maternal GBS carriage, prematurity, prolonged membrane rupture and suspected infection in labour. In the UK, at the time of this modelling exercise (2014–2015), GBS IAP was recommended for women with either of the first two listed risk factors and broad spectrum antibiotics, with an agent active against GBS, recommended for women with suspected infection in labour.^{13 14} Since the completion of the modelling exercise, the RCOG has updated its guideline on GBS.⁷ The main change is that they now recommend offering GBS IAP to all women in confirmed pre-term labour. This is due to the increased risk of EOGBS and mortality in pre-term infants compared with term infants.

In 2017, the UKNSC completed the process of reviewing the evidence on antenatal GBS screening as part of its triennial review process.¹⁵ This review concluded that the recommendation not to implement screening in the UK should not be changed.⁶ This was because the committee considered there to be insufficient evidence on the balance of benefits and harms from culture-based screening and treating women with positive results with IAP.

The discussion on antenatal screening for GBS has taken place in a UK policy context shaped by the Maternity Review,¹⁶ National Health Service Outcomes Frameworks¹⁷ and the Antimicrobial Resistance Strategy.¹⁸ These major policy drivers emphasise issues ranging from patient choice and experience, place of birth, reduction

of neonatal deaths, antibiotic stewardship and reduction of antibiotic usage. The potential impact of screening on these issues, therefore, needs to be considered.

The charity, Group B Strep Support (GBSS), has campaigned for screening since the 1990s. The controversy surrounding the screening policy is reflected in journal debate,^{19–22} and politicians from the main parties in the four UK countries have signed petitions, raised parliamentary questions and led delegations on this issue. A petition in favour of screening, with over 250 000 signatures, was delivered to the ministers and senior officials in January 2017.²³

In part, this interest is stimulated by the implementation of antenatal screening for maternal GBS carriage in a number of developed countries. With few exceptions, retrospective cohort studies from these countries report a decline in the rate of culture positive EOGBS following the introduction of screening.²⁴ For example, surveillance data from the USA reports reduced rates of EOGBS following implementation of each new policy change.¹¹ The retrospective and observational design of these studies makes it difficult to ascertain if the data are complete, or if the reduction is conclusively attributable to screening alone.^{6 25}

In addition, the absence of reports on the effect of screening on the rate of culture-negative sepsis presumed to be due to EOGBS disease limits interpretation further.^{19 26} The difficulty in extrapolating data from different geographical settings has also been noted in relation to EOGBS.²⁷ However, the absence of suitably powered UK studies necessitates the use of available data from the UK and other countries, to estimate the potential impact of screening in the UK.

The UK NSC therefore convened a multidisciplinary, multi-agency expert group in 2014 to consider the available evidence to inform the development of a model to estimate the preventive potential of screening when added to current clinical practice. A pivotal aim of the process was to establish a shared set of assumptions among key stakeholders on a controversial topic.

METHODS

Model structure

A pragmatic deterministic model was developed in Microsoft Excel 2010 to simulate two scenarios in a 1-year UK pregnancy cohort. The first scenario was based on the risk-based management pathway recommended in the UK at the time of the model development (2014–2015).^{13 14} The second scenario was the screening and risk-based scenario. In this scenario, antenatal culture-based screening for GBS was offered at 36 weeks of pregnancy to women not already identified as being at risk through the risk-based strategy. This scenario was based broadly on the screening strategy recommended by the US Centers for Disease Control and Prevention¹¹ and the existing UK risk-based approach. This meant that women with known risk factors who should already be offered




Figure 1 Flow of the maternity cohort into antenatal and intrapartum risk factor groups. EOGBS, early onset GBS; IAP, intrapartum antibiotic prophylaxis; PROM, pre-labour rupture of membranes.

IAP under UK guidelines continued to be offered IAP without screening. Women without these risk factors were offered screening. Women giving birth pre-term, before 37 weeks, were assumed not to receive screening. This was due to the timing of the screening test at 36 weeks and the logistics of transport and laboratory processing time.

Within both scenarios, women were sequentially divided into mutually exclusive groups based on various clinical parameters.

In scenario one, the existing risk-based approach, the clinical characteristics on which the sequential divisions were based were as follows:

- ▶ Mode of birth (elective caesarean section or not).
- ▶ Presence of antenatal risk factors for EOGBS (a previous baby with EOGBS, incidental detection of GBS carriage or no risk factors).
- ▶ Timing of birth (term or pre-term).
- ▶ Presence of intrapartum risk factors for EOGBS (pre-term pre-labour rupture of membranes, suspected infection during labour, or ‘uncomplicated birth’, that is, no intrapartum risk factors).

This resulted in the population being divided into 15 ‘clinical risk groups’. These are depicted in [figure 1](#).

In scenario 2, the screening and risk-based approach, women who had no antenatal risk factors for EOGBS and who gave birth at term were eligible for screening. The population eligible for screening is depicted in [figure 2](#) and was divided into groups based on:

- ▶ Receipt of screening (yes or no).
- ▶ Result of screening (positive or negative for GBS).
- ▶ GBS status at delivery (positive or negative for GBS).
- ▶ Presence of intrapartum risk factors for EOGBS (infection during labour or uncomplicated birth).

This resulted in 23 clinical risk groups in scenario two, comprising the 13 groups from scenario one which would not be eligible for screening plus an additional 10 groups resulting from screening. Risks for EOGBS and related outcomes had been agreed in advance and were applied to these groups to obtain the numbers of affected individuals. The agreed risks are summarised in [tables 1 and 2](#).

[Figure 3](#) depicts the model structure for IAP and outcomes from the model.

To ensure a fair comparison, model outputs were checked to make sure that the number of EOGBS cases was equivalent in both scenarios if no antibiotic prophylaxis was given. In order to achieve this, the GBS




Figure 2 Flow of screening eligible women through the screening scenario. Group B *Streptococcus* (GBS), intrapartum antibiotic prophylaxis (IAP) against early onset GBS (EOGBS).

colonisation transition rates in the model needed to be adjusted to ensure that there was no overall change in GBS carriage rates between the time of screening and the time of delivery.

Consensus building approach

Members of the expert group are listed in table 3. They comprised representatives from UK organisations responsible for the development of guidance, policy and patient advocacy relating to EOGBS, as well as experts with experience in clinical practice fields related to EOGBS, microbiology, epidemiology or statistics.

Members of the group were given the opportunity to comment on the structure of the model and individual model parameters.

Evidence identified in the UK NSC's 2012 triennial review of antenatal GBS screening policy and national guidance documents were used to inform data used in the model.^{6 13 14 25} These were supplemented by a broad search in September 2014 for papers published since the 2012 NSC evidence review (see online supplementary information for details). Relevant national data was also identified from sources such as the Office for National Statistics.

Higher quality evidence, such as systematic reviews^{28 29} and randomised clinical trials, were prioritised as sources of data, as were studies from the UK.^{4 8} A summary of up to five potential values for each input parameter, based on the best available and most relevant evidence identified, was provided to the expert group. Where no evidence regarding an input parameter was identified, this was also noted.

Through a two part survey, the group was asked to consider the relevance of each input parameter, the most appropriate source of the parameter value, and the applicability of the source-study findings to a UK population. Group members could also suggest alternative sources

of data. In the absence of suitable published data for a parameter, the expert group selected a figure based on clinical experience and opinion.

All parameters and assumptions that received over 70% agreement were included in the model unchanged. Those that did not meet this threshold were refined and resubmitted to the expert group. Those that received over 50% agreement in the second round were included in the model. A small number of assumptions went forward to the second round. These focused on the details of the modelled IAP strategy. This included the timing and duration of IAP, whether second-line antibiotics should be included for analysis in the model, the schedule for those receiving IAP and subsequently developing fever in labour and whether IAP uptake and optimum duration should vary by risk group. All proposed changes received 70% or more, except for one which received 65%.

The survey results informed two meetings of the expert group. The first meeting reviewed and agreed about the model structure and input parameters. The second received and discussed the outputs of the model.

Patient and public involvement

Representatives of GBSS were involved as members of the expert group in all stages of the group's work. This included individual discussion with the project lead (DB) regarding the consensus building approach to agreeing to the model's parameters, priorities and outcomes. It also included participation in the two-part survey and workshops to discuss the model.

Results of the model were disseminated to all members of the expert group in the form of a draft report of the modelling project. Comments received from GBSS raised a number of concerns about the outcomes of the model. These concerns centred on the emerging results of an enhanced surveillance study which suggested that the rate of EOGBS had increased

Table 1 Population characteristics and early onset GBS (EOGBS) mortality and morbidity outcomes

Parameter	Input data	Data source
Maternity population characteristics		
Number of live births*	England and Wales – 718235 Scotland – 57202 Northern Ireland – 24890 Total – 800327	Office for National Statistics 2012 live birth data ³⁸ Information Services Division Scotland 2012–2013 data ³⁹ and National Records of Scotland 2012 data ⁴⁰ Northern Ireland Statistics and Research Agency 2012 ⁴¹
Elective caesarean section rate	England – 10.7% Wales – 11.7% Scotland – 12.8% Northern Ireland – 15.3% Total – 11.04%*	Hospital Episode Statistics (HES) 2012–2013 data ⁴² Stats Wales 2012–2013 data ⁴³ ISD Scotland 2012–2013 data ³⁹ Department of Health, Social Services and Public Safety Northern Ireland 2012–2013 data ⁴¹
Pre-term birth rate	8.2%	HES 2012–2013 data ⁴²
Risk factors indicating intrapartum antibiotics in UK guidance		
Proportion of the population with antenatal risk factors for EOGBS		
Previous infant with EOGBS	0.03% of model cohort	Colbourn <i>et al</i> , 2007 ⁹
Incidental group B <i>S treptococcus</i> detection	5.0% of model cohort	Expert group consensus agreement based on Colbourn <i>et al</i> , 2007 ⁹ and Daniels <i>et al</i> , 2011 ⁴⁴
Proportion of the term population with intrapartum risk factors		
Rate of maternal intrapartum infection in term deliveries	2.1%	Daniels <i>et al</i> , 2011 ⁴⁴
Proportion of the pre-term population with additional intrapartum risk factors		
Rate of maternal intrapartum infection in pre-term deliveries	4.1%	Daniels <i>et al</i> , 2011 ⁴⁴
Rate of pre-labour rupture of membranes (PROM) in pre-term deliveries	40%	Royal College of Obstetricians and Gynaecologists (RCOG) Pre-term PROM Green-top guideline 44, 2010 ⁴⁵
EOGBS mortality and morbidity outcomes		
Mortality in pre-term babies with EOGBS	18.3%	RCOG, 2012 ¹⁴ [derived from Heath <i>et al</i> , 2004 ⁴]
Mortality in term babies with EOGBS	5.7%	RCOG, 2012 ¹⁴ [derived from Heath <i>et al</i> , 2004 ⁴]
Morbidity in pre-term babies with EOGBS	6.91%	Expert group consensus agreement [based on Colbourn <i>et al</i> , 2007 ⁹]
Morbidity in term babies with EOGBS	6.89%	Expert group consensus agreement [based on Colbourn <i>et al</i> , 2007 ⁹]

*Excludes stillbirths, miscarriages and terminations; multiple births are only counted once.

and that mortality had decreased.⁶ GBSS suggested that the model should be re-run with the new data and that greater emphasis on EOGBS in term women was needed. It was not possible to address all the concerns without reconvening the expert group in a new modelling exercise. However, GBSS's concern prompted a post hoc analysis in term women with no risk factors indicating IAP. This is the most important group in the context of screening and the post hoc analysis is briefly reported later in this paper.

Outcomes

Outcomes for each scenario were evaluated in a hypothetical UK maternity cohort, over a 1-year period.

The outcomes were total culture positive EOGBS infections, EOGBS mortality and severe morbidity (eg, severe motor, intellectual, visual, hearing or other neurological impairment that meant the child was not able to attend mainstream school), use of IAP and maternal penicillin anaphylaxis. These outcomes were combined to explore the number of additional women needed to be treated with IAP to prevent additional EOGBS cases, deaths and severe disability.

The expert group noted that some of the model input parameters were based on low quality or inconclusive evidence. In order to explore the effect these data may have on outcomes, one-way sensitivity analyses were

Table 2 Estimated early onset GBS (EOGBS) incidence rates in each clinical risk group

Risk group	Input data	Data source
Background incidence/risk group		
No risk factors at term	0.2 per 1000 births	Royal College of Obstetricians and Gynaecologists (RCOG), 2012 ¹⁴
Suspected maternal infection at term	5.29 per 1000 births	RCOG, 2012 ¹⁴
Pre-term birth	2.30 per 1000 births	RCOG, 2012 ¹⁴
Suspected maternal infection at pre-term	5.29 per 1000 births	Expert group consensus agreement based on RCOG, 2012 ¹⁴
Pre-term pre-labour rupture of membranes	2.30 per 1000 births	Expert group consensus agreement based on RCOG, 2012 ¹⁴
Previous baby with EOGBS with no other risk factors	50 per 1000 births	Expert group consensus agreement
Incidence rate/1000 births		
Previous baby with EOGBS with one more additional risk factor	100 per 1000 births	Expert group consensus agreement
Incidental group B <i>Streptococcus</i> detection	2.30 per 1000 births	Expert group consensus agreement based on RCOG, 2012 ¹⁴
GBS carrier in labour with no other antenatal or intrapartum risk factors, delivering at term	0.91 per 1000 births	Expert group consensus agreement based on available data
GBS carrier in labour with suspected maternal infection, delivering at term	24.0 per 1000 births	Expert group consensus agreement based on available data
Not a GBS carrier in labour, delivering at term (with or without suspected maternal infection)	0 per 1000 births	Expert group consensus agreement
Background colonisation rate		
Colonisation rate	22% [Sensitivity analysis: 20%–30%]	Expert group consensus agreement

carried out to look at the impact of varying the following input parameters:

- ▶ Screening uptake rate.
- ▶ Antibiotic delivery in screen positive women.
- ▶ Effectiveness of IAP in preventing EOGBS.
- ▶ Transition rates for GBS status from screening to delivery.

The sensitivity analyses were run using a plausible lower and higher estimate, based on ranges agreed by the expert group. Individual parameters were changed one at a time, leaving all other parameters unchanged to provide discrete analyses of their impact. Input parameter values and data sources are presented in [tables 1, 2, 4 and 5](#).

Results

The model's key results are presented in [table 6](#).

Based on the inputs 800 327 live births were included in the model. Current guidance recommends that women with intact membranes undergoing elective caesarean should not receive GBS IAP in the absence of labour.¹⁴ As such, live births by elective caesarean were excluded, reducing the number included in the analysis to 711 999. Without IAP, there were an estimated 421 cases of culture positive EOGBS, a rate of 0.59/1000 live births. The modelled estimate of deaths and severe disability caused by EOGBS without IAP was 42 and 29 respectively.

In the risk based scenario, 30 666 women were estimated to receive antibiotics in labour and 70 cases of EOGBS prevented. In the screening scenario, a further 96 260 women received IAP on the basis of the screening result. This resulted in the prevention of an additional 52 to 57 cases of EOGBS (range in sensitivity analyses: 40 to 67) which included the prevention of three deaths (range in sensitivity analyses: two to four) and four cases of severe disability (range in sensitivity analyses: 3 to 5). This means that with screening, an additional 1675 to 1854 women receive IAP to prevent one EOGBS case and 24 065 to 32 087, to prevent one EOGBS death. Maternal anaphylaxis remained an extremely rare event in both scenarios, with 0.3 cases in the risk based scenario and 1.7 cases in the screening scenario.

Among women receiving IAP in the model, 8% received clindamycin due to reported penicillin allergy. It was assumed that treatment failure due to clindamycin resistance would be avoided by susceptibility testing in screen-positive women.

The sensitivity analyses did not have a large impact on results (see ranges above). An additional, post hoc analysis, focusing on term women with no risk factors indicating IAP, was undertaken. All parameters remained the same as those described above. However, an increase in




Figure 3 Model structure for women with an indication for intrapartum antibiotic prophylaxis (IAP) and outcomes of babies with early onset GBS (EOGBS).

the rate of EOGBS in carriers in this group along with a decrease in mortality was assumed. When the rate of EOGBS was double that used in the model, the additional number of women receiving IAP to prevent a case of EOGBS was 844 to 926. The reduced mortality rate meant that the number of women receiving IAP to prevent a death from EOGBS remained the same as the model's main estimate of 24 065 to 32 087.

DISCUSSION

The key outcomes of the model are summarised in [box 1](#).

This model suggests that the additional reduction in mortality and severe disability due to EOGBS with culture-based screening for maternal GBS carriage added to the current risk-based approach is likely to be, numerically, very limited in the UK. The addition of screening to the modelled risk-based prevention strategy increased the prevention of EOGBS cases from approximately 16% with the risk-based prevention strategy alone to approximately 25% of the modelled total. Similarly, the proportion of deaths prevented increased from approximately 12% to 19%.

There are a number of reasons for this limited numerical impact. The distribution of EOGBS across the risk groups and the clustering of its worst effects in groups outside the screening population limit the benefits that can be expected. In the model, 54% of the EOGBS cases

and 75% of deaths occurred in groups which were not eligible for screening, notably in the groups of women who already have known risk factors for EOGBS prior to 36 weeks or who give birth pre-term. In addition, not all women carrying GBS at labour would be correctly identified by the test. The model estimated that between 20 916 and 30 726 women who screened negative at 36 weeks gestation would be GBS positive at delivery. This group comprised women whose GBS carriage status was estimated to change from negative at the point of screening to positive at the time of labour. This number includes women truly transitioning in carriage status and also those receiving false negative screening test results.

Expert consensus and examples of non-UK based screening programmes suggest that attrition along the pathway should be expected and that uptake of both screening and IAP would be less than 100%. In addition, the delivery of IAP for sub-optimum durations is thought to reduce its prophylactic effect and was factored into the model. However, it should be noted that estimates of effectiveness are not well grounded in clinical trial evidence,²⁹ and the evidence base exploring duration of administration and prophylactic effect is limited to observational studies.^{30–32}

Prevention of EOGBS as a result of screening has to be considered in relation to the impact on the population as a whole. The likelihood of having a baby affected by

Table 3 Members of the expert group

Expert group member	Position
Professor Catherine Peckham (Chair)	National Health Service (NHS), NHS Infectious Diseases in Pregnancy Screening Programme, Executive Lead/Institute for Child Health
Dr Alison Bedford-Russell	Neonatologist, Birmingham Women's Hospital/Group B Strep Support (GBSS)
Professor Peter Brocklehurst	Director, Birmingham Clinical Trials Unit, University of Birmingham/Royal College of Obstetricians and Gynaecologists (RCOG) Greentop Guideline
Professor Androulla Efstratiou	Head, WHO Global Reference Centre for Diphtheria & Streptococcal Infections and European Centre for Disease Prevention and Control, UK Scientific Coordinator
Professor Paul Heath	Consultant in paediatric infectious diseases, St George's University of London & St Georges University Hospitals NHS Trust, London: BPSU study/RCOG Greentop Guideline
Dr Rhona Hughes	Obstetrician, Edinburgh/RCOG Greentop Guideline
Dr Theresa Lamagni	Senior Epidemiologist and Section Head, Healthcare-Associated Infection & Antimicrobial Resistance Division, National Infection Service, Public Health England (PHE)
Dr Anne Mackie	Director of Screening & Screening Quality Assurance, PHE
Mr John Marshall	Evidence Lead, UK National Screening Committee
Dr Rachel Moll	National Medical Director's Fellow, PHE
Mrs Jane Plumb	Chief Executive, GBSS
Dr Julie Robotham	Statistics, Modelling and Economics Department, PHE
Ms Farah Seedat	PhD student, University of Warwick
Dr Nan Shetty	Consultant Microbiologist and Training Lead, Reference Microbiology Services, PHE
Professor Helen Spiby	Professor of Midwifery, University of Nottingham
Professor Phillip Steer	Emeritus Professor of Obstetrics & Gynaecology, Imperial College/GBSS
Professor Ben Stenson	Neonatologist, Royal Infirmary of Edinburgh/RCOG Greentop Guideline
Professor Mark Turner	Neonatologist, Liverpool Women's NHS Foundation Trust

EOGBS appears to be low in women delivering at term with no known risk factors; a rate of about 0.2/1,000 live births was used in the model. These are the women who would be eligible for screening. Studies of antenatal GBS screening test accuracy mainly focus on accuracy for predicting maternal GBS carriage at delivery, and rarely report on neonatal outcomes or ability of the test to predict these.^{6,28} The positive predictive value of screening in late pregnancy for the outcome of EOGBS has recently been estimated as approximately 0.2% (2 cases of EOGBS per 1000 screen-detected carriers).⁶ This would be the level of risk reported to women who screen positive. Overdetection is a constant concern about screening, and its high rate in this context means that the ability of GBS screening to provide high quality post-test information may need to be questioned.

Overtreatment is a consequence of overdetection. A number of factors contribute to the high rate of overuse of IAP as a consequence of screening. The model estimated that between 16382 and 24065 screen positive women would receive antibiotics when they are no longer carrying the bacterium during labour. In addition, a large proportion of carriers in labour do not transmit the bacterium to the neonate during delivery.^{6,33} Among colonised neonates, only 3% develop EOGBS.³⁴

The absence of a diagnostic or risk refinement strategy, to follow a screen-positive result means that

many thousands of women would receive GBS IAP to manage a very low risk of EOGBS affecting their baby.

Reports of GBS organisms with reduced susceptibility, or resistance to penicillin, have caused concern despite being very rare and of uncertain clinical significance.^{5,11} While clindamycin has previously been recommended for IAP in women with penicillin allergy, the increase in resistance to this antibiotic has led to RCOG recommending that it should no longer be used for this purpose.⁷ In addition to this, the possibility has been raised that intrapartum antibiotics may have long-term effects on the infant gut flora and research into this is in the early stages.⁶

The difficulty in quantifying the harm of GBS IAP may make screening appear to be a harm-free intervention.⁶ However, the use of antibiotics in such a large group of women, the vast majority of whom will not experience benefit, means its fit with the current policy emphasis on prudent antibiotic prescribing goals¹⁸ may be difficult. This is because, as modelled, screening would do little to reduce antibiotic usage in the clinical risk groups who already receive IAP, there is low risk of EOGBS in the screened population, limited impact of screening on the worst outcomes, a lack of evidence to estimate whether outcomes are different for screen-detected and clinically-detected babies with EOGBS and

Table 4 Antibiotic uptake, optimal and suboptimal delivery, anaphylaxis

Parameter	Input data	Data source	
Uptake of antibiotics			
Maternal infection at term	80%	Expert group consensus agreement	
Maternal infection at pre-term	85%	Expert group consensus agreement	
Pre-term pre-labour rupture of membranes (pPROM)	5%	Expert group consensus agreement	
Screened group B <i>Streptococcus</i> (GBS) positive	80% [Sensitivity analysis 70%–90%]	Expert group consensus agreement [based on Albouy-Llaty <i>et al.</i> 2012 ⁴⁶]	
Previous baby with early onset GBS (EOGBS)	90%	Expert group consensus agreement	
Incidental detection of GBS carriage	47%	Moorthy <i>et al.</i> , 2014 ⁴⁷	
Optimal and suboptimal intrapartum antibiotic prophylaxis (IAP) delivery			
Parameter	Optimal IAP (delivered >2 hours)	Suboptimal IAP (delivered <2 hours)	Data source
Groups treated with antibiotics not solely for prevention of EOGBS			
Maternal infection at term	20%	80%	Expert group consensus agreement
Maternal infection at pre-term	10%	90%	Expert group consensus agreement
pPROM	65%	35%	Expert group consensus agreement
Groups receiving IAP solely for the prevention of EOGBS			
Screened GBS positive	65%	35%	Expert group consensus agreement [based on Lin <i>et al.</i> , 2011 ³⁰]
Previous baby with EOGBS	65%	35%	Expert group consensus agreement
Incidental detection of GBS carriage	65%	35%	Expert group consensus agreement
IAP effectiveness in preventing EOGBS			
IAP effectiveness (prevention of EOGBS)	RR [*] 0.17 (Sensitivity analysis 0.05 to 0.30) RR=relative risk	RR [*] 0.58 (Sensitivity analysis 0.525 to 0.65) Suboptimal effectiveness is 50% lower than optimal	Ohlsson <i>et al.</i> , 2014 ²⁹ (optimal) and expert group consensus agreement (sub-optimal)
Allergy, resistance and anaphylaxis rates			
Self-reported penicillin allergy rate	8%		Turrentine <i>et al.</i> , 2009 ⁴⁸ and Zilberman <i>et al.</i> , 2014 ⁴⁹
Clindamycin resistance rate	17%		Public Health England, 2014 (2013 data) ⁵⁰
Maternal penicillin anaphylaxis	1.56 per 100 000 women treated		Mulla <i>et al.</i> , 2010 ⁵¹

a high rate of overtreatment. In addition, the large number of additional women receiving GBS IAP will need to do so in a setting where the antibiotics can be

delivered intravenously. This may reduce the choice of birth setting for these women by removing the option of home birth.

Table 5 Screening test uptake and colonisation status transitions between screening and delivery

Screening uptake		
Parameter	Input data	Data source
Screening uptake rate	90% [Sensitivity analysis – 75%–95%]	Expert group consensus agreement
Colonisation status transition		
'Transition rate' from group B <i>Streptococcus</i> (GBS+) at 36 weeks to GBS- at delivery*	Base case 1: 25% Base case 2: 17% [Sensitivity analysis 11.7%–40%]	Valkenburg-van den Berg <i>et al.</i> 2010, ²⁸ Di Renzo <i>et al</i> 2015 ⁵² and expert group consensus agreement
'Transition rate' from GBS- at 36 weeks to GBS+ at delivery*	Base case 1: 7.1% Base case 2: 4.8% [Sensitivity analysis 3.3%–12%]	Valkenburg-van den Berg <i>et al.</i> 2010, ²⁸ Di Renzo <i>et al</i> 2015 ⁵² and expert group consensus agreement

*NB: Based on the data available, it is not possible to distinguish between women who have an incorrect screening result (ie, false positive or false negative) and whose true colonisation status remains unchanged at delivery, and those who had a correct screening result (true positive or true negative) and then transition to a different colonisation status at delivery.

Strengths and limitations

Antenatal screening for GBS is a controversial topic, and one of the strengths of this study was the use of a

multidisciplinary stakeholder group encompassing a wide range of viewpoints, to gain consensus on the model structure and inputs. To our knowledge, this is the first

Table 6 Model results

Risk group	Outcomes before application of either prevention scenario (baseline)		Scenario 1 (risk-based prevention)		Scenario 2 (risk-based plus screening-based prevention)	
	Number of women in risk group	Early onset GBS (EOGBS) cases	Antibiotic type offered	Number of women receiving antibiotics	Antibiotic type offered	Number of women receiving antibiotics
Neonate affected by EOGBS in a previous pregnancy	214	12	IAP and broad spectrum*	192	IAP and broad spectrum*	192
Incidental detection of maternal GBS carriage	35 600	84	IAP and broad spectrum*	17 005	IAP and broad spectrum*	17 005
Pre-term delivery without the above risk factors	55 446	134	Broad spectrum*	3041	Broad spectrum*	3041
Suspected intrapartum infection at term without the above risk factors	13 036	69	Broad spectrum*	10 428	Broad spectrum* in all, preceded by IAP in 2065 screen-positive women	10 428
Term women with no antenatal or intrapartum risk factors indicating intrapartum antibiotic prophylaxis (IAP) (without screening)	607 703	122	N/A	0	IAP for women who screen group B <i>Streptococcus</i> GBS positive	96 260
Summary	Baseline		Scenario 1		Scenario 2	
Totals for each scenario	Population	800 327 women	Treated with antibiotics	30 666	Treated with antibiotics	126 926
	EOGBS cases	421	EOGBS cases prevented	70	EOGBS cases prevented	122–127
	Deaths from EOGBS	42	Deaths from EOGBS prevented	5	Deaths from EOGBS prevented	8
	Severe disability from EOGBS	29	Severe disability from EOGBS prevented	5	Severe disability from EOGBS prevented	8–9

*Broad spectrum antibiotics given to those with suspected intrapartum infection.

N/A, not applicable.

Box 1 Key outcomes

Summary information relating to the screening plus risk-factor scenario

- ▶ 558 664 women were screened at 36 weeks out of a total of 620 739 women. 122 906 were screen-positive and 435 758 screen-negative. 62 075 remained unscreened due to declining the offer or failure to be offered screening.
- ▶ The addition of screening at 36 weeks resulted in 96 260 women receiving intrapartum antibiotic prophylaxis (IAP) in addition to those already treated within the risk-based strategy alone. Of the screened women, 16 382 to 24 065 received IAP when they no longer carried group B *Streptococcus* (GBS) at delivery.
- ▶ Screening at 36 weeks prevented 52 to 57 cases of early onset GBS (EOGBS), three deaths and four severe morbidities due to EOGBS, in addition to those prevented in the risk-based scenario.
- ▶ In the screening-plus-risk -factor scenario, 1675 to 1854 women received penicillin IAP to prevent a case and 24 065 to 32 087 to prevent a death due to EOGBS, in addition to those prevented in the risk-based scenario.
- ▶ 2065 screen-positive term women who subsequently developed a suspected infection in labour received GBS IAP in addition to broad spectrum antibiotics. Apart from this group, the number of women receiving antibiotics and the type of antibiotic used remained constant in both strategies in the women with risk factors.

time this approach has been taken to inform a model of antenatal GBS screening.

Data sources were identified through the use of UK National Screening Committee evidence reviews and broad systematic searches for subsequently published evidence. Higher quality evidence, such as published systematic reviews were used where available, as well as national sources including guidance and national statistics. However, systematic reviews to support each model input were not feasible. In addition, the expert group provided input on realistic input values in the absence of relevant evidence.

Limited evidence was available for many model parameters. This includes the rate of penicillin anaphylaxis and the proportion of pregnant women reporting allergy to penicillin. Research to inform these and other inputs would improve future modelling exercises. Research into factors which impact EOGBS risk in babies born to women colonised by GBS could be valuable in the development of a post-screening risk refinement strategy to narrow the pool of women receiving antibiotics.

There is uncertainty regarding the model's outputs due to limitations in the underlying evidence. The results are not dissimilar to those estimated by a hospital in Australia which introduced universal screening. Planned and post hoc sensitivity analyses did not substantially affect this. This Australian study estimated that 1190 women received IAP to prevent one case of EOGBS.³⁵ However, it was not clear whether their figure took into account the women who would have received IAP based on the risk-based approach prior to the introduction of screening. In addition, the study was observational and the number of

births during the screening period relatively small; so the estimate of EOGBS cases prevented may not have been very robust.

Since the completion of the modelling exercise, there have been changes to the UK's recommended risk-based strategy, mainly that the RCOG has now recommend offering IAP to all women in established pre-term labour. No analysis was undertaken to estimate the effect of these changes. However, this change should not influence the screening scenario, as women who give birth pre-term would not have had the chance to be screened.

The model focused on a limited set of outcomes. In part, this related to the overall aims of the project, which was to try to establish a set of shared assumptions about a core set of basic parameters and outcomes. However, it was also related to the limitations in the data available; for example, it was not considered possible within the terms of this project to model outcomes relating to mild or moderate disability arising from EOGBS.

The model did not explore qualitative questions, for example whether screening would affect maternal choice, the medicalisation of labour or women's experience of maternity services.

CONCLUSION

This is the first time a consensus-based assessment of the evidence has been developed within a UK-based expert group. A very limited evidence base was encountered and important issues such as the impact of screening on culture-negative disease or on women's birth experience could not be addressed. However, consensus was achieved on the structure of the model and the parameters required for an estimate of the preventive potential of screening.

The resulting outputs focus attention on a hypothetical screening programme which was estimated to combine a low impact, in terms of preventing the worst aspects of EOGBS, with a high impact, in terms of the volume of women categorised as high risk and treated with prophylactic antibiotics in labour.

Expectations about the benefits of interventions, including screening interventions, can be overestimated by both patients and health professionals.^{36 37} The modelling work steered by this expert group provides a useful sense of perspective on antenatal screening for maternal GBS carriage.

Acknowledgements We would like to thank all members of the expert group for contributing to this project. We would like to thank representatives of GBSS for their contribution to this project. We would also like to thank Anelia Boshnakova, Senior Information Specialist, Bazian Ltd. who performed the literature searches to identify evidence to inform development of the model.

Contributors The paper is derived from the 'UK National Screening Committee end of project report: Early onset streptococcal (EOGBS) disease: a report of a modelling exercise prepared for the expert group'. The modelling work was steered by an expert group with representation from the relevant professional groups and the major advisory and standard setting bodies and a patient advocacy group. CP chaired the expert group. The expert group commented on the end of project report, but the preparation of this manuscript was the responsibility of the authors

alone, and has not been reviewed by the expert group, and may not represent those of all expert group members. DB and JM lead the work to define the model's aims, organise the expert group and to manage the process. AW was responsible for producing the model and provided advice throughout the project's life cycle. AW and AB are employees of Bazian Ltd. which received payment to develop the model through contracts with the UK National Screening Committee. All authors contributed to interpretation of the results. JM, DB, and AW drafted the end of project report and this publication, with critical review by all authors. JM acts as guarantor for the paper.

Funding UK National Screening Committee.

Competing interests CP chaired the expert group. JM and DB were both employees of the UK National Screening Committee at the time of preparation of the model. AW is an employee of Bazian Ltd. who received payment to develop the model and write up the project through contracts with the UK National Screening Committee. Bazian Ltd. have also been paid to carry out other literature searches, reviews and models for the National Screening Committee, including reviews of the evidence on antenatal GBS screening. Bazian Ltd. is part of The Economist Group, and holds contracts with public and private organisations within the healthcare industry.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Baker CJ, Barrett FF. Transmission of group B streptococci among parturient women and their neonates. *J Pediatr* 1973;83:919–25.
- Vergnano S, Menson E, Kennea N, et al. Neonatal infections in England: the NeonIN surveillance network. *Arch Dis Child Fetal Neonatal Ed* 2011;96:F9–14.
- Okike IO, Johnson AP, Henderson KL, et al. Incidence, etiology, and outcome of bacterial meningitis in infants aged <90 days in the United Kingdom and Republic of Ireland: prospective, enhanced, national population-based surveillance. *Clin Infect Dis* 2014;59:e150–e7.
- Heath PT, Balfour G, Weisner AM, et al. Group B streptococcal disease in UK and Irish infants younger than 90 days. *Lancet* 2004;363:292–4.
- Lamagni TL, Keshishian C, Efstratiou A, et al. Emerging trends in the epidemiology of invasive group B streptococcal disease in England and Wales, 1991–2010. *Clin Infect Dis* 2013;57:682–8.
- UK National Screening Committee. *The UK NSC recommendation on Group B Streptococcus screening in pregnancy review of screening*. London: Public Health England, 2016.
- Royal College of Obstetricians and Gynaecologists. Prevention of early-onset neonatal group b streptococcal disease: green-top guideline No. 36. *BJOG* 2017;124:e280–305.
- Vergnano S, Embleton N, Collinson A, et al. Missed opportunities for preventing group B streptococcus infection. *Arch Dis Child Fetal Neonatal Ed* 2010;95:F72–3.
- Colbourn T, Asseburg C, Bojke L, et al. Prenatal screening and treatment strategies to prevent group B streptococcal and other bacterial infections in early infancy: cost-effectiveness and expected value of information analyses. *Health Technol Assess* 2007;11:1–226.
- Hamada S, Vearncombe M, McGeer A, et al. Neonatal group B streptococcal disease: incidence, presentation, and mortality. *J Matern Fetal Neonatal Med* 2008;21:53–7.
- Verani JR, McGee L, Schrag SJ. Prevention of perinatal group B streptococcal disease—revised guidelines from CDC, 2010. *MMWR Recomm Rep* 2010;59:1–36.
- SMI B 58. *Detection of carriage of group B streptococci (Public Health England, 2015*. England: SMI B 58.
- National Institute for Health and Care Excellence. *Neonatal infection (early onset): antibiotics for prevention and treatment*. CG149. London: National Institute for Health and Care Excellence, 2012.
- Royal College of Obstetricians and Gynaecologists. *The prevention of early-onset neonatal Group B streptococcal disease: Green-top guideline No. 36*. 2nd edition. London: Royal College of Obstetricians and Gynaecologists, 2012.
- UK National Screening Committee. *UK NSC: evidence review process*. London: Public Health England.
- National Maternity Review. *Better births: improving outcomes of maternity services in England*. London: NHS England, 2016.
- Department of Health. *NHS outcomes framework 2015 to 2016*. London: Department of Health, 2014.
- Department of Health. *UK 5 Year antimicrobial resistance strategy 2013 to 2018*. London: Department of Health, 2013.
- Brocklehurst P. Screening for Group B streptococcus should be routine in pregnancy: AGAINST: current evidence does not support the introduction of microbiological screening for identifying carriers of Group B streptococcus. *BJOG* 2015;122:368.
- Steer PJ. FOR: the case for screening. *BJOG* 2015;122:369.
- Plumb J, Clayton G. Group B streptococcus infection: risk and prevention. *Pract Midwife* 2013;16:27–30.
- Stewart M, Scamell M, McFarlane A. Professionals respond to GBS article. *Pract Midwife* 2013;16:8–9.
- Petition calling for routine group. Petition calling for routine group B Strep tests in the UK goes to parliament. <https://www.midirs.org/group-b-step-tests-uk-parliament/>
- Phares CR, Lynfield R, Farley MM, et al. Epidemiology of invasive group B streptococcal disease in the United States, 1999–2005. *JAMA* 2008;299:2056–65.
- Ltd B. *Screening for Group B streptococcal infection in pregnancy: external review against programme appraisal criteria for the UK national screening committee*. London: UK National Screening Committee, 2012.
- Carbonell-Estrany X, Figueras-Aloy J, Salcedo-Abizanda S, et al. Probable early-onset group B streptococcal neonatal sepsis: a serious clinical condition related to intrauterine infection. *Arch Dis Child Fetal Neonatal Ed* 2008;93:F85–9.
- McQuaid F, Plumb J. More needs to be done to prevent Group B strep infection in the UK. *Br J Midwifery* 2015;23:418–23.
- Valkenburg-van den Berg AW, Houtman-Roelofs RL, Oostvogel PM, et al. Timing of group B streptococcus screening in pregnancy: a systematic review. *Gynecol Obstet Invest* 2010;69:174–83.
- Ohlsson A, Shah VS. Intrapartum antibiotics for known maternal Group B streptococcal colonization. *Cochrane Database Syst Rev* 2014;6:Cd007467.
- Lin FY, Weisman LE, Azimi P, et al. Assessment of intrapartum antibiotic prophylaxis for the prevention of early-onset group B Streptococcal disease. *Pediatr Infect Dis J* 2011;30:759–63.
- Fairlie T, Zell ER, Schrag S. Effectiveness of intrapartum antibiotic prophylaxis for prevention of early-onset group B streptococcal disease. *Obstet Gynecol* 2013;121:570–7.
- Illuzzi JL, Bracken MB. Duration of intrapartum prophylaxis for neonatal group B streptococcal disease: a systematic review. *Obstet Gynecol* 2006;108:1254–65.
- Daniels J, Gray J, Pattison H, et al. Rapid testing for group B streptococcus during labour: a test accuracy study with evaluation of acceptability and cost-effectiveness. *Health Technol Assess* 2009;13:1–154.
- Colbourn T, Gilbert R. An overview of the natural history of early onset group B streptococcal disease in the UK. *Early Hum Dev* 2007;83:149–56.
- Angstetra D, Ferguson J, Giles WB. Institution of universal screening for Group B streptococcus (GBS) from a risk management protocol results in reduction of early-onset GBS disease in a tertiary obstetric unit. *Aust N Z J Obstet Gynaecol* 2007;47:378–82.
- Hoffmann TC, Del Mar C. Clinicians' expectations of the benefits and harms of treatments, screening, and tests: a systematic review. *JAMA Intern Med* 2017;177:407–19.
- Hoffmann TC, Del Mar C. Patients' expectations of the benefits and harms of treatments, screening, and tests: a systematic review. *JAMA Intern Med* 2015;175:274–86.
- Office for National Statistics. *Births in England and Wales, 2012* <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/livebirths/bulletins/birthsummarytablesenglandandwales/2013-07-10>
- Information Services Division Scotland. *Births in Scottish hospitals, 2013* <http://www.isdscotland.org/Health-Topics/Maternity-and-Births/Publications/data-tables.asp?id=1285#1285>
- National Records of Scotland. *Births, deaths and other vital events - preliminary annual figures, 2012* <https://www.nrscotland.gov.uk/statistics-and-data/statistics/statistics-by-theme/vital-events/general-publications/births-deaths-and-other-vital-events-preliminary-annual-figures/2012>
- Northern Ireland Statistics and Research Agency. *Births in Northern Ireland*. Belfast: Northern Ireland Statistics and Research Agency,

2012. http://webarchive.nationalarchives.gov.uk/20160512114115/http://www.nisra.gov.uk/archive/demography/publications/births_deaths/births_2012.pdf
42. Health and Social Care Information Centre. Hospital episode statistics: NHS maternity statistics - England, 2012 – 13. <http://digital.nhs.uk/catalogue/PUB12744>
 43. Wales S. Number of deliveries in Wales by delivery method and length of stay: 2012-2013. 2015 <https://statswales.gov.wales/Catalogue/Health-and-Social-Care/NHS-Primary-and-Community-Activity/Maternity/NumberOfDeliveriesInWales-by-DeliveryMethod-LengthOfStay>
 44. Daniels JP, Gray J, Pattison HM, *et al*. Intrapartum tests for group B streptococcus: accuracy and acceptability of screening. *BJOG: An International Journal of Obstetrics and Gynaecology* 2011;118:257–65.
 45. Royal College of Obstetricians and Gynaecologists. *Preterm prelabour rupture of membranes: green-top guideline No. 44*. London: Royal College of Obstetricians and Gynaecologists, 2010.
 46. Albouy-Llaty M, Nadeau C, Descombes E, *et al*. Improving perinatal Group B streptococcus screening with process indicators. *J Eval Clin Pract* 2012;18:727–33.
 47. Moorthy V, Bakour S, Lawson K. PLD.40 Effective identification and management of Group B streptococcus in pregnancy and labour. *Arch Dis Child Fetal Neonatal Ed* 2014;99:A117.3–21.
 48. Turrentine MA, Ramirez MM, Mastrobattista JM. Cost-effectiveness of universal prophylaxis in pregnancy with prior group B streptococci colonization. *Infect Dis Obstet Gynecol* 2009;2009:1–11.
 49. Zilberman D, Williams SF, Kurian R, *et al*. Does genital tract GBS colonization affect the latency period in patients with preterm premature rupture of membranes not in labor prior to 34 weeks? *J Matern Fetal Neonatal Med* 2014;27:338–41.
 50. Public Health England. *Voluntary surveillance of pyogenic and non-pyogenic streptococcal bacteraemia in England, Wales and Northern Ireland: 2013*. London: Public Health England, 2014.
 51. Mulla ZD, Ebrahim MS, Gonzalez JL. Anaphylaxis in the obstetric patient: analysis of a statewide hospital discharge database. *Ann Allergy Asthma Immunol* 2010;104:55–9.
 52. Di Renzo GC, Melin P, Berardi A, *et al*. Intrapartum GBS screening and antibiotic prophylaxis: a European consensus conference. *J Matern Fetal Neonatal Med* 2015;28:766–82.