

# Poultry\_Science\_Journal\_\_Volu me\_6\_Issue\_1\_Pages\_51-62- 2017.pdf

*by* T Yudiarti

---

**Submission date:** 15-Aug-2018 07:02PM (UTC+0700)

**Submission ID:** 990136297

**File name:** Poultry\_Science\_Journal\_\_Volume\_6\_Issue\_1\_Pages\_51-62-2017.pdf (561.78K)

**Word count:** 7594

**Character count:** 38867



## The Physiological Responses to Dietary Administration of Zinc Bacitracin and *Bacillus* Mixture on Low-Weight Day-Old Chicks

Sugiharto S, Yudiarti T, Isroli I & Widiastuti E

39

Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Central Java, Indonesia

Poultry Science Journal 2018, 6(1): 51-62

### Keywords

Probiotic  
In-feed antibiotic  
Multi-strains *Bacillus*  
Low weight day-old chick

### Corresponding author

Sugiharto Sugiharto  
[sgh\\_undip@yahoo.co.id](mailto:sgh_undip@yahoo.co.id)

### Article history

Received: November 16, 2016  
Revised: December 21, 2017  
Accepted: January 20, 2018

### Abstract

This study investigated the responses of low-weight day-old chicks to zinc bacitracin and *Bacillus* mixture on growth performance, hematology, intestinal selected microbiota population and carcass characteristics. A total of 192 unsexed Lohman MB-202 day-old broiler chicks were randomly allotted to four dietary treatment groups of 48 chicks each (6 replicates of 8 chicks) and fed for 35 days. The four treatments were CONT (basal diet without supplementation), AGP (basal diet with 0.04% zinc bacitracin), PROB (basal diet supplemented with 0.5% *Bacillus* mixture), and PROB+AGP (basal diet supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin). There were no significant differences in final body weight, feed intake, and feed conversion ratio of broilers. The relative weight of thymus was higher ( $P = 0.01$ ) in AGP than that in PROB and PROB+AGP birds. Birds in PROB+AGP had lower ( $P = 0.04$ ) values of alanine aminotransferase (ALT) enzyme compared to those in CONT and AGP groups, but the difference was not significant when compared with birds in PROB group. Serum creatinine was lower ( $P < 0.01$ ) in PROB+AGP than in CONT and AGP birds. Birds in AGP group tended ( $P = 0.09$ ) to have higher globulin concentration in the serum compared to other birds. There was also a tendency ( $P = 0.07$ ) for PROB+AGP birds to have higher albumin to globulin ratio compared with other birds. There was more lactic acid bacteria in ileal digesta ( $P = 0.05$ ) in PROB+AGP than in AGP birds, but the difference was not significant when compared to CONT and PROB birds. Birds in AGP group had higher ( $P = 0.05$ ) relative breast weight compared to other birds. In conclusion, *Bacillus* mixture did not affect performance and hematological parameters, but increased the intestinal population of lactic acid bacteria in broiler chicks.

### Introduction

It is well documented that the weight of one-day-old broiler chicks determines their future weight at market age (Tona *et al.*, 2004; Mendes *et al.*, 2011; Toghyani *et al.*, 2011). For example, chicks with low initial weights (34.40–35.22 g) had lower final body weights (BW) than those

with heavy hatching weights (39.29–41.30 g) (Mendes *et al.*, 2011). In another study, Michalczuk *et al.* (2011) showed that hatching chicks with BW higher than 42 g presented higher BW at day 36 compared to those with BW lower than or equal to 39 g at hatch. Similar findings have been observed in Pekin duck

13

Please cite this article as: Sugiharto S, Yudiarti T, Isroli I & Widiastuti E. 2018. The Physiological Responses to Dietary Administration of Zinc Bacitracin and *Bacillus* Mixture on Low-Weight Day-Old Chicks. *Poult. Sci. J.* 6(1): 51-62.

© 2018 PSJ. All Rights Reserved

(Sözcü and Ipek, 2017). Birds with a heavier initial BW may have more developed digestive organs (such as gizzard and intestine), which may improve digestive capacity and thus growth performance of the birds (Sözcü and Ipek, 2017). In these respects, it is discouraged to use chicks with low initial BW in commercial broiler farms as their rearing will reduce economical benefits. Chicks with low initial BW are usually produced from the junior breeder flock (25–30-week-old) (Tona *et al.*, 2004; Ulmer-Franco *et al.*, 2010).

Due to the regular egg production cycle, broiler breeders would produce chicks from junior breeder flocks at specific times. Instead of culling chicks with low hatching weights, improving the management and rearing conditions can be implemented (Butcher and Nilipour, 2002; Ulmer-Franco *et al.*, 2010). Another concern related to chicks from young breeder flock is a high mortality rate (Wyatt *et al.*, 1985). The reasons behind this are unclear but may be associated with poor immune function. In order to be as productive as the heavier chicks, these chicks need distinct management and rearing conditions (Butcher and Nilipour, 2002; Ulmer-Franco *et al.*, 2010). For example, providing extra vitamins and minerals to these chicks during the rearing period may improve their performance (Butcher and Nilipour, 2002). Some vitamins and minerals are essential for optimizing metabolism, and thus growth performance (Islam *et al.*, 2004; Ajuwon *et al.*, 2011) and immune functions in broiler chickens (Sanda, 2015; Sugiharto, 2016).

In-feed antibiotics have long been used as growth promoters, antimicrobial agents, and immune stimulants in broiler production (Lee *et al.*, 2012; Sugiharto, 2016). In the case of low day-old weight chicks, in-feed antibiotics are essential for improving the growth and health of broilers. However, long-term application of such antibiotics may result in antibiotic residues in broiler meat which can pose risk to human health (Sugiharto, 2016). Hence, alternatives to in-feed antibiotics are important, especially for day-old low-weight chicks. Among the in-feed antibiotic substitutes, probiotics – which contains a wide spectrum of *Bacillus* strains – has attracted considerable interest from poultry nutritionists. Compared to *Lactobacillus*-based probiotic cultures, which is commonly used as a feed additive in poultry nutrition, probiotics

containing a wide spectrum of *Bacillus* strains show better viability and stability, especially during feed processing and storage (Simon, 2005). The ability to form spores allows *Bacillus*-based probiotics to survive not only in the extreme conditions of the gastrointestinal tract, but to also be more stable during processing and feed storage (Elshagabee *et al.*, 2017). Therefore, probiotics that contain a wide spectrum of *Bacillus* strains seem to be more effective and practical in broiler production. In our previous study, we prepared a select mixture of *Bacillus* probiotics with vitamins and minerals (which we termed called “*Bacillus* mixture”), and found an improvement in the organ development and immune competence of broiler chicks (Isroli *et al.*, 2017).

The objective of the present study was to investigate the response of low day-old weight chicks to the administration of zinc bacitracin and *Bacillus* mixture in terms of growth performance, blood parameters, intestinal microbiota populations, and carcass characteristics. In this study, the effects associated with *Bacillus* mixture was compared with zinc bacitracin without vitamins and minerals supplementation in low-weight chicks. To date, zinc bacitracin is still commonly used in Indonesian commercial poultry producers.

#### Materials and Methods

A total of 192 one-day-old chicks (unsexed Lohmann meat broilers; BW =  $35.20 \pm 0.25$  g; mean  $\pm$  standard deviation) were used in the present trial. Upon arrival, the chicks were weighed individually and randomly allotted to one of four groups of 48 chicks each (6 replicates of 8 chicks). The chicks were raised in an open-sided broiler house with wire floor pens for the entire trial period. Light was provided 24 h per day. To control the temperature and humidity, the broiler house was equipped with a plastic curtain, light bulb as heater, and blower fan. The dietary treatment groups were CONT (basal diet without any supplementation), AGP (basal diet with 0.04% zinc bacitracin), PROB (basal diet supplemented with 0.5% *Bacillus* mixture), and PROB+AGP (basal diet supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin).

The study lasted for 195 days, and for the entire study period, the feed (in mash form) and water were provided *ad libitum*. The coccidiostat, enzymes, and other feed additives/supplements were not included in the basal diet. The basal

diet was formulated (Table, 1) to meet or exceed the Indonesian National Standards for Broiler Feed (SNI, 2006). To test the efficacy of the supplements in preventing the overgrowth of *Clostridium perfringens*, the diet was formulated to increase the *C. perfringens* proliferation in the intestine of birds by including bone meal, chicken feather meal, meat bone meal, and viscous cereals (M'Sadeq et al., 2015). The *Bacillus* mixture contained 12.10 log cfu/g multi-strains *Bacillus* (i.e., *Bacillus cereus* strain SIIA\_Pb\_E3, *Bacillus licheniformis* strain FJAT-29133, *Bacillus megaterium* strain F4-2-27 and *Bacilli* 64 pp. 11CM31Y12), 0.100 mg vitamin A, 0.018 mg vitamin D<sub>3</sub>, 0.100 mg vitamin E, 1200 mg Ca, 750 mg P, 0.08 mg Mg, 0.006 mg Co,

0.045 mg Cu, 0.015 mg Se, 0.180 mg S, 0.010 mg Zn, 0.060 mg KCl, 0.030 mg I, 0.060 mg Fe and 0.100 mg Mn. The *Bacillus* mixture was prepared in our laboratory, and zeolite was used as a carrier for the bacteria. The novel *Bacillus* strains have recently been isolated from the rumen content of cow, and exhibit antibacterial activity (*in vitro*) against *Escherichia coli* ATCC 25922 (unpublished data). These bacterial cultures also showed probiotic activities in broiler chicks during the brooding period *in vivo* (Isroli et al., 2017). The experiment was performed under the standard procedures of rearing and treating of farm animals stated in the law of the Republic of Indonesia number 18, 2009 regarding animal husbandry and health.

**Table 1.** Ingredients and chemical composition of basal diet

Items (% , unless otherwise noted)	Composition
Maize (CP 8.5%)	45.5
Soybean meal (CP 46%)	17.0
Wheat flour	10.0
Bread flour	5.00
Rice bran	4.45
Crude palm oil	3.50
Corn gluten meal	3.60
Distiller dried grains <sup>1</sup>	3.00
Meat bone meal	2.80
Hydrolyzed chicken feather meal	2.00
Bone meal	1.50
Lysine	0.55
Methionine	0.37
L-threonine	0.08
Salt	0.15
Premix <sup>2</sup>	0.50
<i>Analyzed composition</i>	
Metabolizable energy (Kcal/kg) <sup>3</sup>	3290
Crude protein	21.7
Crude fat	5.90
Crude fiber	6.79
Ash	10.9

<sup>1</sup>The dried residue remaining after the starch extraction from maize.

<sup>2</sup>Mineral-vitamin premix contained (62 g of diet): Ca 2.250 g, P 0.625 g, Fe 3.570 mg, Cu 0.640 mg, Mn 5.285 mg, Zn 0.003 mg, Co 0.001 mg, Se 0.013 mg, 10.016 mg, vitamin A 375 IU, vitamin D 150 IU, vitamin E 0.080 mg.

<sup>3</sup>Value was calculated according to formula (Bolton, 1967) as follow: 40.81 [0.87 [crude protein + 2.25 crude fat + nitrogen free extract] + 2.5].

Throughout the study period, birds were only vaccinated with commercial Newcastle disease virus (NDV) vaccine through eye drops and drinking water at day 4 and 18 of the experiment, respectively. Live BW and accumulative feed intake (FI) of broiler chicks were recorded at the end of experiment. Blood was collected from the bird's wing veins (6 birds per treatment) and placed in vacutainers

containing ethylene diamine tetra acetic acid (EDTA) for the determination of complete blood counts. For the serum biochemical analysis, the rest of the blood was collected in the vacutainers without anticoagulant, permitted to clot at room temperature, and centrifuged at 448 g for 15 min. The serum was frozen until analyses (Sugiharto et al., 2017). After blood sampling, the birds were slaughtered, de-feathered, and eviscerated. The

internal organs were immediately obtained and weighed (Sugiharto *et al.*, 2017). For the microbiological and short chain fatty acids (SCFAs) analysis, digesta were expelled by gently squeezing from the ileum (from Meckel's diverticulum to a point 4 cm proximal to the ileocecal junction) and cecum into the sterile sample bottles. The samples of breast muscle were collected for the determination of crude protein and fat content.

Complete blood counts were determined using a hematology analyzer (Prima Fully-auto Hematology Analyzer, PT. Prima Alkesindo Nusantara, Jakarta, Indonesia). The serum NDV antibody titers were measured according to hem agglutination inhibition (HI) assay (Villegas, 1987). The titers were presented as geometric mean titers ( $\text{Log}_2$ ). To determine the levels of total triglyceride, total cholesterol, high-density lipoprotein-cholesterol (HDL-c), and low-density lipoprotein-cholesterol (LDL-c) cholesterol, uric acid, and creatinine in serum, we used enzymatic colorimetric/color methods. The spectrophotometric/photometric tests were used to determine total protein, albumin, alanine aminotransferase (ALT), and aspartate aminotransferase (AST) in the serum of broilers. Globulin was calculated from the difference between total protein and albumin in serum. These biochemical analyses in serum were conducted using kit DiaSys Diagnostic System GmbH, Holzheim, Germany) according to the manufacturer's instructions. The bacterial populations in the ileal and cecal digesta of birds were determined (6 birds per treatment) according to Sugiharto *et al.* (2017) with few modifications. Coliform bacteria were enumerated on MacConkey agar (Merck KGaA,

Darmstadt, Germany) after aerobic incubation at 30°C for 24 h as red colonies. The number of lactic acid bacteria (LAB) was determined on de Man, Rogosa and Sharpe (MRS; Merck KGaA, Darmstadt, Germany) agar following an aerobic incubation at 38°C for 48 h. The number of *C. perfringens* was determined on tryptose sulfite cycloserine (TSC; Merck KGaA, Darmstadt, Germany) agar plates following anaerobic incubation at 38°C for 48 h. Analysis of SCFAs concentrations in the cecal digesta was conducted by gas chromatography according to the conditions described by Sugiharto *et al.* (2015). Proximate analysis was conducted to analyze the crude protein and fat contents of meat according to the standard methods (AOAC, 1995). Crude protein content was determined according to the Kjeldahl method, while crude fat was determined by the Soxhlet extraction method using petroleum ether.

The data were analyzed according to a completely randomized design by ANOVA using the General Linear Models Procedure in SAS (1985). The pen was considered as the experimental unit. The differences ( $P < 0.05$ ) among groups were further analyzed using Duncan's multiple-range test.

## Results

### Performance of broiler chicks

Data on the performance of broiler chicks are presented in Table 2. There were no significant differences in final BW, and feed conversion ratio (FCR) of broiler chicks between the treatment groups. However, birds in AGP and PROB groups tended ( $P = 0.06$ ) to have higher feed intake than birds in CONT and PROB+AGP groups.

**Table 2.** Effects of zinc bacitracin and *Bacillus* mixture on performance of broiler chicks at 35 days of age

Items	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P- value
	CONT	AGP	PROB	PROB+AGP		
Live BW (g)	909	972	955	917	40.8	0.66
Accumulative FI (g)	1,899	2,057	2,057	1,957	46.8	0.06
FCR (g/g)	2.19	2.22	2.22	2.24	0.04	0.98

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diet supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diets supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error.

### Internal organ weights of broiler chicks

The relative weights of internal organs of broiler chicks are shown in Table 3. The relative weight of thymus was higher ( $P = 0.01$ ) in AGP than in

PROB and PROB+AGP birds. All other internal organs assessed had similar relative weights across the treatment groups.

**Table 3.** Effects of zinc bacitracin and *Bacillus* mixture on relative weights of internal organs in broiler chicks

Items (% live BW)	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P-value
	CONT	AGP	PROB	PROB+AGP		
Heart	0.83	0.85	0.91	0.81	0.07	0.77
Liver	3.17	3.14	3.63	3.26	0.29	0.62
Proventriculus	0.43	0.47	0.46	0.50	0.04	0.66
Gizzard	1.20	1.15	1.16	1.18	0.10	0.98
Spleen	0.22	0.14	0.26	0.13	0.05	0.23
Thymus	0.24 <sup>ab</sup>	0.30 <sup>a</sup>	0.15 <sup>b</sup>	0.20 <sup>b</sup>	0.03	0.01
Bursa of Fabricius	0.18	0.16	0.22	0.32	0.07	0.37
Duodenum	0.64	0.70	0.68	0.65	0.04	0.73
Jejunum	1.12	1.14	1.13	1.16	0.09	0.99
Ileum	1.04	1.01	0.94	0.84	0.07	0.16
Caecum	0.49	0.50	0.56	0.66	0.10	0.63
Pancreas	0.32	0.35	0.27	0.32	0.03	0.24

<sup>a,b</sup> Means in the row with different letters show significant differences ( $P < 0.05$ ).

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diets supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diets supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error.

#### Hematological parameters of broiler chicks

Complete blood counts and serum biochemical parameters of broilers are shown in Tables 4 and 5, respectively. In general, there were no substantial differences in complete blood counts of broiler chicks. Regarding serum biochemical parameters, birds in PROB+AGP had lower ( $P = 0.04$ ) levels of ALT enzyme compared to CONT and AGP groups, but the difference was not significant when compared to birds in PROB

groups. Serum creatinine was lower ( $P < 0.01$ ) in PROB+AGP birds than in CONT and AGP birds, but the difference was not significant when compared to PROB birds. Birds in AGP group tended ( $P = 0.09$ ) to have a higher globulin concentration in the serum compared to birds in other treatments. PROB+AGP birds also tended to have higher albumin to globulin ratio when compared to other birds ( $P = 0.07$ ).

**Table 4.** Effects of zinc bacitracin and *Bacillus* mixture on complete blood counts in broiler chicks

Items	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P-value
	CONT	AGP	PROB	PROB+AGP		
Hemoglobin (g/dL)	8.76	9.74	9.38	9.07	0.79	0.84
Erythrocytes ( $10^6/\mu\text{L}$ )	2.05	2.26	2.11	2.06	0.16	0.80
Hematocrit (%)	29.1	31.3	30.3	30.0	2.76	0.95
MCV (fL)	140	138	144	145	2.80	0.31
MCH (pg)	42.5	42.9	44.3	43.8	1.03	0.56
MCHC (g/dL)	30.4	31.1	30.9	30.3	0.52	0.58
Leukocytes ( $10^3/\mu\text{L}$ )	19.1	21.8	25.6	22.1	2.36	0.36
Heterophils ( $10^3/\mu\text{L}$ )	0.46	0.66	0.68	0.63	0.10	0.40
Eosinophils ( $10^3/\mu\text{L}$ )	0.84	1.10	1.15	1.00	0.10	0.16
Lymphocytes ( $10^3/\mu\text{L}$ )	17.8	20.0	23.7	20.5	2.43	0.37
Thrombocytes ( $10^3/\mu\text{L}$ )	24.4	17.0	19.7	17.5	3.88	0.52

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diets supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diets supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error.

#### Intestinal bacterial populations of broiler chicks

The numbers of LAB in ileal digesta were higher ( $P = 0.05$ ) in PROB+AGP than in AGP birds, but the difference was not significant when

compared to CONT and PROB birds (Table 6). There were no significant differences in the counts of coliform and *C. perfringens* in both the ileal and cecal digesta of broilers.

**Table 5.** Effects of zinc bacitracin and *Bacillus* mixture on serum biochemical parameters and antibody titer in broiler chicks

Items <sup>#</sup>	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P-value
	CONT	AGP	PROB	PROB+AGP		
Total cholesterol (mg/dL)	133	140	142	153	14.5	0.81
HDL-c (mg/dL)	69.0	73.0	71.7	69.8	6.89	0.97
LDL-c (mg/dL)	54.8	72.6	53.2	74.2	12.4	0.51
Total triglyceride (mg/dL)	45.1	54.1	47.7	49.2	6.38	0.78
AST (U/L)	237	262	298	232	30.2	0.42
ALT (U/L)	4.27 <sup>a</sup>	6.28 <sup>a</sup>	2.15 <sup>ab</sup>	0.60 <sup>b</sup>	1.35	0.04
Total protein (g/dL)	2.67	3.15	2.63	2.63	0.20	0.21
Albumin (g/dL)	1.28	1.54	1.40	1.48	0.11	0.37
Globulin (g/dL)	1.39	1.62	1.24	1.15	0.13	0.09
A/G ratio	0.95	0.98	1.16	1.36	0.11	0.07
Uric acid (mg/dL)	4.09	4.26	7.44	4.69	1.31	0.26
Creatinine (mg/dL)	0.25 <sup>a</sup>	0.30 <sup>a</sup>	0.16 <sup>ab</sup>	0.07 <sup>b</sup>	0.04	<0.01
Antibody titer against NDV (Log <sub>2</sub> GMT)	2.60	2.60	3.60	2.20	0.64	0.47

<sup>a,b</sup> Means in the row with different letters show significant differences ( $P < 0.05$ ).

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diets supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diets supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error.

<sup>#</sup>HDL-c: high-density lipoprotein cholesterol, LDL-c: low-density lipoprotein cholesterol, AST: aspartate aminotransferase, ALT: alanine aminotransferase, A/G ratio: albumin to globulin ratio, NDV: Newcastle disease virus; GMT: geometric mean titer.

**Table 6.** Effects of zinc bacitracin and *Bacillus* mixture on selected bacterial populations in the intestine of broiler chicks

Items (log cfu/g)	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P-value
	CONT	AGP	PROB	PROB+AGP		
Ileum						
Coliform	7.25	7.14	7.00	7.05	0.12	0.46
<i>C. perfringens</i>	5.42	6.29	4.71	5.48	0.61	0.37
LAB <sup>#</sup>	9.23 <sup>ab</sup>	9.10 <sup>b</sup>	9.42 <sup>ab</sup>	9.65 <sup>a</sup>	0.13	0.05
Caecum						
Coliform	7.21	7.13	7.45	7.51	0.15	0.24
<i>C. perfringens</i>	6.35	5.83	6.12	7.36	0.51	0.19
LAB <sup>#</sup>	9.42	9.56	9.54	9.78	0.15	0.41

<sup>a,b</sup> Means in the row with different letters show significant differences ( $P < 0.05$ ).

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diet supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diet supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error; <sup>#</sup>LAB: lactic acid bacteria.

#### Concentrations of short chain fatty acids in cecal digesta of broiler chicks

The concentrations of SCFAs in cecal digesta of broilers did not differ across treatment groups (Table, 7).

#### Carcass traits of broiler chicks

Table 8 shows the carcass characteristics of

broiler chicks. Birds in AGP group had higher ( $P = 0.05$ ) relative weight of breast compared to other birds. No significant differences were observed in the relative weight of eviscerated carcass, breast, thigh, drumstick, wing, and abdominal fat of broilers. Crude protein and fat contents of breast meat were also similar across treatments.

**Table 7.** Effects of zinc bacitracin and *Bacillus* mixture on concentrations of short chain fatty acids in cecal digesta of broiler chicks

Items (mmol/kg)	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P-value
	CONT	AGP	PROB	PROB+AGP		
Acetic acid	76.1	65.0	62.8	66.3	9.70	0.77
Propionic acid	36.0	34.0	34.0	34.6	4.92	0.99
Butyric acid	21.2	22.7	23.7	21.9	3.81	0.97

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diet supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diet supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error.

**Table 8.** Effects of zinc bacitracin and *Bacillus* mixture on carcass characteristics of broiler chicks

Items	Dietary treatments <sup>†</sup>				SE <sup>‡</sup>	P-value
	CONT	AGP	PROB	PROB+AGP		
	% live weight					
Eviscerated carcass	60.2	63.4	61.5	61.3	1.58	0.56
	% Eviscerated carcass					
Breast	32.0 <sup>b</sup>	33.8 <sup>a</sup>	31.9 <sup>b</sup>	31.7 <sup>b</sup>	0.54	0.05
Thigh	16.4	15.6	16.8	16.8	0.46	0.26
Drumstick	15.4	15.4	15.7	15.1	0.76	0.98
Wing	14.2	13.6	14.3	14.8	0.44	0.31
Abdominal fat	0.69	0.99	0.81	0.61	0.25	0.73
	% proximate analysis					
Crude protein	23.2	23.7	22.7	23.1	0.38	0.34
Crude fat	0.53	0.60	0.57	0.57	0.04	0.65

<sup>a,b</sup> Means in the row with different letters show significant differences ( $P < 0.05$ ).

<sup>†</sup>CONT: basal diet without any supplementation, AGP: basal diet with 0.04% zinc bacitracin, PROB: basal diet supplemented with 0.5% *Bacillus* mixture, PROB+AGP: basal diet supplemented with 0.5% *Bacillus* mixture and 0.04% zinc bacitracin; <sup>‡</sup>SE: standard error.

## Discussion

A number of studies have reported inferior growth performance in broilers with low hatching weight compared to those with heavier hatchling weight (Butcher and Nilipour, 2002; Tona et al., 2004; Ulmer-Franco et al., 2010; Mendes et al., 2011; Michalczyk et al., 2011; Toghyani et al., 2011). Internal organs in broilers with low initial BW are also smaller (Sözcü and Ipek, 2017). As such, this study did not include a group of chicks with heavier hatching weight as negative control. As expected, the BW of broilers at 35 days of age in our study did not reach the target weight of Lohmann meat broilers, which have hatching weights of 42 g and weights of 2 kg on day 35 (Aviagen, 2007). The slower growth rate of birds with low hatching weights has been attributed to the slower differentiation of satellite cells due to the lower levels of myogenin and cyclin-dependent kinase inhibitor (p21) (both of which play an important role in muscle differentiation) (Andrés and Walsh, 1996). Satellite cells are the primary contributors to DNA synthesis for post-hatch muscle growth (Daughtry et al., 2017), so their slower differentiation may imply slower muscle hypertrophy or skeletal muscle growth of broiler chicks in later life. Supplementing of probiotic and antibiotic had no significant impact on the growth performance of birds in the present study, and the reasons for their ineffectiveness on growth performance in chicks with low initial BW remain largely unknown. Satellite cells are maximally active shortly after hatching and are very responsive to nutrition (Powell et al., 2013). In commercial hatcheries, broiler chicks often

rely on nutrients from the remaining yolk sac during handling and transport. Indeed, chicks produced from young breeder flocks had smaller proportion of yolk sac (Vieira and Moran Jr., 1998; Ulmer-Franco et al., 2010). It could be possible that chicks with low initial BW may have lower availability of nutrients during the post-hatch period compared to those with heavier initial BW, which could slow proliferation and differentiation of satellite cells in broiler chicks (Li et al., 2012). Therefore, the growth-promoting effects of probiotics and antibiotic zinc bacitracin may be hindered by the delay in post-hatch muscle hypertrophy during the first week of broiler life, which may also slow down the growth of broilers at subsequent ages. Further, a number of other factors may also attenuate the growth-promoting effects of probiotics and in-feed antibiotics in the present study. For example, the quality of feed provided to the chicks contained 6.79% crude fiber, which is higher than the standard fiber content in broiler feed. In this study, there was a tendency for birds in AGP and PROB groups to have higher feed intake than birds in CONT and PROB+AGP groups. However, treatment with in-feed antibiotics or probiotics did not affect the FCR of broiler chicks. Similar results were reported by Olnood et al. (2015) and Mahmoud et al. (2017), who showed no substantial effects of antibiotic zinc bacitracin and probiotic *Bacillus subtilis*, respectively, on the FCR of broiler chicks.

Lymphoid organ weight is a common marker of immune status in broiler chickens, as the



development of immune organs is crucial for the optimal production of immunoglobulins (Heckert *et al.*, 2002). In this study, the relative weight of thymus was lower in birds supplemented with *Bacillus* mixture or combination of *Bacillus* mixture and zinc bacitracin, compared to birds fed zinc bacitracin. Previous studies revealed an increase in thymus weight in birds following infections. For example, thymus size increased in birds after infections with *Chlamydia psittaci* HJ strain virus (Chu *et al.*, 2016), H7N3 avian influenza (HPAI) virus (Kapczynski *et al.*, 2013), and infectious bursal disease virus (IBDV) (Mutinda *et al.*, 2015). Therefore, the lower relative weight of thymus in birds treated with *Bacillus* mixture may be attributed to a lower degree of infections in the birds though this should be interpreted with caution as Rimondi *et al.* (2014) reported a significant reduction in thymus-to-body weight ratios in white leghorn chicks inoculated with Argentinean chicken anemia virus. Reasons for these differences across studies may be related to differences in the types or strains of birds tested, their holding conditions, the microorganisms used for infection (types, pathogenicity and doses), and the duration of infection. Nevertheless, birds in PROB and PROB+AGP tended to have lower globulin levels and higher A/G ratio when compared with those in AGP birds. This finding may support a lower infection incidence in birds supplemented with *Bacillus* mixture. Indeed, their serum globulin and lower A/G ratio have been shown to be associated with infections in birds (e.g. with *E. coli* in broiler chicks and Pekin ducks, Sharma *et al.*, 2015; Ogunbanwo *et al.*, 2004; Facon *et al.*, 2014). In our study, ALT enzyme levels were lower in birds in PROB+AGP and PROB compared to those in CONT and AGP groups. A decrease in serum ALT has been reported in broiler chicks in association with yeast supplementation (Aluwong *et al.*, 2013) and commercial probiotics containing *Bacillus subtilis*, *Bacillus coagulans* and *Saccharomyces boulardii* (Haque *et al.*, 2017). With regard to the role of vitamins and minerals, Huff *et al.* (1992) showed decreased ALT levels with supplementation of vitamin-mineral premixes in broiler chickens. Low ALT levels may be associated with a healthy liver or the absence of a toxicant-response in broiler chicks (Sugiharto *et al.*, 2016). Sharma *et al.* (2015) reported that an *E. coli* infection was associated with increased

ALT levels in bird serum. Hence, the lower serum ALT levels that we observed in probiotic-treated birds may be attributed to low levels of infections. In the present study, serum creatinine levels were lower in PROB+AGP and PROB birds compared to CONT and AGP birds. Previous studies by Mahmoud (2015) and Ismail (2017) showed an increase in serum creatinine levels in ducks and chickens infected by avian influenza viruses and NDV, respectively. Salim *et al.* (2011) showed a decrease in creatinine levels in broilers that were infected with *Salmonella typhimurium* but were supplemented with *Bacillus subtilis*. Probiotics seem to protect the chicks from infections or reduce the adverse effects of infections.

Results in the present study showed that treatment with *Bacillus* mixture was capable of increasing the population of LAB in the ilea of broiler chicks. Gao *et al.* (2017) also found a greater population of *Lactobacillus* in the gut of broilers fed *Bacillus subtilis*. Our data therefore suggest that *Bacillus* mixture treatment may favor the growth of good bacteria in the intestine of chicks, possibly because *Bacillus* spp. can produce antibacterial substances/antimicrobial peptides (AMPs) (Teixeira *et al.*, 2013), increase total SCFAs concentration in the ceca (Fujiwara *et al.*, 2009), and lower the intestinal pH (Knaul *et al.*, 2011). Such conditions may inhibit the growth of harmful bacteria and support the growth of beneficial bacteria such as LAB in the intestine of broilers. In regards to the antibacterial activity of *Bacillus*-based probiotic, our study did not show a significant effect of probiotic treatment on the numbers of coliform bacteria and *C. perfringens* in the intestine of broilers. There were also no effects on the concentrations of SCFAs in the cecal contents of broiler chicks, which is similar to findings from Olmood *et al.* (2015). However, Milián *et al.* (2013) showed an increase in acetic, propionic as well as total SCFAs in the cecum of broiler chicks treated with *Bacillus subtilis*. These differences across studies may be due to differences in probiotic strains and fibre content in diets.

In-feed antibiotic treatment resulted in higher relative weights of breast muscle in broiler chicks in our study. Similarly, Izat *et al.* (1990) reported that feeding bambermycins (2.2 ppm) or bacitracin methylene disalicylate (27.5 ppm) increased the weight of carcass and breast meat in broiler chicks. Denli *et al.* (2003) showed that feeding broiler chicks the antibiotic flavomycin

resulted in higher carcass weight. However, Landy *et al.* (2011) showed no effect of flavophospholipol on the carcass weight of broiler chicks. The higher breast weight may be related to the higher carcass and live weights of broilers (Sözcü and Ipek, 2017).

### Conclusion

Treatment with *Bacillus* mixture increases the intestinal population of LAB in low weight day-old broiler chicks. However, such treatment does

### References

- Ajuwon OR, Idowu OMO, Afolabi SA, Kehinde BO, Oguntola OO & Olatunbosun KO. 2011. The effects of dietary copper supplementation on oxidative and antioxidant systems in broiler chickens. *Archivos de Zootecnia*, 60: 275-282. DOI: 10.4321/S0004-05922011000200012
- Aluwong T, Hassan FB, Raji MA, Kawu MU, Dzenda T & Ayo JO. 2013. Effect of different levels of supplemental yeast on performance indices, serum enzymes and electrolytes of broiler chickens. *African Journal of Biotechnology*, 12: 5480-5485. DOI: 10.5897/AJB12.2588
- Andrés V & Walsh K. 1996. Myogenin, cell cycle withdrawal, and phenotypic differentiation are temporally separable events that precede cell fusion upon myogenesis. *Journal of Cell Biology*, 132: 657-666.
- AOAC. 1995. Official methods of analysis (16<sup>th</sup> ed). Association of Official Analytical Chemists, Washington DC, USA.
- Aviagen.2007.<http://www.incubatrici.padovan.it/allegati/LOHMANN.pdf> (accessed on 20 December 2017)
- Bolton W. 1967. Poultry Nutrition. MAFF Bulletin No.174, HMSO, London.
- Butcher GD & Nilipour AH. 2002. Broiler performance from cull eggs. IFAS Extension, University of Florida VM125; <https://edis.ifas.ufl.edu/pdffiles/VM/VM09200.pdf> (accessed on 20 December 2017)
- Chu J, Zhang Q, Zhang T, Han E, Zhao P, Khan A, He C & Wu Y. 2016. *Chlamydia psittaci* infection increases mortality of avian influenza virus H9N2 by suppressing host immune response. *Scientific Report*, 6: 29421, DOI: 10.1038/srep29421
- Daughtry MR, Berio E, Shen Z, Suess EJ, Shah N, Geiger AE, Berguson ER, Dalloul RA, Persia ME, Shi H & Gerrard DE. 2017. Satellite cell-mediated breast muscle regeneration decreases with broiler size. *Poultry Science*, 96: 3457-3464. DOI: 10.3382/ps/pex068
- Denli M, Okan F & Celik K. 2003. Effect of dietary probiotic, organic acid and antibiotic supplementation to diets on broiler performance and carcass yield. *Pakistan Journal of Nutrition*, 2: 89-91. DOI: 10.3923/pjn.2003.89-91
- Elshagabee FMF, Rokana N, Gulhane RD, Sharma C & Panwar H. 2017. *Bacillus* as potential probiotics: status, concerns, and future perspectives. *Frontiers in Microbiology*, 8: 1490. DOI: 10.3389/fmicb.2017.01490
- Facon C, Roman Y & Guerin J-L. 2014. Assessment of inflammatory status in poultry using plasma protein electrophoresis as a diagnostic tool. *Revue de Médecine Vétérinaire*, 165: 305-312.
- Fujiwara K-I, Yamazaki M, Abe H, Nakashima K, Yakabe Y, Otsuka M, Ohbayashi Y, Kato Y, Namai K, Toyokuni A, Miyaguchi Y & Nakamura Y. 2009. Effect of *Bacillus subtilis* var. *natto* fermented soybean on growth performance, microbial activity in the caeca and cytokine gene expression of domestic meat type chickens. *The Journal of Poultry Science*, 46: 116-122. DOI: 10.2141/jpsa.46.116
- Gao Z, Wu H, Shi L, Zhang X, Sheng R, Yin F & Gooneratne R. 2017. Study of *Bacillus subtilis* on growth performance, nutrition metabolism and intestinal microflora of 1 to 42 d broiler chickens. *Animal Nutrition*, 3: 109-113. DOI: 10.1016/j.aninu.2017.02.002
- Haque MI, Ahmad N & Miah MA. 2017. Comparative analysis of body weight and serum biochemistry in broilers supplemented with some selected probiotics and antibiotic growth promoters. *Journal of Advanced Veterinary and Animal Research*, 4: 288-294. DOI: 10.5455/javar.2017.d226

- 12 Heckert RA, Estevez I, Russek-Cohen E & Pettit RR. 2002. Effects of density and perch availability on the immune status of broilers. *Poultry Science*, 81: 451-457. DOI: 10.1093/ps/81.4.451
- 22 Huff WE, Kubena LF, Harvey RB & Philips TD. 1992. Effect of vitamin-mineral supplementation on growing chicks. *Poultry Science*, 71: 64-69. DOI: 10.3382/ps.0710064
- Islam M, 16 Bhuiyan MER, Begum MIA, Miah MA & Myenuddin M. 2004. Effect of vitamin-mineral premix supplementation on body weight and certain haemato-biochemical values in broiler chickens. *Bangladesh Journal of Veterinary Medicine*, 2: 45-48. DOI: 10.3329/bjvm.v2i1.1934
- Isn 53 HTH. 2017. Biochemical and hematological studies on the effect of neem (*Azadirachta Indica*) leaves aqueous extract on Newcastle disease vaccine and infection in broiler chickens. *International Journal of Recent Scientific Research*, 8: 15876-15884. DOI: 10.24327/ijrsr.2017.0803.0002
- 73 Isroli I, Yudiarti T, Widiastuti E & Sugiharto S. 2017. Probiotic *Bacillus* plus vitamins and minerals enhanced haemoglobin values and relative weight of ileum and improved feed conversion ratio of broilers during brooding period. *Livestock Research for Rural Development*, 29. <http://www.lrrd.org/40/lrrd29/11/sgh29212.html>
- 40 Izat AL, Colberg M, Reiber MA, Adams MH, Skinner 26 T, Cabel MC, Stilborn HL & Waldroup PW. 1990. Effects of different antibiotics on performance, processing characteristics, and parts yield of broiler chickens. *Poultry Science*, 69: 1787-1791. DOI: 10.3382/ps.0691787
- Kapczynski DR, Pantin-Jackwood M, Guzman SG, Ricardez Y, Spackman E, Bertran K, Suarez DL & Swayne DE. 2013. Characterization of the 2012 highly pathogenic avian influenza H7N3 virus isolated from poultry in an outbreak in Mexico: pathobiology and vaccine protection. *Journal of Virology*, 87: 9086-9096. DOI: 10.1128/JVI.00666-13
- Knap I, Kehlet AB, Bennedsen M, Mathis GF, Hofacre CL, Lumpkins BS, Jensen MM, Raun M & Lay A. 2011. *Bacillus subtilis* (DSM17299) significantly reduces *Salmonella* in broilers. *Poultry Science*, 90: 1690-1694. DOI: 10.3382/ps.2010-01056
- 1 Landy N, Ghulamkari Gh & Toghyani M. 2011. Performance, carcass characteristics, and immunity in broiler chickens fed dietary neem (*Azadirachta indica*) as alternative for an antibiotic growth promoter. *Livestock Science*, 142: 305-309. DOI: 10.1016/j.livsci.2011.08.017
- 57 Lee KW, Lillehoj HS, Lee SH, Jang SI, Park MS, Bautista DA, Ritte 46 D, Hong YH, Siragusa GR & Lillehoj EP. 2012. Effect of dietary antimicrobials on immune status in broiler chickens. *Asian-Australasian Journal of Animal Science*, 25: 382-392. DOI: 10.5713/ajas.2011.11259
- 45 Li Y, Yang X, Ni Y, Decuyper E, Buyse 11 eraert N, Grossmann R & Zhao R. 2012. Early-age feed restriction affects viability and gene expression of satellite cells isolated from the gastrocnemius muscle of broiler chicks. *Journal of Animal Science and Biotechnology*, 3: 33. DOI: 10.1186/2049-1891-3-33
- 18 Mahmoud EA. 2015. Hemato-biochemical and pathological changes on avian influenza in naturally infected domestic ducks in Egypt. *Veterinary World*, 8: 1177-1182. DOI: 10.14202/vetworld.2015.1177-1182
- Mahmoud KZ, Obei 23 BS, Al-Sadi MZ & Hatahet Sh R. 2017. Effect of *Bacillus subtilis* supplementation and dietary crude protein level on growth performance and intestinal morphological changes of meat type chicken. *Livestock Science*, 195: 99-104. DOI: 10.1016/j.livsci.2016.11.015
- Mendes AS, Paixao SJ, Restelatto R, Ruffatti R, Possenti JC, de Moura DJ, Morello GMZ & de Carvalho TMR. 2011. Effects of initial body weight and litter material on broiler production. *Brazilian Journal of Poultry Science*, 13: 165-170. DOI: 10.1590/S1516-635X2011000300001
- Michalczuk M, Stepińska M & Lukaszewicz M. 2011. Effect of the initial body weight of Ross 308 chicken broilers on the rate of growth. *Annals of Warsaw University of Life Sciences - SGGW, Animal Science*, 49: 121-125.
- Milián G, Rondón AJ, Pérez M, Bocourt R, Rodríguez 34 Z, Ranilla MJ, Rodríguez M & Carro MD. 2013. Evaluation of *Bacillus subtilis* biopreparations as growth promoters in chickens. *Cuban Journal of Agricultural Science*, 47: 61-66. 4
- M'Sadeq SA, Wu S, Swick RA & Choct M. 2015. Towards the control of necrotic enteritis in broiler chickens with in-feed antibiotics phasing-out worldwide. *Animal Nutrition*, 1: 1-11. DOI: 10.72 5/j.aninu.2015.02.004
- Mutinda WU, Nja 21 W, Nyaga PN, Bebora LC, Mbuthia PG, Kemboi D, Githinji JWK &

- Muriuki A. 2015. Isolation of infectious bursal disease virus using indigenous chicken embryos in Kenya. *International Scholarly Research Notices*. DOI: 10.1155/2015/464714
- Ogunbanwo ST, Sanni AI & Onilude AA. 2004. Influence of bacteriocin in the control of *Escherichia coli* infection of broiler chickens in Nigeria. *World Journal of Microbiology and Biotechnology*, 20: 51-56. DOI: 10.1023/B:WIBI.0000013311.43842.74
- Olnood CG, Beski SSM, Choct M & Iji PA. 2015. Novel probiotics: their effects on growth performance, gut development, microbial community and activity of broiler chickens. *Animal Nutrition*, 1: 184-191. DOI: 10.1016/j.aninu.2015.07.003
- Powell DJ, McFarland DC, Cowieson AJ, Muir WI & Velleman SG. 2013. The effect of nutritional status on myogenic satellite cell proliferation and differentiation. *Poultry Science*, 92: 2163-2173. DOI: 10.3382/ps.2013-03107
- Rimondi A, Pinto S, Olivera V, Dibárbora M, Pérez-Filgueira M, Craig MI & Pereda A. 2014. Comparative histopathological and immunological study of two field strains of chicken anemia virus. 45: 102. DOI: 10.1186/s13567-014-0102-y
- Salim HA, Abd-Allah OA & Fararh KM. 2011. Effect feeding probiotic on haematological, biochemical properties and immune response in broiler. *Benha Veterinary Medical Journal*, 22: 35-43.
- Sanda ME. 2015. Effects of vitamin-mineral supplement on the immune response of broilers to Newcastle Disease vaccination. *International Journal of Agricultural and Veterinary Sciences*, 1:10-13. DOI: 10.18819/ija.2015.1542
- SAS (Statistical Analysis System). 1985. SAS/STAT 5. User's Guide. SAS Institute Inc. Cary, North Carolina.
- Sharma V, Jakhar KK, Nehra V & Kumar S. 2015. Biochemical studies in experimentally *Escherichia coli* infected broiler chicken supplemented with neem (*Azadirachta indica*) leaf extract. *Veterinary World*, 8: 1340-1345. DOI: 10.14202/vetworld.2015.1340-1345
- Simon O. 2005. Micro-organisms as feed additives - probiotics. *Advances in Pork Production*, 16: 161-167. <http://www.prairieswine.com/pdf/2451.pdf> (accessed on 20 December 2017).
- SNI (Indonesian National Standard). 2006. Standard for broiler feed (SNI 01-3930-2006). National Standardization Agency of Indonesia, Jakarta, Indonesia (article in Bahasa).
- Sözcü A & Ipek A. 2017. Effects of egg weight on chick and organ development, growth and slaughter traits in Pekin ducks. *Journal of Biological & Environmental Sciences*, 11: 87-92. <http://jbes.uludag.edu.tr/PDFDOSYALAR/32/mak04.pdf> (accessed on 20 December 2017)
- Sugiharto S, Lauridsen C & Jensen BB. 2015. Gastrointestinal ecosystem and immunological responses in *E. coli* challenged pigs after weaning fed liquid diets containing whey permeate fermented with different lactic acid bacteria. *Animal Feed Science and Technology*, 207: 278-282. DOI: 10.1016/j.anifeeds.2015.06.019
- Sugiharto S. 2016. Role of nutraceuticals in gut health and growth performance of poultry. *Journal of the Saudi Society of Agricultural Sciences*, 15: 99-111. DOI: 10.1016/j.jssas.2014.06.001
- Sugiharto S, Yudiarti T & Isroli I. 2016. Performances and haematological profile of broilers fed fermented dried cassava (*Manihot esculenta* Crantz). *Tropical Animal Health and Production*, 48: 1337-1341. DOI: 10.1007/s11250-016-1098-2
- Sugiharto S, Yudiarti T, Isroli I, Widiastuti E & Putra FD. 2017. Effect of dietary supplementation with *Rhizopus oryzae* or *Clrysonilia crassa* on growth performance, blood profile, intestinal microbial population, and carcass traits in broilers exposed to heat stress. *Archives Animal Breeding*, 60: 347-356. DOI: 10.5194/aab-60-347-2019
- Teixeira ML, Rosa AD & Brandelli A. 2013. Characterization of an antimicrobial peptide produced by *Bacillus subtilis* subsp. *Spizezinii* showing inhibitory activity towards *Haemophilus parasuis*. *Microbiology*, 159: 980-988. DOI: 10.1099/mic.0.062828-0
- Toghyani M, Toghyani M, Gheisari A, Ghalamkari G & Eghbalsaied S. 2011. Evaluation of cinnamon and garlic as antibiotic growth promoter substitutions on performance, immune responses, serum biochemical and haematological parameters in broiler chicks. *Livestock Science*, 138: 167-173. DOI: 10.1016/j.livsci.2010.12.018
- Tona K, Onagbesan O, De Ketelaere B, Decuyper E & Bruggeman V. 2004. Effects of age of broiler breeders and egg storage on egg quality, hatchability, chick quality, chick weight and

- chick post-hatch growth to forty two days. The Journal of Applied Poultry Research, 13: 10-18. DOI: 10.1093/japr/13.1.10
- 6 Ulmer-Franco AM, Fasenko GM & Christopher EEO. 2010. Hatching egg characteristics, chick quality, and broiler performance at 2 breeder flock ages and from 3 egg weights. Poultry Science, 89: 2735-2742. DOI: 10.3382/ps.2009-00403
- 25 Vieira SL & Moran Jr. ET. 1998. Eggs and chicks from broiler breeders of extremely different age. The Journal of Applied Poultry Research, 7: 372-376. DOI: 10.1093/japr/7.4.372
- Villegas P. 1956. Avian virus diseases laboratory manual. College of Veterinary Medicine, University of Georgia, Athens, Georgia, USA.
- 20 Wyatt CL, Weaver WD & Beane WL. 1985. Influence of egg size, egg shell quality and post hatch holding time on broiler performance. Poultry Science, 64: 2049-2055. DOI: 10.3382/ps.0642049

ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

7%

STUDENT PAPERS

---

PRIMARY SOURCES

---

- |   |  |     |
|---|--|-----|
| 1 | ZHOU, Qian, Si-si WANG, Guang YANG, Wen ZHAO, and Hui-ling LI. "Development and evaluation of a herbal formulation with anti-pathogenic activities and probiotics stimulatory effects", Journal of Integrative Agriculture, 2016.<br>Publication | <1% |
| 2 | saarcagri.net<br>Internet Source   | <1% |
| 3 | anis.au.dk<br>Internet Source  | <1% |
| 4 | journal-cdn.frontiersin.org<br>Internet Source   | <1% |
| 5 | info.ibaraki.ac.jp<br>Internet Source  | <1% |
| 6 | Løtvedt, Pia, and Per Jensen. "Effects of Hatching Time on Behavior and Weight Development of Chickens", PLoS ONE, 2014.<br>Publication  | <1% |
-

7

Vasudevan Gowthaman, Shambu Dayal Singh, Kuldeep Dhama, Palani Srinivasan, Sellappan Saravanan, Thippichettypalayam R. Gopala Krishna Murthy, Muthannan Andavar Ramakrishnan. "Molecular Survey of Respiratory and Immunosuppressive Pathogens Associated with Low Pathogenic Avian Influenza H9N2 Subtype and Virulent Newcastle Disease Viruses in Commercial Chicken Flocks", The Journal of Poultry Science, 2017

Publication

<1%

8

Submitted to University of New South Wales

Student Paper

<1%

9

[www.lume.ufrgs.br](http://www.lume.ufrgs.br)

Internet Source

<1%

10

Gayatri Suresh, Ratul Kumar Das, Satinder Kaur Brar, Tarek Rouissi, Antonio Avalos Ramirez, Younes Chorfi, Stephane Godbout. "Alternatives to antibiotics in poultry feed: molecular perspectives", Critical Reviews in Microbiology, 2017

Publication

<1%

11

J Vierck. "SATELLITE CELL REGULATION FOLLOWING MYOTRAUMA CAUSED BY RESISTANCE EXERCISE", Cell Biology International, 2000

Publication

<1%

12

[helda.helsinki.fi](http://helda.helsinki.fi)

Internet Source

&lt;1%

13

[journals.co.za](http://journals.co.za)

Internet Source

&lt;1%

14

[sci.ui.edu.ng](http://sci.ui.edu.ng)

Internet Source

&lt;1%

15

[ansci.osu.edu](http://ansci.osu.edu)

Internet Source

&lt;1%

16

[bdresearchpublications.com](http://bdresearchpublications.com)

Internet Source

&lt;1%

17

Submitted to University of Newcastle upon Tyne

Student Paper

&lt;1%

18

[actavet.vfu.cz](http://actavet.vfu.cz)

Internet Source

&lt;1%

19

[efsa.onlinelibrary.wiley.com](http://efsa.onlinelibrary.wiley.com)

Internet Source

&lt;1%

20

ATASOY, Fatih, ONBASILAR, Esin Ebru and APAYDIN, Seval. "Denizli ve ticari tavuk sürülerinde yumurta kalite özelliklerinin karşılaştırılması", Lalahan Hayvancılık Araştırma Enstitüsü, 2001.

Publication

&lt;1%

21

Submitted to University of the Free State

Student Paper

&lt;1%



22 Rashid, Mohammad Harun Ar, Nazim Ahmad, Mohammad Rohul Amin, and Mohammad Lalmoddin Mollah. "Effects of selected vitamins and minerals on growth rate and hematological parameters in broilers", Asian Journal of Medical and Biological Research, 2016.  
Publication

---

23 [m.scirp.org](http://m.scirp.org)  
Internet Source

---

24 Submitted to Universitas Negeri Semarang  
Student Paper

---

25 Maiorka, A, E Santin, AVF Silva, KS Routman, JM Pizauro Jr., and M Macari. "Effect of broiler breeder age on pancreas enzymes activity and digestive tract weight of embryos and chicks", Revista Brasileira de Ciãncia Avãcola, 2004.  
Publication

---

26 [www.nzfsa.govt.nz](http://www.nzfsa.govt.nz)  
Internet Source

---

27 [besjournals.onlinelibrary.wiley.com](http://besjournals.onlinelibrary.wiley.com)  
Internet Source

---

28 Submitted to Florida Memorial College  
Student Paper

---

29 Submitted to University of Mosul  
Student Paper

---

30	<a href="http://www.pork.org">www.pork.org</a> Internet Source	<1%
31	<a href="http://pjms.com.pk">pjms.com.pk</a> Internet Source	<1%
32	<a href="http://www.wjgnet.com">www.wjgnet.com</a> Internet Source	<1%
33	A. Asghar Saki, P. Zamani, M. Rahmati, H. Mahmoudi. "The effect of cage density on laying hen performance, egg quality, and excreta minerals", The Journal of Applied Poultry Research, 2012 Publication	<1%
34	<a href="http://repository.unhas.ac.id">repository.unhas.ac.id</a> Internet Source	<1%
35	Submitted to Khon Kaen University Student Paper	<1%
36	A. Bandegan. "Standardized ileal amino acid digestibility in wheat distillers dried grains with solubles for broilers", Poultry Science, 12/01/2009 Publication	<1%
37	Ihara, Yasuhiro, Hideki Hyodo, Shin Sukegawa, Hiroshi Murakami, and Fumiki Morimatsu. "Isolation, characterization, and effect of administration in vivo, a novel probiotic strain	<1%

from pig feces : Effect of the Novel Probiotics in Swine", Animal Science Journal, 2013.

Publication

38

Rimondi, Agustina, Silvina Pinto, Valeria Olivera, Marina Dibárhora, Mariano Pérez-Filgueira, María Craig, and Ariel Pereda. "Comparative histopathological and immunological study of two field strains of chicken anemia virus", Veterinary Research, 2014.

Publication

<1%

39

[iopscience.iop.org](http://iopscience.iop.org)

Internet Source

<1%

40

ŞEHU, Adnan, YALÇIN, Sakine and ÖNOL, Ahmet G.. "Farklı enerji düzeylerindeki rasyonlara virginiamycin katkısının bıldırcınlarda besi performansı ve karkas randımanı üzerine etkisi", TUBITAK, 2002.

Publication

<1%

41

[www.revmedvet.com](http://www.revmedvet.com)

Internet Source

<1%

42

Commercial Chicken Meat and Egg Production, 2002.

Publication

<1%

43

Gilbert C. Sigua. "Recycling biosolids and lake-dredged materials to pasture-based animal agriculture: alternative nutrient sources for

<1%

forage productivity and sustainability. A review",  
Agronomy for Sustainable Development,  
01/2009

Publication

---

44

Submitted to University of New England

Student Paper

<1%

---

45

[bmcgenomics.biomedcentral.com](http://bmcgenomics.biomedcentral.com)

Internet Source

<1%

---

46

Leon J Broom. "Gut barrier function: Effects of  
(antibiotic) growth promoters on key barrier  
components and associations with growth  
performance", Poultry Science, 2018

Publication

<1%

---

47

Tingtao Chen. "Application of denaturing  
gradient gel electrophoresis to microbial  
diversity analysis in Chinese Douchi", Journal of  
the Science of Food and Agriculture, 2012

Publication

<1%

---

48

Submitted to Volunteer State Community  
College

Student Paper

<1%

---

49

Rude, B. J., and D. L. Rankins. "Effect of Hay  
and Molasses on the Utilization of Broiler Litter-  
Based Diets by Beef Cattle", Journal of Applied  
Animal Research, 1999.

Publication

<1%

---

50

B Kierończyk, M Sassek, E Pruszyńska-Oszmałek, P Kołodziejcki, M Rawski, S Świątkiewicz, D Józefiak. "The physiological response of broiler chickens to the dietary supplementation of the bacteriocin nisin and ionophore coccidiostats", Poultry Science, 2017

Publication

---

<1%

51

Takahashi, Masahide. "Effect of New Composite Probiotic Microecology Additives on Growth Performance and Immunity in Chicks", Medicine and Biopharmaceutical, 2016.

Publication

---

<1%

52

S. Yalçın, S. Özkan, E. Coskuner, G. Bilgen, Y. Delen, Y. Kurtulmus, T. Tanyalçın. "Effects of strain, maternal age and sex on morphological characteristics and composition of tibial bone in broilers", British Poultry Science, 2010

Publication

---

<1%

53

Sumaira Saleem, Gulzar Muhammad, Muhammad Ajaz Hussain, Syed Nasir Abbas Bukhari. " A comprehensive review of phytochemical profile, bioactives for pharmaceuticals, and pharmacological attributes of ", Phytotherapy Research, 2018

Publication

---

<1%

54

Okorie-Kanu, Christian O., Onyinye J. Okorie-Kanu, and John O. A. Okoye. "Blood

<1%

biochemistry responses of ducks infected with a velogenic Newcastle disease virus",  
Comparative Clinical Pathology, 2016.

Publication

55

[aas.bf.uni-lj.si](http://aas.bf.uni-lj.si)

Internet Source

<1%

56

[iai.asm.org](http://iai.asm.org)

Internet Source

<1%

57

[proteomesci.biomedcentral.com](http://proteomesci.biomedcentral.com)

Internet Source

<1%

58

L. Billard, E. Song, M. Y. Shim, G. M. Pesti.  
"Interpreting experiments on egg production--  
Statistical considerations", Poultry Science,  
2013

Publication

<1%

59

[www.ijat-aatsea.com](http://www.ijat-aatsea.com)

Internet Source

<1%

60

[curis.ku.dk](http://curis.ku.dk)

Internet Source

<1%

61

Jaksch, D.. "The effect of ozone treatment on  
the microbial contamination of pork meat  
measured by detecting the emissions using  
PTR-MS and by enumeration of  
microorganisms", International Journal of Mass  
Spectrometry, 20041215

Publication

<1%

62

Maneewan, Chamroon, Apichai Mekbungwan, Wasin Charerntantanakul, Kohsho Yamauchi, and Koh-en Yamauchi. "Effects of dietary *Centella asiatica* (L.) Urban on growth performance, nutrient digestibility, blood composition in piglets vaccinated with *Mycoplasma hyopneumoniae* : Effect of *Centella asiatica* on piglets with mycoplasma", *Animal Science Journal*, 2014.

Publication

&lt;1%

63

Vázquez, Ramón Silva, Lorenzo Antonio Durán Meléndez, Eduardo Santellano Estrada, Carlos Rodríguez Muela, Guillermo Villalobos Villalobos, Gerardo Méndez Zamora, and Michael E. Hume. "Performance of broiler chickens supplemented with Mexican oregano oil (*Lippia berlandieri* Schauer)", *Revista Brasileira de Zootecnia*, 2015.

Publication

&lt;1%

64

Yamauchi, K., T. Buwjoom, K. Koge, and T. Ebashi. "Histological alterations of the intestinal villi and epithelial cells in chickens fed dietary sugar cane extract", *British Poultry Science*, 2006.

Publication

&lt;1%

65

Sadeghi, Ghorbanali, Mahmood Habibian, Ali Raei, Danial Farhadi, and Naser Khateri. "Effects of dietary supplementation of *Ferulago*

&lt;1%

angulata (Schlecht.) Boiss powder on growth performance, carcass characteristics, and gut microflora and pH in broiler chicks", *Comparative Clinical Pathology*, 2015.

Publication

---

66

Houshmand, M., K. Azhar, I. Zulkifli, M. H. Bejo, and A. Kamyab. "Effects of prebiotic, protein level, and stocking density on performance, immunity, and stress indicators of broilers", *Poultry Science*, 2012.

Publication

---

67

Tagang, Aluwong, B Hassan Fatima, A Raji Moshood, U Kawu Mohamed, Dzenda Tavershima, and O Ayo Joseph. "Effect of different levels of supplemental yeast on performance indices, serum enzymes and electrolytes of broiler chickens", *AFRICAN JOURNAL OF BIOTECHNOLOGY*, 2013.

Publication

---

68

Lizama, L.C.. "Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets", *Animal Feed Science and Technology*, 198805

Publication

---

69

M F Karimy, B Sutrisno, A Agus, A E Suryani, L Istiqomah, E Damayanti. "Aflatoxin effect on erythrocyte profile and histopathology of broilers

<1%

<1%

<1%

<1%



given different additives", IOP Conference  
Series: Earth and Environmental Science, 2017

Publication

70

[repository.uaf.edu.pk](http://repository.uaf.edu.pk)

Internet Source

<1%

71

Hesong Wang, Xueqin Ni, Xiaodan Qing, Lei Liu, Jing Lai, Abdul Khalique, Guangyao Li, Kangcheng Pan, Bo Jing, Dong Zeng. "Probiotic Enhanced Intestinal Immunity in Broilers against Subclinical Necrotic Enteritis", *Frontiers in Immunology*, 2017

Publication

<1%

72

Sultan, Serageldeen, Nabila Osman, Moemen A. Mohamed, Ahmed I. Ahmed, Rabab A. M. Abdallah, Marwa Faraouk, and Mokhtar Taha. "Infectious bursal disease vaccine ameliorates velogenic Newcastle disease virus infection in immunopotential chickens", *Comparative Clinical Pathology*, 2015.

Publication

<1%

73

[www.neliti.com](http://www.neliti.com)

Internet Source

<1%

74

[jcea.agr.hr](http://jcea.agr.hr)

Internet Source

<1%

75

[flex.flinders.edu.au](http://flex.flinders.edu.au)

Internet Source

<1%

76

Raghunathan Srianand, Patrick Petitjean, Cédric Ledoux, Gary Ferland, Gargi Shaw. "The VLT-UVES survey for molecular hydrogen in high-redshift damped Lyman  $\alpha$  systems: physical conditions in the neutral gas", Monthly Notices of the Royal Astronomical Society, 2005

Publication

&lt;1%

77

Y. Pinchasov. "Comparison of post-hatch holding time and subsequent early performance of broiler chicks and Turkey poults", British Poultry Science, 3/1993

Publication

&lt;1%

78

Velmurugu RAVINDRAN. "Performance and welfare of broilers as affected by stocking density and zinc bacitracin supplementation", Animal Science Journal, 2/2006

Publication

&lt;1%

79

PUVAČA, N., D. LJUBOJEVIĆ, LJ. KOSTADINOVIĆ, J. LEVIĆ, N. NIKOLOVA, B. MIŠČEVIĆ, T. KÖNYVES, D. LUKAČ, and S. POPOVIĆ. "Spices and herbs in broilers nutrition: hot red pepper (*Capsicum annum* L.) and its mode of action", World s Poultry Science Journal, 2015.

Publication

&lt;1%

80

M. R. Daughtry, E. Berio, Z. Shen, E. J. R. Suess et al. "Satellite cell-mediated breast

&lt;1%

muscle regeneration decreases with broiler size", Poultry Science, 2017

Publication

81

[jast.modares.ac.ir](http://jast.modares.ac.ir)

Internet Source

<1%

82

A. Daneshmand, GH. Sadeghi, A. Karimi, A. Vaziry. "Effect of oyster mushroom (*Pleurotus ostreatus*) with and without probiotic on growth performance and some blood parameters of male broilers", *Animal Feed Science and Technology*, 2011

Publication

<1%

83

A. Biswas. "Effect of higher levels of dietary selenium on production performance and immune responses in growing Japanese quail", *British Poultry Science*, 8/1/2006

Publication

<1%

84

S. M. ABDULRAHIM. "Effect of *Lactobacillus acidophilus* and zinc bacitracin as dietary additives for broiler chickens", *British Poultry Science*, 3/1/1999

Publication

<1%

85

S. Sugiharto, B. B. Jensen, M. S. Hedemann, C. Lauridsen. " Comparison of casein and whey in diets on performance, immune responses and metabolomic profile of weanling pigs challenged with F4 ", *Canadian Journal of Animal Science*,

<1%

2014

Publication

---

86

[www.ifrj.upm.edu.my](http://www.ifrj.upm.edu.my)

Internet Source

<1%

---

87

Ahmed M. Erfan, Abdullah A. Selim, Mahmoud M. Naguib. "Characterization of full genome sequences of chicken anemia viruses circulating in Egypt reveals distinct genetic diversity and evidence of recombination", *Virus Research*, 2018

Publication

<1%

---

88

[austinpublishinggroup.com](http://austinpublishinggroup.com)

Internet Source

<1%

---

89

Verma, J., T.S. Johri, B.K. Swain, and S. Ameena. "Effect of graded levels of aflatoxin, ochratoxin and their combinations on the performance and immune response of broilers", *British Poultry Science*, 2004.

Publication

<1%

---

90

Saleh, Ahmed A. Amber, Khairy Magd, Moha. "Integrative effects of feeding *Aspergillus awamori* and fructooligosaccharide on growth performance a", *BioMed Research International*, Annual 2014 Issue

Publication

<1%

---

91

X. Zhang, D. Zhang, L. Wang, D. Liu. "Molecular cloning and functional characterization of

<1%

myogenin promoter in Min pig", *Acta Agriculturae Scandinavica, Section A — Animal Science*, 2017

Publication

---

92

Darryl C Wilkie, Andrew G Van Kessel, Lisa J White, Bernard Laarveld, Murray D Drew. "Dietary amino acids affect intestinal populations in broiler chickens ", *Canadian Journal of Animal Science*, 2005

<1%

Publication

---

93

Sanja J. Popović, Ljiljana M. Kostadinović, Nikola M. Puvača, Bojana M. Kokić, Ivana S. Čabarkapa, Olivera M. Đuragić. "Potential of wormwood (*Artemisia absinthium*) as a feed supplement in rabbit diet: effect on controlling rabbit coccidiosis, antioxidative systems and growth performance", *Veterinarski arhiv*, 2017

<1%

Publication

---

94

Storlien, T. M., E. Prestlie, K. A. Beauchemin, T. A. McAllister, A. Iwaasa, and O. M. Harstad. "Supplementation with crushed rapeseed causes reduction of methane emissions from lactating dairy cows on pasture", *Animal Production Science*, 2015.

<1%

Publication

---

95

Pereverzeva, E.. "Intravenous tolerance of a nanoparticle-based formulation of doxorubicin in

<1%

healthy rats", Toxicology Letters, 20080421

Publication

---

96

"PERFORMANCE AND TIBIA  
CHARACTERISTICS OF BROILERS FED  
DIETS CONTAINING RAW OR TREATED OAK  
ACORN (QUE", Journal of Animal and Plant  
Sciences, Feb 28 2018 Issue

<1%

Publication

---

Exclude quotes On

Exclude matches Off

Exclude bibliography On