

vanced_Veterinary_and_Animal_
Research_-_5_3_332-
342.Agust_18.pdf
by T Yudiarti

Submission date: 20-Nov-2018 01:08PM (UTC+0700)

Submission ID: 1042454077

File name: vanced_Veterinary_and_Animal_Research_-_5_3_332-342.Agust_18.pdf (706.41K)

Word count: 8124

Character count: 41683

Original Article

The effect of fungi-origin probiotic *Chrysonilia crassa* in comparison to selected commercially used feed additives on broiler chicken performance, intestinal microbiology, and blood indices

Sugiharto Sugiharto^{1#}, Turrini Yudiarti¹, Isroli Isroli¹, Endang Widiastuti¹, Hanny Indrat Wahyuni¹ and Tri Agus Sartono¹

• Received: June 14, 2018 • Revised: Aug 1, 2018 • Accepted: Aug 2, 2018 • Published Online: Aug 5, 2018



AFFILIATIONS

¹Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Central Java Province, Indonesia.

ABSTRACT

Objective: The objective of the current study was to investigate the influence of probiotic *Chrysonilia crassa* in comparison with zinc bacitracin, commercial probiotic *Bacillus subtilis*, and herbal medicine waste on growth, intestinal microbiology, and blood indices of broilers.

Materials and methods: Three hundreds of Lohmann day-old chicks were allocated to control diet (basal diet; CONT), basal diet with antibiotic zinc bacitracin (AZB), basal diet with *B. subtilis* (PROB), basal diet with *C. crassa* (PROC), and basal diet with herbal medicine waste (HERBW). Sample collections were conducted on day 34 of the experiment.

Results: PROB showed greater ($P<0.05$) body weight than CONT chicks. Leukocytes and lymphocytes numbers were higher ($P<0.05$) in HERBW than in birds of other treatments. Compared to CONT and HERBW, PROC birds showed higher ($P<0.05$) level of vaccine titer to Newcastle disease virus. CONT had lower ($P<0.05$) and higher ($P<0.05$) total protein and globulin, and the ratio of albumin to globulin (A/G ratio) in serum, respectively, compared to other chicks. Higher level ($P<0.05$) of albumin was observed in PROB relative to CONT and PROC. Lower uric acid ($P<0.05$) was seen in PROC when compared with CONT and PROB. PROC had higher ($P<0.05$) aspartate aminotransferase than AGP, PROB, and HERBW. Ileal coliform was decreased ($P<0.05$) in PROB and PROC, relative to CONT and HERBW.

Conclusion: Probiotics were capable of improving the growth, immune responses, and intestinal bacterial populations of broilers. The effects of probiotics *C. crassa* corresponded to that of commercial probiotic *B. subtilis* and antibiotic growth promoters.

CORRESPONDENCE:

#Sugiharto Sugiharto,
 Department of Animal Science, Faculty of
 Animal and Agricultural Sciences,
 Diponegoro University, Semarang, Central
 Java Province, Indonesia.
 E-mail: sgb_ndip@yahoo.co.id

KEYWORDS

Antibiotic growth promoters; Broiler; *Chrysonilia crassa*; Growth; Health; Probiotic

INTRODUCTION

Antibiotics have long been included sub-therapeutically in diets as an antimicrobial agent as well as growth promoters in broiler production. The use of antibiotic growth promoters (AGPs) is, however, no longer permitted in most countries due to the concern about antimicrobial resistance both in birds and humans as consumers (Abudabos et al., 2017a). Indeed, there are some problems related to performances and health upon the removal of in-feed antibiotics in broiler production (Sugiharto, 2016). The health problems in poultry are often the result of the imbalance of microbial community in the gut, which may be aggravated by the withdrawal of antibiotics from feeds (Abid et al., 2016). The change in the balance of bacteria in the gut has been shown to induce immunological disturbances in broiler chickens (Simon et al., 2016; Schokker et al., 2017). This immune stress may lead to physiological disorder and thus negatively affect broiler performances (Feng et al., 2012). Given the interplay between gut microbiota and immune system (Simon et al., 2016), feeding any compounds that can maintain the balanced microbiota in the intestine seems beneficial to support the healthy growth of broiler chickens post in-feed antibiotics removal.

The potentials of probiotics in establishing and maintaining the balance of intestinal microbiota and improving the immunological competences and growth performances of broiler chickens have been documented. However, there is still a hesitation among poultry nutritionists to incorporate probiotics in broiler rations since the results of some *in vivo* experiments with probiotics have been inconsistent (Sugiharto, 2016). Several studies pointed out that the benefits of probiotics for poultry and other farm animals are hosts- and species/strain-specific (Sugiharto et al., 2015a; Sugiharto, 2016; Abudabos et al., 2017b). For the latter reason, attempts are still needed to find novel probiotic microorganisms that can be a satisfactory alternative to in-feed antibiotics for broiler chickens. In our earlier study, we have isolated the fungus *Chrysonilia crassa* from the ileum of the Indonesian indigenous chickens (Yudiarti et al., 2012a) and shown its probiotic activities *in vitro* (Yudiarti et al., 2013). In the *in vivo* experiment, the fungus was capable of controlling bacterial population in the intestine and improving the duodenal villi development of the Indonesian indigenous chicks (Yudiarti et al., 2012b). Our recent study further confirmed the benefits of *C. crassa* in improving the physiological conditions and antioxidant status of broilers under heat stress (Sugiharto et al., 2017a). In the present

study, the benefits of *C. crassa* were evaluated specifically for broiler chickens in comparison with commercial probiotics and other feed additives.

Herbal products such as Chinese herbal medicine have long been employed to substitute the use of AGPs for broilers. The use of such products resulted in improved intestinal bacterial community and growth rate in broiler chicks (Guo et al., 2004; Han et al., 2012; Zawqari et al., 2016). However, the application of herbal products in poultry production seems to be impractical as it has to compete with humans for medicine (Elfahmi et al., 2014). In Indonesia, herbal medicine (called *Jamu*) industries have been growing substantially in the recent years. This may consequently increase the waste generated during *Jamu* production (Soetrisnanto et al., 2012). According to Kisworo et al. (2016), most of the *Jamu* waste contains phenols, tannins, flavonoids, alkaloids, saponins, and essential oils that in general possess antibacterial activities. Taking the latter properties into consideration, feeding *Jamu* waste was, therefore, expected to improve the microbial community and thus, the well-being of broiler chickens. To date, the comparative effects among probiotic *C. crassa*, antibiotic zinc bacitracin (AZB), commercial probiotic *Bacillus subtilis*, and herbal medicine waste as the alternatives to AGPs on the production and health performances of broiler chickens have never been documented. Therefore, the objective of the current study was to evaluate the influence of fungi-origin probiotic *C. crassa* in comparison with AZB, commercial probiotic *B. subtilis*, and herbal medicine waste on growth performance, intestinal microbiology, and blood parameters of broilers.

MATERIALS AND METHODS

Ethical approval: The rearing and handling of chicks during the experiment were conducted according to the animal ethics mentioned in the Law of the Republic of Indonesia number 18, 2009.

Probiotics and herbal medicine waste preparations: Probiotic *C. crassa* was prepared based on Sugiharto et al. (2017a). Briefly, *C. crassa* was retrieved from the fungal stock (preserved on a potato dextrose agar [PDA; Merck KGaA, Darmstadt, Germany] and kept at 4°C). The fungal colonies were streaked on PDA and incubated at 38°C for 2 days. After being dislodged from the PDA, the fungal mycelia were diluted in sterile distilled water (100 ml). The sterile rice bran [200 g; 85.9% dry matter (DM)] was then inoculated with the above-mentioned suspension. The inoculum contained 1×10^{12} cfu/ml for each inoculation. Following 4 days of aerobic incubation

at room temperature, the number of fungal colonies grown in rice bran was determined. To do proximate analysis, a sample of fermented rice bran (100 g) was obtained. Standard plate count method showed that the fungi growing in rice bran were 1×10^{11} cfu/gm. Based on proximate analysis (AOAC, 1995), *C. crassa*-fermented rice bran contained (as a dry basis) 2,659 kcal/kg gross energy (GE), 11.6% crude protein (CP), 5.64% crude fat (CF), 23.1% crude fiber, and 12.0% total ash. The commercial probiotic (Baymix Grobig) was obtained from PT. Bayer Indonesia (Jakarta, Indonesia). It contained *B. subtilis* at a minimum 1×10^{10} viable spores/gm. The herbal medicine waste was obtained from PT. Sidomuncul, Tbk. (Semarang, Central Java, Indonesia). The waste was generated from the production of *Jamu* "Tolak Angin" composing of *Foeniculi Fructus*, *Isorae Fructus*, *Caryophylli folium*, *Zingiberis Rhizoma*, *Menthae arvensis Herba*, *Phyllanthi Herba*, *Mel depuratum*, *Myristicace semen*, *Amomi Fructus*, *Centellae Herba*, *Parkiae semen*, and *Oryza sativa*. Proximate analysis showed that the herbal medicine waste contained (as a dry basis) 2,189 kcal/kg GE, 11.6% CP, 3.28% CF, 40.4% crude fiber, and 4.89% total ash.

Experimental design, broiler management, and analyses: Three hundreds of day-old chicks [body weight (BW) 41.13 ± 0.22 g; mean \pm SD] Lohmann meat broilers were placed in an open-sided ventilated broiler house with rice husk as bedding material. They were randomly allocated to five dietary groups, i.e., control diet (basal diet with no additive; CONT), basal diet containing 0.04% of AZB, basal diet containing 0.01% of commercial probiotic *B. subtilis* (PROB), basal diet containing 1% of *C. crassa*-fermented rice bran (PROC), and basal diet with 1% of herbal medicine waste (HERBW). The basal diet was prepared as a single feed throughout the study period (34 days) and formulated to comply the Indonesian National Standards for broiler feed requirement (Table 1).

The feeds were prepared in mash form and provided *ad libitum* throughout the trial. Coccidiostat was not included in the basal feed. The feed additives were appended "on top" after feed formulation. The amount of *C. crassa*-fermented rice bran added to the basal feed was based on Sugiharto et al. (2017a) while the other commercial feed additives were according to the manufacturer's instructions. Vaccinations were conducted on chicks using commercial Newcastle disease virus (NDV) vaccine at days 4 and 18 through eye drop and drinking water, respectively. BW and feed consumption were recorded weekly.

On day 34, blood collection was conducted on one bird per pen (six birds per dietary group) from the birds' wing veins. For the analysis of complete blood counts, the blood was put in ethylenediaminetetraacetic acid vacutainers. For the analysis of serum biochemistry, blood was put in vacutainers with no anticoagulant. To make the serum, the blood was permitted to clot for 2 hours at room temperature and then, centrifuged at $5,000 \times g$ for 10 min. The same birds as blood sampled were killed by *halal* neck cut after being weighed (Alshelmani et al., 2016). The internal organs and commercial cuts of broilers were obtained and weighed. For the analyses of microbiology and short-chain fatty acids (SCFAs), the contents of ileum and cecum were aseptically collected.

Table 1. Ingredients and chemical composition (as a dry basis) of the basal diet

Items	Composition (% , unless otherwise noted)
Maize	45.5
Soybean meal	17.0
Wheat flour	10.0
Bread flour	5.00
Rice bran	4.45
Crude palm oil	3.50
Corn gluten meal	3.60
Distiller dried grains	3.00
Meat bone meal	2.80
Chicken feather meal	2.00
Bone meal	1.50
Lysine	0.55
Methionine	0.37
L-threonine	0.08
Salt	0.15
Premix ¹	0.50
Chemical composition:	
Metabolizable energy (kcal/kg) ²	3,300
DM	89.6
Crude protein	21.9
Crude fat	6.40
Crude fiber	5.62
Ash	6.39

¹Mineral-vitamin premix provided (per kg of feed) Ca 2.250 gm, P 0.625 gm, Fe 3.570 mg, Cu 0.640 mg, Mn 5.285 mg, Zn 0.003 mg, Co 0.001 mg, Se 0.013 mg, I 0.016 mg, vit A 375 IU, vit D 150 IU, vit E 0.080 mg

²Values were obtained based on the formula according to Bolton (1967), in which metabolizable energy = $40.81 [0.87 (CP + 2.25 CF + \text{nitrogen-free extract}) + 2.5]$

The complete blood counts were determined using a hematology analyzer (Prima Fully-auto Hematology Analyzer, PT. Prima Alkesindo Nusantara, Jakarta, Indonesia). The serum antibody titer to NDV vaccine was assayed on the basis of hemagglutination inhibition assay (Villegas, 1987). Total triglyceride was determined in serum according to the enzymatic method of glycerol-

3-phosphate oxidase. Cholesterol and its fractions were assayed based on the enzymatic determination of cholesterol oxidase. The enzymatic colorimetric analysis with glucose oxidase was employed to determine the blood glucose level. The levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) enzymes in serum were assayed according to the IFCC procedure (www.ifcc.org) with no pyridoxal phosphate. Photometric analyses were conducted to determine the levels of total protein and albumin in serum. The difference between total protein and albumin values was considered as globulin values. Uric acid concentration in serum was determined according to the enzymatic color test. The entire serum biochemical analyses were conducted in duplicate based on the manufacturer's protocols (DiaSys Diagnostic System GmbH, Holzheim, Germany).

The microbiological analysis of the ileal and cecal digesta was conducted according to [Sugiharto et al. \(2017a\)](#) with few modifications. The numbers of coliform and lactose-negative enterobacteria were enumerated on MacConkey agar (Merck KGaA). The inoculated cultures were incubated aerobically at 38°C for 24 h and then, counted as red and colorless colonies, respectively. The sum of coliform and lactose-negative enterobacteria was regarded

as enterobacteria. To enumerate the number of lactic acid bacteria (LAB), de Man, Rogosa and Sharpe (MRS; Merck KGaA) agar was used. Bacterial enumeration was conducted after the culture was incubated anaerobically at 38°C for 48 h. The population of *Clostridium perfringens* were enumerated on the tryptose sulphite cycloserine (TSC; Merck KGaA) agar. The black colonies were counted after the plates were incubated anaerobically at 38°C for 48 h. Gas chromatography was used to determine the SCFAs concentration in the cecal digesta of broilers. The analysis was performed according to [Sugiharto et al. \(2015a\)](#).

Results in the present study were analyzed by ANOVA using GLM Procedure (SAS Inst. Inc., Cary, NC, USA). The pen was considered as the unit of the experiment. Duncan's multiple-range test was carried out when significant differences among treatments exist.

RESULTS

Table 2 shows the data on final BW, feed intake, and feed conversion ratio (FCR) of broilers. Birds in PROB group had higher ($P < 0.05$) final BW than in CONT, but the significant difference was not observed compared to

Table 2. The effect of fungi-origin probiotic *C. crassa* and other feed additives on performances of broiler chickens

Items	Dietary treatments					SE	P value
	CONT	AGP	PROB	PROC	HERBW		
Final BW, gm/bird	1,620 ^b	1,663 ^{ab}	1,729 ^a	1,663 ^{ab}	1,691 ^{ab}	23.7	0.04
Feed intake, gm/bird	2,906	2,952	3,031	2,970	3,013	30.3	0.05
FCR	1.84	1.82	1.80	1.83	1.83	0.02	0.54

^{a,b}Means with different superscripts in each row are significantly different

CONT: basal diet, AGP: basal diet supplemented with zinc bacitracin, PROB: basal diet with *B. subtilis*, PROC: basal diet with *C. crassa*, HERBW: basal diet with herbal medicine waste, and SE: standard error

Table 3. The effect of fungi-origin probiotic *C. crassa* and other feed additives on internal organs of broiler chickens

Items (% live BW)	Dietary treatments					SE	P value
	CONT	AGP	PROB	PROC	HERBW		
Heart	0.42	0.44	0.43	0.44	0.45	0.03	0.96
Liver	2.16	2.34	2.21	2.14	2.14	0.13	0.78
Bile	0.06	0.05	0.04	0.05	0.05	0.01	0.61
Proventriculus	0.40	0.42	0.37	0.37	0.43	0.02	0.08
Gizzard	1.35 ^a	1.18 ^b	1.19 ^b	1.32 ^{ab}	1.38 ^a	0.05	0.03
Duodenum	0.60	0.59	0.55	0.56	0.59	0.04	0.94
Jejunum	1.12	0.99	0.99	1.05	1.17	0.07	0.24
Ileum	0.89	0.89	0.77	0.77	0.78	0.06	0.30
Caeca	0.30	0.26	0.28	0.28	0.35	0.03	0.25
Pancreas	0.25	0.27	0.25	0.26	0.28	0.02	0.81
Spleen	0.07	0.07	0.10	0.09	0.08	0.01	0.14
Thymus	0.28	0.28	0.29	0.36	0.28	0.04	0.59
Bursa of Fabricius	0.10	0.21	0.13	0.19	0.15	0.04	0.37

^{a,b}Means with different superscripts in each row are significantly different

CONT: basal diet, AGP: basal diet supplemented with zinc bacitracin, PROB: basal diet with *B. subtilis*, PROC: basal diet with *C. crassa*, HERBW: basal diet with herbal medicine waste, and SE: standard error

Table 4. The effect of fungi-origin probiotic *C. crassa* and other feed additives on the hematological parameter of broilers

Items	Dietary treatments					SE	P value
	CONT	AGP	PROB	PROC	HERBW		
Complete blood counts							
Hemoglobin (gm/dL)	9.63	9.93	10.3	9.75	9.80	0.35	0.76
Erythrocytes (10 ⁶ /μL)	2.23	2.31	2.37	2.26	2.23	0.08	0.72
Hematocrit (%)	30.8	31.3	32.2	30.6	30.7	1.18	0.86
MCV (fl)	138	136	136	135	138	1.36	0.51
MCH (pg)	43.3	43.0	43.3	43.1	44.0	0.54	0.71
MCHC (gm/dL)	31.4	31.7	31.9	31.9	31.9	0.36	0.85
Leukocytes (10 ³ /μL)	17.5 ^b	20.2 ^b	20.6 ^b	20.9 ^b	27.2 ^a	1.74	<0.01
Heterophils (10 ³ /μL)	0.53	0.45	0.62	0.62	0.63	0.10	0.62
Eosinophils (10 ³ /μL)	0.85	0.95	1.00	1.05	1.10	0.12	0.60
Lymphocytes (10 ³ /μL)	16.1 ^b	18.8 ^b	19.0 ^b	19.2 ^b	25.5 ^a	1.60	<0.01
Thrombocytes (10 ³ /μL)	16.7	17.5	17.7	18.3	17.3	1.68	0.97
Biochemical parameters							
Total cholesterol (mg/dL)	142	159	175	164	153	10.9	0.28
HDL (mg/dL)	32.3	27.7	21.7	27.7	31.0	3.14	0.17
LDL (mg/dL)	84.5	94.8	114	95.0	86.4	10.9	0.37
Total triglycende (mg/dL)	125	181	201	197	177	33.5	0.53
AST (U/l)	260 ^{ab}	249 ^b	247 ^b	317 ^a	209 ^b	21.4	0.03
ALT (U/l)	1.02	0.80	1.73	0.70	0.60	0.50	0.46
Total protein (gm/dL)	2.97 ^b	3.96 ^a	4.05 ^a	3.62 ^a	3.76 ^a	0.19	<0.01
Albumin (gm/dL)	1.38 ^b	1.47 ^{ab}	1.62 ^a	1.48 ^{ab}	1.49 ^{ab}	0.05	0.04
Globulin (gm/dL)	1.59 ^b	2.49 ^a	2.44 ^a	2.18 ^a	2.27 ^a	0.17	<0.01
A/G ratio	0.91 ^a	0.60 ^b	0.67 ^b	0.70 ^b	0.67 ^b	0.06	0.01
Uric acid (mg/dL)	13.0 ^{ab}	8.80 ^{bc}	14.2 ^a	7.14 ^c	8.23 ^{bc}	1.66	0.02
Creatinine (mg/dL)	0.39	0.35	0.37	0.27	0.44	0.05	0.29
Antibody titer against NDV (Log ₂ GMT)	4.67 ^b	5.33 ^{ab}	5.60 ^{ab}	6.00 ^a	4.67 ^b	0.35	0.04

^{ab}: Means with different superscripts in each row are significantly different

CONT: basal diet, AGP: basal diet supplemented with zinc bacitracin, PROB: basal diet with *B. subtilis*, PROC: basal diet with *C. crassa*, HERBW: basal diet with herbal medicine waste, MCV: mean corpuscular volume, MCH: Mean corpuscular hemoglobin, MCHC: mean corpuscular hemoglobin concentration, A/G ratio: albumin to globulin ratio, GMT: geometric mean titer, and SE: standard error

that in AGP, PROC, and HERBW birds. There was an obvious tendency ($P=0.05$) that birds in PROB and HERBW groups had higher feed intake than CONT birds. FCR did not differ ($P>0.05$) among the birds.

The weight of gizzard relative to BW was lower ($P<0.05$) in AGP and PROB compared to that in CONT and HERBW birds. No significant divergence was observed with regards to other internal organs among the birds (Table 3).

Data on the complete blood count of broiler chicks are shown in Table 4. The numbers of leukocytes and lymphocytes were greater ($P<0.05$) in HERBW than in other birds. There was no significant difference with regards to the numbers of heterophils, eosinophils, thrombocytes, erythrocytes, hemoglobin, and the percentage of hematocrit. Data on serum antibody titer and biochemical parameters are presented in Table 4. Compared to CONT and HERBW, PROC birds had

higher ($P<0.05$) level of vaccine titer to NDV. CONT had lower ($P<0.05$) and higher ($P<0.05$) total protein and globulin, and the ratio of albumin to globulin (A/G ratio) in serum, respectively, compared to other chicks. Higher level ($P<0.05$) of albumin was observed in PROB relative to CONT and PROC. Lower uric acid ($P<0.05$) was seen in PROC when compared with CONT and PROB. The AST enzyme was higher ($P<0.05$) in PROC than in AGP, PROB, and HERBW, but not different when compared with CONT birds. The differences in serum creatinine, ALT enzyme, triglycerides, total cholesterol, and low-density lipoprotein (LDL) and high-density lipoprotein (HDL) cholesterol were not observed among the birds.

Data on the numbers of microorganisms and pH of ileal and cecal digesta are shown in Table 5. Coliform bacteria were lower ($P<0.05$) in the ileal digesta of PROB and PROC than in CONT and HERBW birds, but the substantial difference was not observed in comparison to AGP. The numbers of *Clostridium perfringens* tended

Table 5. The effect of fungi-origin probiotic *C. crassa* and other feed additives on microbial populations and pH in the ileal and cecal digesta of broilers

Items (log cfu/gm, except pH)	Dietary treatments					SE	P value
	CONT	AGP	PROB	PROC	HERBW		
Ileum							
Coliform	6.68 ^a	6.21 ^{ab}	5.89 ^b	5.94 ^b	6.68 ^a	0.23	0.04
Lactose-negative enterobacteria	2.88	3.26	2.48	3.69	2.64	1.09	0.93
Enterobacteriaceae	9.56	9.47	8.37	9.63	9.32	0.95	0.88
LAB	8.50	8.45	8.32	8.18	8.15	0.25	0.82
<i>Clostridium perfringens</i>	5.94	3.88	3.46	3.97	5.94	0.77	0.07
pH	5.73	6.02	5.67	5.72	5.87	0.16	0.53
Cecum							
Coliform	6.39	6.18	5.95	6.12	6.64	0.18	0.10
Lactose-negative enterobacteria	5.13	4.46	5.43	5.25	5.00	0.45	0.62
Enterobacteriaceae	11.5	10.7	11.4	11.4	11.7	0.46	0.59
LAB	8.49	8.60	8.57	8.58	8.87	0.11	0.20
<i>Clostridium perfringens</i>	6.39	5.95	6.08	6.67	6.00	0.28	0.35
pH	6.30	6.18	6.17	6.35	6.58	0.16	0.38

^{a,b}Means with different superscripts in each row are significantly different

CONT: basal diet, AGP: basal diet supplemented with zinc bacitracin, PROB: basal diet with *B. subtilis*, PROC: basal diet with *C. crassa*, HERBW: basal diet with herbal medicine waste, cfu: colony forming units, SE: standard error

($P=0.07$) to be lower in the ileal digesta of AGP, PROB, and PROC than in that of CONT and HERBW birds. The significant difference was not observed in terms of pH both in the ileal and cecal digesta and microbial populations in the cecal digesta of broilers across the treatment groups.

The significant divergence was not seen with respect to the SCFAs concentrations in the cecal contents and carcass traits of broiler chicks (data not presented).

DISCUSSION

In the present study, commercial probiotic *B. subtilis* showed the most substantial effect on the growth performance of broilers, especially when compared to the control. This result has also been confirmed by other authors (Gao et al., 2017). The latter workers suggested that *B. subtilis* improved the digestibility and metabolic process, and hence increased the growth rate of broilers. Such commercial probiotic has also been documented to improve the intestinal health, and therefore more energy was partitioned into growth rather than maintenance (Gao et al., 2017). Indeed, there was no significant difference in term of final BW among PROB, AGP, PROC, and HERBW birds in this study. With regard particularly to *C. crassa*, the absence of difference in final BW between PROC and AGP birds may, therefore, suggest the opportunity of the fungal probiotic to substitute the use of AGPs in broiler production. With regard to feed intake, PROB and HERBW birds tended to consume more feed than CONT birds in this study. In accordance with our result, feeding *B. subtilis* has been

reported to increase feed intake in broiler chickens challenged with *Salmonella Gallinarum* (Park et al., 2017). The mechanism by which probiotic *B. subtilis* increased feed intake in broilers was not definitely known, but probiotics may improve the intestinal microbial ecosystem and increase digestive enzyme activity which aids the nutrient digestion (Sugiharto, 2016). The increased feed intake in HERBW birds was concomitant with Guo et al. (2004) reporting an increased feed intake in female broiler chicks fed Chinese herbal medicine. In such case, herbal preparations may improve the flavor of feeds, increase the secretion of digestive fluids, and improve the bacterial community in the gut of chicks (Wenk, 2003).

Results in the present study showed that dietary supplementation with AGPs or probiotic *B. subtilis* resulted in the lower relative weight of gizzard as compared to other treatment birds. Concomitantly, Barbieri et al. (2015) reported a decreased relative weight of gizzard in 21 days of age broiler chicks fed AGP savilamycin. With regard to probiotic treatment, our present finding was in contrast to most published papers reporting an increase (Khaliq and Ebrahimzad, 2016) or no change (Barbieri et al., 2015; Sugiharto et al., 2017a) in the relative weight of gizzard of broilers. At present, the reason for the lower relative weight of gizzard in AGP and PROC birds remains unknown. Perhaps, the relatively higher final BW in AGP and PROC birds (as the denominator in the calculation) resulted in lower values of the relative gizzard weight. Indeed, the significant difference was not observed with regard to the

absolute weight of gizzard across the birds in the present study (data not presented).

Compared to other birds, feeding herbal medicine waste resulted in higher numbers of leukocytes and lymphocytes. The latter data may be attributed to the enhanced immune responses in HERBW birds (Sugiharto et al., 2017b). Considering the lower serum antibody titer against NDV vaccine (and thus lower protection against NDV) in HERBW chicks, the increased numbers of leukocytes and lymphocytes may, therefore, be associated with the acute viral infection or internal stress (Sugiharto et al., 2017b) in the respective birds. In this current study, serum antibody titer against NDV vaccine was significantly higher in PROC than in CONT and HERBW birds. This may suggest that feeding probiotic *C. crassa* was capable of protecting the birds from potential infection by NDV. Corresponding report was revealed by Haghghi et al. (2005) showing an increased level of serum antibody (immunoglobulin M) against sheep red blood cells in broiler chicks treated with a probiotic mixture containing *Lactobacillus acidophilus*, *Bifidobacterium bifidum*, and *Streptococcus faecalis*. The latter authors suggested that probiotics may provoke the immune cells to secrete cytokines, which in turn stimulate the production of antibody.

Relative to CONT, total protein and globulin concentrations in serum increased in the treated chicks. It has widely been known that serum total protein may reflect protein deposition in the body and thus the growth rate of farm animals (Ghazalah and Ali, 2008). Consistent with this, relatively higher final BW was observed in the treated birds as compared to CONT birds. With regard to globulin, the high level of this protein fraction has been associated with the improved immune defense system in broilers (Abdel-Fattah et al., 2008). Hence, the higher concentration of serum globulin may implicate in better disease protection in the treated birds. The latter suggestion was also endorsed by the data that albumin to globulin ratio (A/G ratio) was lower in the treated birds than in the control birds. Note that lower serum A/G ratio was attributed to the better immune competence of broilers (Ghazalah and Ali, 2008; Sugiharto et al., 2016). In the current trial, albumin concentration was higher in the serum of PROB as compared to CONT birds. Given that serum albumin level was positively correlated with growth rate and nitrogen retention in animals (de Ridder et al., 2012), the higher albumin concentration in PROB may, therefore, promote protein deposition and thus growth rate in the respective broilers. In poultry, serum uric acid concentration has been used to indicate muscle

proteolysis (Sugiharto et al., 2017a), infections (Lin et al., 2015), and stress conditions (Babacanoglu et al., 2013). In this study, serum uric acid was lowest in birds supplemented with probiotic *C. crassa*. This finding may, therefore, suggest that the fungus *C. crassa* was capable of protecting the birds from the above detrimental conditions. In this study, *C. crassa*-supplemented birds had a higher level of AST enzyme as compared to other treated birds. This may suggest that feeding *C. crassa* led to the toxicological effect on birds (Sugiharto et al., 2016). It was possible that phytic acid content in rice bran (as a carrier/media for growing the fungus) negatively affected the liver condition (Shibata et al., 2012), resulting in increased level of AST enzyme. This notion should, however, be accounted with caution since the level of ALT enzyme and the liver relative weight were not different among the birds in this current study.

It has previously been revealed that feeding fungal probiotic (e.g., *Acremonium charticola*) (Sugiharto et al., 2017b) and probiotic *B. subtilis* (Jeong and Kim, 2014) was able to reduce the number of coliform in the gastrointestinal tract of broiler chicks. In agreement with this, our present data demonstrated the lower number of coliform bacteria in the ileal content of PROB and PROC as compared to CONT and HERBW birds. In the current study, fungal and bacterial probiotic supplementation was also able to control the population of *C. perfringens* in the ileum of broilers, which was comparable to the effect of AGPs supplementation. The concomitant result has been confirmed by La Ragione and Woodward (2003), in which feeding *B. subtilis* spores reduced the intestinal colonization by *C. perfringens* in broiler chicks. The mechanisms by which the fungal and bacterial probiotics controlled the population of pathogenic bacteria in the intestine of birds are largely unknown. In respect to probiotic *B. subtilis*, this probiotic may, however, decrease the potentially pathogenic bacteria through competitive exclusion (La Ragione and Woodward, 2003). With regard particularly to the fungus *C. crassa*, our present finding was consistent with Yudiarti et al. (2012b) who previously showed the potential of the fungus in controlling the bacterial population (not specific to a certain bacteria) in the intestine of the Indonesian native chickens. According to Sugiharto et al. (2015b; 2017a), one possible mechanism by which the fungal probiotic controls the population of pathogenic microorganism could be that the fungus produced some antibacterial substances that disrupt the biological functions of pathogens. Taken together, dietary inclusion with the fungus *C. crassa* was beneficial in improving the intestinal microbial population of broiler chickens.

In the present study, the influence of treatments was not observed with regard to the concentrations of SCFAs in the cecal digesta of broilers. With regard particularly to probiotic and AZB, [Olnood et al. \(2015\)](#) previously reported no notable effect of probiotic *Lactobacillus* spp. and zinc bacitracin on SCFAs (acetate, propionate, and butyrate) concentration in the contents of ileum and cecum of 21 and 35 days of age broilers. Unlike this, [Milián et al. \(2013\)](#) reported the increased concentrations of acetate, propionate, and total SCFAs in broiler chicks with feeding *B. subtilis*. Moreover, [Murugesan et al. \(2014\)](#) showed a higher concentration of butyrate in broilers fed probiotic *Bacillus licheniformis* strain while the concentrations of acetate, propionate, and total SCFAs did not change with the treatment. The definite reason for these divergent findings remains unclear, but the different species and strains of probiotic microorganisms used as well as the conditions of the study may be responsible. In respect to the effect of herbal medicine waste, corresponding result was reported in pig by [Hanczakoska and Swiatkiewicz \(2012\)](#), at which herbal extract had no impact on the concentrations of acetate, propionate, butyrate, and total SCFAs in the ileum and cecum ([Hanczakoska and Swiatkiewicz, 2012](#)).

⁵⁶ The lack of influence of dietary treatments on the carcass traits of broilers was observed in the present study. This result was consistent with that of previously reported by [Sugiharto et al. \(2017a\)](#) showing no effects of the probiotic *C. crassa* on carcass characteristics of broiler chickens. With regard to AGPs and probiotic *B. subtilis*, treatment with such additives has also been reported not to affect the carcass traits of broiler chickens ([Nunes et al., 2012](#)). In term of treatment with herbal medicine waste, our present data were in accordance with that of reported by [Onu \(2010\)](#) showing no effect of two herbal spices (garlic and ginger) on carcass traits of broilers. However, [Puvača et al. \(2016\)](#) reported conversely, in that, feeding herbal drugs (garlic, black pepper, and hot red pepper) increased carcass weight in broilers. The definite explanation for these divergent results was unknown, but the difference in the composition and nature of herbal products used seemed to be responsible.

Overall, the effect of probiotic *C. crassa* on broiler growth and health performances corresponded to that of commercial probiotic *B. subtilis* and AGPs. On this basis, probiotic *C. crassa* can be a new candidate for AGPs substitute in broiler rations. However, the use of probiotic *C. crassa* as a feed additive for broilers seemed to be less economic especially when compared with probiotic *B. subtilis*. Indeed, the study showed that in

general, filamentous fungi had a slow growth rate in the culture media, and therefore need longer time to proliferate (in our case, *C. crassa* was grown in rice bran for 4 days). Moreover, the longer incubation time (for fungal growth) may imply in more accumulation of contaminants in the fungal culture ([Miyashira et al., 2010](#)). With regard to *B. subtilis*, these bacteria need a shorter time to proliferate, which is about 24 h ([Pant et al., 2015](#)).

CONCLUSION

In conclusion, treatment with probiotics was capable of improving the growth performance, immune responses, and intestinal bacterial populations of broiler chicks. The effects of fungi-origin probiotic *C. crassa* corresponded to that of commercial probiotic *B. subtilis* and AGPs.

ACKNOWLEDGEMENT

We would like to thank PT. Bayer Indonesia (Jakarta, Indonesia) and PT. Sidomuncul, Tbk. (Semarang, Indonesia), which provided commercial probiotic *B. subtilis* (Baymix Grobig) and herbal medicine waste, respectively. The study was fully funded by Diponegoro University through non-tax state revenue (PNBP), no. 275-046/UN7.5.1/PG/2017, 23 March 2017.

CONFLICT OF INTEREST

The authors have no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

AUTHORS' CONTRIBUTION

SS planned, conducted the experiment, and prepared the article. TY, EW, HIW, and TAS carried out the *in vivo* trial and corrected the article and II conducted the analysis of statistic.

REFERENCES

1. Abdel-Fattah SA, El-Sanhoury MH, El-Mednay NM, Abdel-Azeem F. Thyroid activity, some blood constituents, organs morphology and performance of broiler chicks fed supplemental organic acids. International Journal of Poultry Science. 2008; 7:215–222. <https://doi.org/10.3923/ijps.2008.215.222>
2. Abid SA, Azeem T, Chaudhary ZI, Ahmad M, Rehman ZU, Umar S. Emerging threat of necrotic enteritis in poultry and its control without use of antibiotics: a review. Journal of Animal and Plant Science. 2016; 26:1556–1667.

3. Abudabos AM, Alyemni AH, Dafalla YM, Khan RU. Effect of organic acid blend and *Bacillus subtilis* alone or in combination on growth traits, blood biochemical and antioxidant status in broiler exposed to *Salmonella typhimurium* challenge during the starter phase. *Journal of Applied Animal Research*. 2017b; 45:538–542. <https://doi.org/10.1080/09712119.2016.1219665>
4. Abudabos AM, Alyemni AH, Dafalla YM, Khan RU. The effect of phytochemicals on growth traits, blood biochemical and intestinal histology in broiler chickens exposed to *Clostridium perfringens* challenge. *Journal of Applied Animal Research*. 2017a; 46:691–695. <https://doi.org/10.1080/09712119.2017.1383258>
5. Alshelmani MI, Loh TC, Foo HL, Sazili AQ, Lau WH. Effect of feeding different levels of palm kernel cake fermented by *Paenibacillus polymyxa* ATCC 842 on nutrient digestibility, intestinal morphology, and gut microflora in broiler chickens. *Animal Feed Science and Technology*. 2016; 216:216–224. <https://doi.org/10.1016/j.anifeedsci.2016.03.019>
6. AOAC. Official methods of analysis of the association of analytical chemists. Washington DC, USA. 1995.
7. Babacanoglu E, Yalçın S, Uysal S. Evaluation of a stress model induced by dietary corticosterone supplementation in broiler breeders: effects on egg yolk corticosterone concentration and biochemical blood parameters. *British Poultry Science*. 2013; 54(6):677–685. <https://doi.org/10.1080/00071668.2013.847901>
8. Barbieri A, do Valle Polycarpo G, Cardoso RGA, da Silva KM, Dadalt JC, Madeira AMBN, de Sousa RLM, de Albuquerque R, Cruz-Polycarpo VC. Effect of probiotic and organic acids in an attempt to replace the antibiotics in diets of broiler chickens challenged with *Eimeria* spp. *International Journal of Poultry Science*. 2015; 14:606–614. <https://doi.org/10.3923/ijps.2015.606.614>
9. Bolton W. Poultry nutrition. MAFF Bulletin No.174, HMSO, London. 1967
10. de Ridder K, Levesque CL, Htoo JK, de Lange CFM. Immune system stimulation reduces the efficiency of tryptophan utilization for body protein deposition in growing pigs. *Journal of Animal Science*. 2012; 90:3485–3491. <https://doi.org/10.2527/jas.2011-4830>
11. Elfahmi, Woerdenbag HJ, Kayser O. *Jamur*. Indonesian traditional herbal medicine towards rational phytopharmacological use. *Journal of Herbal Medicine*. 2014; 4:51–73. <https://doi.org/10.1016/j.hermed.2014.01.002>
12. Feng Y, Yang XJ, Wang YB, Li WL, Liu Y, Yin RQ, Yao JH. Effects of immune stress on performance parameters, intestinal enzyme activity and mRNA expression of intestinal transporters in broiler chickens. *Asian-Australasian Journal of Animal Science*. 2012; 25:701–707. <https://doi.org/10.5713/ajas.2011.11377>
13. Gao Z, Wu H, Shi L, Zhang X, Sheng R, Yin F, Gooneratne R. Study of *Bacillus subtilis* on growth performance, nutrition metabolism and intestinal microflora of 1 to 42 d broiler chickens. *Animal Nutrition*. 2017; 3:109–113. <https://doi.org/10.1016/j.aninu.2017.02.002>
14. Ghazalah AA, Ali AM. Rosemary leaves as a dietary supplement for growth in broiler chickens. *International Journal of Poultry Science*. 2008; 7:234–239. <https://doi.org/10.3923/ijps.2008.234.239>
15. Guo FC, Kwakkel RP, Soede J, Williams BA, Verstegen MW. Effect of a Chinese herb medicine formulation, as an alternative for antibiotics, on performance of broilers. *British Poultry Science*. 2004; 45:793–797. <https://doi.org/10.1080/00071660400012741>
16. Haghighi HR, Gong J, Gyles CL, Hayes MA, Sanei B, Parvizi P, Gisavi H, Chambers JR, Sharif S. Modulation of antibody-mediated immune response by probiotics in chickens. *Clinical and Diagnostic Laboratory Immunology*. 2005; 12:1387–1392. <https://doi.org/10.1128/CDLI.12.12.1387-1392.2005>
17. Han X, Piao XS, Zhang HY, Li PF, Yi JQ, Zhang Q, Li P. *Forsythia suspensa* extract has the potential to substitute antibiotic in broiler chicken. *Asian-Australasian Journal of Animal Science*. 2012; 25:569–576. <https://doi.org/10.5713/ajas.2011.11425>
18. Hanczakoska E, Swiatkiewicz M. Effect of herbal extracts on piglet performance and small intestinal epithelial villi. *Czech Journal of Animal Science*. 2012; 57:420–429. <https://doi.org/10.17221/6316-CJAS>
19. Jeong JS, Kim IH. Effect of *Bacillus subtilis* C-3102 spores as a probiotic feed supplement on growth performance, noxious gas emission, and intestinal microflora in broilers. *Poultry Science*. 2014; 93:3097–3103. <https://doi.org/10.3382/ps.2014-04086>
20. Khaliq M, Ebrahimnezhad Y. The effects of probiotic Yeasture® using at different rearing periods on performance, intestinal microbial population and carcass traits in broiler chickens. *Veterinary Science Development*. 2016; 6.
21. Kisworo AN, Agus A, Kustantinah, Suwignyo B. Physicochemical characteristics identification and secondary metabolite analysis of solid herbal waste as source of feed rich fiber and supplement for ruminants. *Animal Production*. 2016; 18:75–84. <https://doi.org/10.20884/1.anprod.2016.18.2.535>
22. La Ragione RM, Woodward MJ. Competitive exclusion by *Bacillus subtilis* spores of *Salmonella enterica* serotype Enteritidis and *Clostridium perfringens* in young chickens. *Veterinary Microbiology*. 2003; 94:245–256. [https://doi.org/10.1016/S0378-1135\(03\)00077-4](https://doi.org/10.1016/S0378-1135(03)00077-4)
23. Lin H, Huang Q, Guo X, Liu P, Liu W, Zou Y, Zhu S, Deng G, Kuang J, Zhang C, Cao H, Hu G. Elevated level of renal xanthine oxidase mRNA transcription after nephropathogenic infectious bronchitis virus infection in growing layers. *Journal of Veterinary Science*. 2015; 16:423–429. <https://doi.org/10.4142/jvs.2015.16.4.423>
24. Milián G, Rondón AJ, Pérez M, Bocourt R, Rodríguez Z, Ranilla MJ, Rodríguez M, Carro MD. Evaluation of *Bacillus*

- subtilis* biopreparations as growth promoters in chickens. Cuban Journal of Agricultural Science. 2013; 47(1):61–66.
25. Miyashita CH, Tanigushi DG, Gugliotta AM, Santos DYAC. Comparison of radial growth rate of the mutualistic fungus of *Atta sexdens rubropilosa* forel in two culture media. Brazilian Journal of Microbiology. 2010; 41:506–511. <https://doi.org/10.1590/S1517-83822010000200035>
 26. Murugesan GR, Romero LF, Persia ME. Effects of protease, phytase and a *Bacillus* sp. direct-fed microbial on nutrient and energy digestibility, ileal brush border digestive enzyme activity and cecal short-chain fatty acid concentration in broiler chickens. PLoS ONE. 2014; 9:e101888. <https://doi.org/10.1371/journal.pone.0101888>
 27. Nunes JO, Bertechini AG, de Brito JÁG, Fassani ÉJ, Mesquita FR, Makiyama L, Meneghetti C. Evaluation of the use of probiotic (*Bacillus subtilis* C-3102) as additive to improve performance in broiler chicken diets. Revista Brasileira de Zootecnia. 2012; 41:2374–2378. <https://doi.org/10.1590/S1516-35982012001100012>
 28. Olnood CG, Beski SSM, Choct M, Iji PA. Novel probiotics: their effects on growth performance, gut development, microbial community and activity of broiler chickens. Animal Nutrition. 2015; 1:184–191. <https://doi.org/10.1016/j.aninu.2015.07.003>
 29. Onu PN. Evaluation of two herbal spices as feed additives for finisher broilers. Biotechnology in Animal Husbandry 2010; 26:383–392. <https://doi.org/10.2298/BAH1006383O>
 30. Pant G, Prakasha A, Pavana JVP, Bera S, Deviram GVNS, Kumar A, Panchpuri M, Prasunaa RG. Production, optimization and partial purification of protease from *Bacillus subtilis*. Journal of Taibah University for Science. 2015; 9:50–55. <https://doi.org/10.1016/j.jtusci.2014.04.010>
 31. Park JH, Kim YM, Kang DK, Kim IH. Effect of dietary *Bacillus subtilis* C14 and RX7 strains on growth performance, blood parameter, and intestinal microbiota in broiler chickens challenged with *Salmonella gallinarum*. Journal of Poultry Science. 2017; 54:236–241. <https://doi.org/10.2141/jpsa.0160078>
 32. Puvača NM, Kostadinović LM, Đuragić OM, Ljubojević DB, Mišević BM, Könyves TL, Popović SJ, Lević, Nikolova NB. Influence of herbal drugs in broiler chicken nutrition on primal carcass cuts quality assessments. Food Feed Research. 2016; 43:43–50. <https://doi.org/10.5937/FFR1601043P>
 33. Schokker D, Jansman AJM, Veninga G, de Bruin N, Vastenhouw SA, de Bree FM, Bossers A, Rebel JMJ, Smits MA. Perturbation of microbiota in one-day old broiler chickens with antibiotic for 24 hours negatively affects intestinal immune development. BMC Genomics. 2017; 18:241. <https://doi.org/10.1186/s12864-017-3625-6>
 34. Shibata T, Yoneda K, Araki T, Nikki T. Effect of phytic acid dietary level on growth performance and serum components in broiler chickens. Journal of Poultry Science. 2012; 49:111–115. <https://doi.org/10.2141/jpsa.011112>
 35. Simon K, Verwoolde MB, Zhang J, Smidt H, de Vries Reilingh G, Kemp B, Lammers A. Long-term effects of early life microbiota disturbance on adaptive immunity in laying hens. Poultry Science. 2016; 95:1543–1554. <https://doi.org/10.3382/ps/pew088>
 36. Soetrisnanto D, Christwardana M, Hadiyanto. Application of phytoremediation for herbal medicine waste and its utilization for protein production. Reaktor. 2012; 14:129–134. <https://doi.org/10.14710/reaktor.14.2.129-134>
 37. Sugiharto S, Lauridsen C, Jensen BB. Gastrointestinal ecosystem and immunological responses in *E. coli* challenged pigs after weaning fed liquid diets containing whey permeate fermented with different lactic acid bacteria. Animal Feed Science and Technology. 2015a; 207:278–282. <https://doi.org/10.1016/j.anifeedsci.2015.06.019>
 38. Sugiharto S, Yudiarti T, Isroli I, Widiastuti E, Putra FD. Effect of dietary supplementation with *Rhizopus oryzae* or *Chrysonilia crassa* on growth performance, blood profile, intestinal microbial population, and carcass traits in broilers exposed to heat stress. Archives Animal Breeding. 2017a; 60:347–356. <https://doi.org/10.5194/aab-60-347-2017>
 39. Sugiharto S, Yudiarti T, Isroli I, Widiastuti E, Putra FD. Intestinal microbial ecology and hematological parameters of broiler fed cassava waste pulp fermented with *Acremonium charticola*. Veterinary World. 2017b; 10:324–330. <https://doi.org/10.14202/vetworld.2017.324-330>
 40. Sugiharto S, Yudiarti T, Isroli I. Functional properties of filamentous fungi isolated from the Indonesian fermented dried cassava, with particular application on poultry. Mycobiology. 2015b; 43:415–422. <https://doi.org/10.5941/MYCO.2015.43.4.415>
 41. Sugiharto S, Yudiarti T, Isroli I. Performances and haematological profile of broilers fed fermented dried cassava (*Manihotesculenta crants*). Tropical Animal Health and Production. 2016; 48:1337–1341. <https://doi.org/10.1007/s11250-016-1098-2>
 42. Sugiharto S. Role of nutraceuticals in gut health and growth performance of poultry. Journal of the Saudi Society of Agricultural Sciences. 2016; 15:99–111. <https://doi.org/10.1016/j.jssas.2014.06.001>
 43. Villegas P. Avian virus diseases laboratory manual. College of Veterinary Medicine. University of Georgia, Athens, Georgia, USA. 1987.
 44. Wenk C. Herbs and botanicals as feed additives in monogastric animals. Asian-Australasian Journal of Animal Science. 2003; 16:282–289. <https://doi.org/10.5713/ajas.2003.282>
 45. Yudiarti T, Yunianto VDBI, Isroli I, Murwani R. *In vitro* test of probiotic strain of seven species of fungi which isolated from gastrointestinal tract of indigenous chicken. Proceeding of the 6th International Seminar of Indonesian Society for Microbiology and the 11th Congress of

- Indonesian Society for Microbiology 2013 (ISISM-ISMCong), Pontianak, West Borneo, Indonesia. 2013.
46. Yudiarti T, Yuniarto VDBI, Murwani R, Kusdiyantini E. Isolation of fungi from the gastrointestinal tract of indigenous chicken. *Journal of the Indonesian Tropical Animal Agriculture*. 2012a; 37:115–120.
<https://doi.org/10.14710/jitaa.37.2.115-120>
47. Yudiarti T, Yuniarto VDBI, Murwani R, Kusdiyantini E. The Effect of *Chrysonilia crassa* additive on duodenal & caecal morphology, bacterial & fungal number, and productivity of Ayam Kampung. *International Journal of Science and Engineering*. 2012b; 3:26–29.
<https://doi.org/10.12777/ijse.3.2.26-29>
48. Zawqari MH, Al-Baddany AA, Al-Baadani HH, Alhidary IA, Khan RU, Aqil GM, Abdurab A. Effect of feeding dried sweet orange (*Citrus sinensis*) peel and lemon grass (*Cymbopogon citratus*) leaves on growth performance, carcass traits, serum metabolites and antioxidant status in broiler during the finisher phase. *Environmental Science and Pollution Research*. 2016; 23:17077–17082.
<https://doi.org/10.1007/s11356-016-6879-7>

This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 Licence (<http://creativecommons.org/licenses/by/4.0>)

ORIGINALITY REPORT

14%

SIMILARITY INDEX

8%

INTERNET SOURCES

12%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Mahmud, Muhammad, Peter Shaba, Helen Yisa, James Gana, Usman Zubairu, Ruth Ndagimba, and Silas Ndagi. "Comparative efficacy of Diazepam, Ketamine and Diazepam-Ketamine combination for sedation or anesthesia in cockerel chickens", Journal of Advanced Veterinary and Animal Research, 2014.
Publication 1%

- 2** João H. Reis, Roger R. Gebert, Mauricio Barreta, Matheus D. Baldissera et al. "Effects of phytogenic feed additive based on thymol, carvacrol and cinnamic aldehyde on body weight, blood parameters and environmental bacteria in broilers chickens", Microbial Pathogenesis, 2018
Publication 1%

- 3** Daneshmand, A.. "Effect of oyster mushroom (Pleurotus ostreatus) with and without probiotic on growth performance and some blood 1%

parameters of male broilers", Animal Feed Science and Technology, 20111125

Publication

4	Submitted to Mansoura University Student Paper	1%
5	Endang Kusumanti, Sugiharto Sugiharto. "Effect of dietary supplementation of binahong leaf meal, betel nut meal or their combination on serum albumin and globulin, fecal endoparasites and bacterial counts in milk of Saanen goats suffering from subclinical mastitis", Agriculture and Natural Resources, 2017 Publication	1%
6	ejournal.undip.ac.id Internet Source	1%
7	bdvets.org Internet Source	<1%
8	dspace.stir.ac.uk Internet Source	<1%
9	Chen G. Olnood, Sleman S.M. Beski, Mingan Choct, Paul A. Iji. "Novel probiotics: Their effects on growth performance, gut development, microbial community and activity of broiler chickens", Animal Nutrition, 2015 Publication	<1%

10

www.jofamericanscience.org

Internet Source

<1%

11

www.ajfand.net

Internet Source

<1%

12

Lene Lind Mikkelsen, Mogens Jakobsen, Bent Borg Jensen. "Effects of dietary oligosaccharides on microbial diversity and fructo-oligosaccharide degrading bacteria in faeces of piglets post-weaning", *Animal Feed Science and Technology*, 2003

Publication

<1%

13

"Direct-Fed Microbials and Prebiotics for Animals", Springer Nature America, Inc, 2012

Publication

<1%

14

www.i-scholar.in

Internet Source

<1%

15

theses.gla.ac.uk

Internet Source

<1%

16

Zeng, Zhikai, Xiao Xu, Qiang Zhang, Ping Li, Panfeng Zhao, Qingyun Li, Jundi Liu, and XiangShu Piao. "Effects of essential oil supplementation of a low-energy diet on performance, intestinal morphology and microflora, immune properties and antioxidant activities in weaned pigs : Essential oil for weaned pigs", *Animal Science Journal*, 2014.

<1%

17 media.neliti.com <1 %
Internet Source

18 Sugiharto, Sugiharto, Turrini Yudiarti, and Isroli Isroli. "Assay of Antioxidant Potential of Two Filamentous Fungi Isolated from the Indonesian Fermented Dried Cassava", *Antioxidants*, 2016.
Publication

19 Molist, F.. "Effects of the insoluble and soluble dietary fibre on the physicochemical properties of digesta and the microbial activity in early weaned piglets", *Animal Feed Science and Technology*, 20090316
Publication

20 www.jove.com <1 %
Internet Source

21 Choubey, Mahipal, Ashok Kumar Pattanaik, Shalini Baliyan, Narayan Dutta, Sunil E. Jadhav, and Kusumakar Sharma. "Dietary supplementation of a novel phytogetic feed additive: effects on nutrient metabolism, antioxidant status and immune response of goats", *Animal Production Science*, 2015.
Publication

22 www.blueberry.org <1 %
Internet Source

23

Mehdi Qorbanpour, Taha Fahim, Faramin Javandel, Mehran Nosrati et al. "Effect of Dietary Ginger (*Zingiber officinale* Roscoe) and Multi-Strain Probiotic on Growth and Carcass Traits, Blood Biochemistry, Immune Responses and Intestinal Microflora in Broiler Chickens", *Animals*, 2018

Publication

<1%

24

www.omicsonline.org

Internet Source

<1%

25

Ruixia Lan, Hoainam Tran, Inho Kim. "Effects of probiotic supplementation in different nutrient density diets on growth performance, nutrient digestibility, blood profiles, fecal microflora and noxious gas emission in weaning pig", *Journal of the Science of Food and Agriculture*, 2017

Publication

<1%

26

Sugiharto Sugiharto, Turrini Yudiarti, Isroli Isroli. "Functional Properties of Filamentous Fungi Isolated from the Indonesian Fermented Dried Cassava, with Particular Application on Poultry", *Mycobiology*, 2015

Publication

<1%

27

"Abstracts of 52nd EASD Annual Meeting", *Diabetologia*, 2016

Publication

<1%

28	<p>Marta Kupryś-Caruk, Monika Michalczuk, Beata Chabłowska, Ilona Stefańska, Danuta Kotyrba, Marta Parzeniecka-Jaworska. "Efficacy and safety assessment of microbiological feed additive for chicken broilers in tolerance studies", Journal of Veterinary Research, 2018</p> <p>Publication</p>	<1%
29	<p>scholar.sun.ac.za</p> <p>Internet Source</p>	<1%
30	<p>kb.psu.ac.th</p> <p>Internet Source</p>	<1%
31	<p>sasas.co.za</p> <p>Internet Source</p>	<1%
32	<p>doiserbia.nb.rs</p> <p>Internet Source</p>	<1%
33	<p>www.pagepress.org</p> <p>Internet Source</p>	<1%
34	<p>Naila Chand, Shabana Naz, Ziaur Rehman, Rifat Ullah Khan. "Blood biochemical profile of four fast-growing broiler strains under high ambient temperature", Applied Biological Chemistry, 2018</p> <p>Publication</p>	<1%
35	<p>Zhang, H. Y., X. S. Piao, Q. Zhang, P. Li, J. Q.</p>	<1%

Yi, J. D. Liu, Q. Y. Li, and G. Q. Wang. "The effects of Forsythia suspensa extract and berberine on growth performance, immunity, antioxidant activities, and intestinal microbiota in broilers under high stocking density", Poultry Science, 2013.

Publication

36

MOHAMMADREZA POURAKBARI, ALIREZA SEIDAVI, LEILA ASADPOUR, ANDRÉS MARTÍNEZ. "Probiotic level effects on growth performance, carcass traits, blood parameters, cecal microbiota, and immune response of broilers", Anais da Academia Brasileira de Ciências, 2016

Publication

<1%

37

preview-
biotechnologyforbiofuels.biomedcentral.com

Internet Source

<1%

38

orbi.uliege.be

Internet Source

<1%

39

www.researchgate.net

Internet Source

<1%

40

Steven C. Ricke. "Gastrointestinal Ecology of Salmonella Enteritidis in Laying Hens and Intervention by Prebiotic and Nondigestible Carbohydrate Dietary Supplementation", Elsevier BV, 2017

<1%

41 Sugiharto Sugiharto. "Role of nutraceuticals in gut health and growth performance of poultry", *Journal of the Saudi Society of Agricultural Sciences*, 2016
Publication <1%

42 hal-anses.archives-ouvertes.fr
Internet Source <1%

43 edepot.wur.nl
Internet Source <1%

44 www.ifa-tulln.boku.ac.at
Internet Source <1%

45 www.scilit.net
Internet Source <1%

46 eurasianjvetsci.org
Internet Source <1%

47 www.epa.hu
Internet Source <1%

48 tatzk.berkeley.edu
Internet Source <1%

49 Takahisa Gono. "VAD with or without subsequent high-dose melphalan followed by autologous stem cell support in AL amyloidosis: Japanese experience and criteria for patient selection", *Amyloid The Journal of* <1%

Protein Folding Disorders, 12/1/2004

Publication

50

dikikomarudin8.blogspot.com

Internet Source

<1%

51

lib.dr.iastate.edu

Internet Source

<1%

52

spasb.ro

Internet Source

<1%

53

Jannigje G. Kers, Francisca C. Velkers, Egil A. J. Fischer, Gerben D. A. Hermes, J. A. Stegeman, Hauke Smidt. "Host and Environmental Factors Affecting the Intestinal Microbiota in Chickens", *Frontiers in Microbiology*, 2018

Publication

<1%

54

Vázquez, Ramón Silva, Lorenzo Antonio Durán Meléndez, Eduardo Santellano Estrada, Carlos Rodríguez Muela, Guillermo Villalobos Villalobos, Gerardo Méndez Zamora, and Michael E. Hume. "Performance of broiler chickens supplemented with Mexican oregano oil (*Lippia berlandieri* Schauer)", *Revista Brasileira de Zootecnia*, 2015.

Publication

<1%

55

Dahiya, J.P.. "Potential strategies for controlling necrotic enteritis in broiler chickens in post-

<1%

antibiotic era", Animal Feed Science and
Technology, 20060804

Publication

56

Mohsen Soltani, Sayed Ali Tabeidian,
Gholamreza Ghalamkari, Amir Hossein Adeljoo
et al. "Effect of dietary extract and dried areal
parts of Rosmarinus officinalis on
performance, immune responses and total
serum antioxidant activity in broiler chicks",
Asian Pacific Journal of Tropical Disease, 2016

<1%

Publication

57

Miao Yu, Chunlong Mu, Chuanjian Zhang,
Yuxiang Yang, Yong Su, Weiyun Zhu. "Marked
Response in Microbial Community and
Metabolism in the Ileum and Cecum of
Suckling Piglets After Early Antibiotics
Exposure", Frontiers in Microbiology, 2018

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

vanced_Veterinary_and_Animal_Research_-_5_3_332-342.Agust_18.pdf

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
