

1 **Ar-Ar age constraints on the timing of Havre Trough opening and**
2 **magmatism.**

3

4 Richard Wysoczanski^{a*}, Graham Leonard^b, James Gill^c, Ian Wright^d, Andrew
5 Calvert^e, William McIntosh^f, Brian Jicha^g, John Gamble^{hi}, Christian Timm^{b,j},
6 Monica Handler^h, Elizabeth Drewes^k, and Alex Zohrab^h.

7

8 *^aNational Institute of Water & Atmospheric Research, Private Bag 14901, Wellington, New*
9 *Zealand*

10 *^bGNS Science, 1 Fairway Drive, Avalon, Lower Hutt, New Zealand*

11 *^cDepartment of Earth and Planetary Sciences, University of California, Santa Cruz,*
12 *California, USA*

13 *^dVice Chancellor's Office, University of Canterbury, Private Bag 4800, Christchurch 8141,*
14 *New Zealand*

15 *^eVolcano Science Center, US Geological Survey, Menlo Park, California, USA*

16 *^fNew Mexico Geochronology Research Laboratory, New Mexico Tech, Socorro, USA*

17 *^gDepartment of Geoscience, University of Wisconsin—Madison, 1215 West Dayton Street,*
18 *Madison WI 53706*

19 *^hSchool of Geography, Environment & Earth Science, Victoria University of Wellington, New*
20 *Zealand*

21 *ⁱSchool of Biological, Earth & Environment Sciences, University College Cork, Ireland*

22 *^jGEOMAR, Helmholtz Centre for Ocean Research, RD 4, Wischhofstrasse 1-3, 24148 Kiel,*
23 *Germany*

24 *^kAlaska Science Center, Alaska Science Center, U.S. Geological Survey, 4210 University*
25 *Drive, Anchorage, AK 99508, USA*

26

27 **Abstract**

28 The age and style of opening of the Havre Trough back-arc system is uncertain due to a lack
29 of geochronologic constraints for the region. ⁴⁰Ar/³⁹Ar dating of 19 volcanic rocks from
30 across the southern Havre Trough and Kermadec Arc was conducted in three laboratories to
31 provide age constraints on the system. The results are integrated and interpreted as suggesting
32 that this subduction system is young (< 2 Ma) and coeval with opening of the continental

33 Taupo Volcanic Zone of New Zealand. Arc magmatism was broadly concurrent across the
34 breadth of the Havre Trough.

35

36 **Keywords**

37 Havre Trough, Kermadec Arc, Ar-Ar, magmatism, back-arc basin, rifting

38

39 **Introduction**

40 The present-day Kermadec Arc and associated Havre Trough back-arc basin is the youngest
41 in a series of Cenozoic volcanic arcs that have developed along the northern New Zealand
42 margin in response to convergence of the Pacific and Australian Plates (Mortimer et al.,
43 2010; Herzer et al., 2011; Bassett et al., 2016). The Kermadec Arc - Havre Trough (KAHT)
44 subduction system is the central portion of a contiguous arc system, with the Tonga Arc –
45 Lau Basin back-arc system to the north, and the Taupo Volcanic Zone (TVZ) of continental
46 New Zealand to the south (Figure 1) (Smith and Price, 2006). The predecessor to the
47 Kermadec Arc, the Miocene-Pliocene Colville Arc (Skinner, 1986; Ballance et al., 1999),
48 rifted apart in response to rollback of the Pacific Plate (Sdrolias and Muller, 2006; Wallace et
49 al., 2009), forming the Havre Trough and resulting in the establishment of the modern
50 Kermadec Arc front. The Colville Ridge and Kermadec Ridge are the remnants of the
51 Colville Arc (Figure 1).

52 The age of opening of the Havre Trough and establishment of the Kermadec Arc is
53 not clear owing to a paucity of age data. In part, this is due to the inherent difficulty in
54 obtaining reliable radioisotopic ages on young, glassy, and vesicular submarine volcanic
55 rocks with low potassium content, and in part due to tectonic complexity, and until recently,
56 limited seafloor sampling in the region. Here, we present ^{40}Ar - ^{39}Ar ages on seafloor volcanic
57 samples from across the southern KAHT subduction system that have important implications
58 for both the age and style of opening of the Havre Trough.

59

60 **Models for opening of the Havre Trough**

61 Several models have been proposed to explain the tectono-magmatic evolution of the Havre
62 Trough and Kermadec Arc, but the process and timing of opening remains contentious.
63 Malahoff et al. (1982), based on airborne magnetic studies and seismic lines over the
64 southern and central portions of the KAHT, tentatively interpreted the Havre Trough to be
65 undergoing spreading, centred on an axial ridge. They interpreted residual magnetic

66 anomalies to indicate a ca. 1.8 Ma age of opening of the basin. Wright (1993), however,
67 interpreted swath mapping data as showing that at least the southern Havre Trough lacked a
68 medial spreading ridge, and hence interpreted back-arc rifting rather than spreading as the
69 mode of extension. Further, Wright (1993), suggested that initiation of rifting occurred at ca.
70 5 Ma, although this age was constrained by extrapolation of geodetic data on continental New
71 Zealand rather than on direct age data from within the Havre Trough.

72 Subsequent models for Havre Trough opening agreed that the system was rifting but
73 have varied in the process and style of rifting being proposed. Wright et al. (1996) suggested
74 that Havre Trough opening and magmatism progressed eastward with time. Parson and
75 Wright (1996) further argued that there was a latitudinal progression from full oceanic
76 spreading in the Lau Basin to the north, to basin rifting in the TVZ to the south. The southern
77 Havre Trough was considered to be in an intermediate phase of rifting that was concentrated
78 along the axial zone of the trough. Ruellan et al. (2003), on the basis of multibeam
79 bathymetry and seismic reflection data, concluded that the southward propagation of
80 spreading was oversimplified, and that southward migration of subduction of the Louisville
81 Seamount Chain had effectively locked the KAHT. They proposed that opening of the Havre
82 Trough was initially fast and pervasive, and then relatively quiescent as the system became
83 locked. Wysoczanski et al. (2010), on the basis of morphological similarities, suggested that
84 the Havre Trough was in a similar state of rifting to the Valu Fa Ridge and Western Lau
85 Basin, and that it also was in a state of “disorganised spreading” (Martinez and Taylor, 2006)
86 whereby diffuse patches of extension localised in deep rifts precedes longitudinally traceable
87 axial ridges characteristic of true ocean spreading systems. This model reconciled the oceanic
88 spreading model of Malahoff et al. (1982) with models of rifting, and is similar to the Parson
89 and Wright (1996) final stage of rifting (their “Phase 4”) preceding full spreading.

90

91 **Analytical Methods and Results**

92 A total of 19 volcanic rocks of variable composition dredged from across the KAHT (Table
93 1) have been dated by Ar-Ar step heating. The sample set is diverse, including samples from
94 five arc front volcanoes, two volcanoes in the central Havre Trough (Gill and Rapuhia), a
95 deep central Havre Trough basin (Ngatoro Rift) with a short axial ridge in its southern extent,
96 and a cross-arc seamount chain (Rumble V Ridge) that spans the breadth of the Havre
97 Trough, from Rumble V to the Colville Ridge (Figure 2). Geochemical data for all the
98 samples have previously been reported, and the source of those data, together with new Ar-

99 Ar ages presented here, are shown in Table 1. With the exception of one andesite and one
100 dacite from the volcanic arc front, all samples are basalts or basaltic andesites (Figure 3).

101 Ar-Ar analyses were performed in three laboratories (USGS, Menlo Park; New
102 Mexico Institute of Mining and Technology (NMIMT), Socorro; and University of
103 Wisconsin-Madison), initially as four smaller and separate studies. The datasets are combined
104 here as one larger study to place constraints on the age of the KAHT (Table 1, Figure 2). All
105 ages presented in Table 1 include 2σ uncertainties and full details of the analytical techniques
106 are given in the Supplementary File.

107 The majority of ages for the arc front volcanoes are <0.06 Ma, although two samples,
108 from Clark (C/1) and Rumble III (X333) have slightly older mean ages of 0.11 Ma and 0.12
109 Ma respectively. Uncertainties on arc front samples however are large, with most ages having
110 2σ uncertainties of zero age, and most ages are zero within analytical uncertainty.

111 Three samples from Rumble V Ridge have ages of <0.11 Ma, overlapping those of
112 the arc front volcanoes within uncertainty. The Ngatoro Rift samples have older ages
113 between 0.20 Ma and 0.68 Ma.

114 To the north, two samples from Rapuhia Ridge, a volcanic ridge extending southwest
115 from Rapuhia volcano in the centre of the Havre Trough, yielded ages of 0.05 ± 0.05 Ma and
116 0.11 ± 0.03 Ma. These ages are marginally older than, but within error of, ages derived from
117 the active volcanic arc front. They are on average younger than the samples from Rumble V
118 Ridge [see above], and notably younger than most of the Ngatoro Rift samples. Three
119 samples analysed from Gill volcano, a back-arc volcano in the Havre Trough that lies
120 between Rapuhia Ridge and the Colville Ridge (Figure 1), have ages significantly older than
121 all other samples, at 0.88 ± 0.05 Ma, 0.97 ± 0.03 Ma and 1.19 ± 0.04 Ma.

122

123 Discussion

124 The presented Ar-Ar ages are from samples that span almost the entire width of the southern
125 Havre Trough and thus provide important constraints on the manner and timing of its
126 opening.

127 A first order observation is that the oldest ages reported here, from a back-arc
128 stratovolcano (Gill volcano: Wysoczanski et al., 2010) in the western part of the Havre
129 Trough, are 0.9 - 1.2 Ma (Table 1, Figure 2). However, because Gill volcano sits on a rifted
130 basin floor, the implied age of rifting must be older. This age is similar to a preferred Ar-Ar
131 age of 1.1 ± 0.4 Ma reported for a basalt from the western Havre Trough (Mortimer et al.,

132 2007) sampled 450 km to the north of, and along strike from, Gill volcano, and to a $1.25 \pm$
133 0.06 Ma U-Pb zircon age from a tonalite xenolith from Raoul Island (Mortimer et al., 2010).
134 In addition, Mortimer et al. (2007) reported an Ar-Ar age of $1.2 \text{ Ma} \pm 0.8$ for a basalt from
135 the Northland Plateau (Figure 1), which they considered to be related to westernmost Colville
136 Ridge volcanism. Together, these ages show no evidence for magmatic activity in the Havre
137 Trough before c. 1.2 Ma, and as noted by Mortimer et al. (2010) suggest that magmatism was
138 active across the full width of the KAHT and west of the Colville Ridge at this time (Figure
139 2). Furthermore, one of our plateau ages from Gill volcano is 875 ± 50 ka, and thus it is
140 conceivable that the age of magmatism for the Havre Trough is younger than 1.2 Ma, and
141 possibly < 1 Ma.

142 Using the 19 new Ar/Ar ages presented in this study and two previously reported by
143 Mortimer et al. (2007; 2010), we now have sufficient geochronologic data to interpret the age
144 of the Havre Trough. In addition, Ballance et al. (1999) reported eight K-Ar ages of c. 2 Ma
145 or younger for the Kermadec Ridge and three K-Ar ages from the eastern Havre Trough,
146 which were near zero age (the oldest at 0.15 ± 0.12 Ma). These ages for the Havre Trough are
147 all significantly younger than the c. 5 Ma age of rifting proposed by Wright (1993). However,
148 we note that all current age data are from surficial seafloor volcanics, and future sampling
149 (especially from sub-seafloor drilling) may yield older ages that would require a
150 reinterpretation of the results presented here.

151 The young age of magmatism, if correct, provides three important implications for the
152 tectonic development of the Havre Trough.

153 Firstly, magmatism and translocation of the modern Kermadec Arc front did not occur
154 in a monotonic eastward progression. Notably, there is near- zero age arc magmatism in the
155 central portion of the Havre Trough at Rapuhia Ridge, and magmatism related to Rumble V
156 Ridge does not young to the east (Figure 4). The Rumble V Ridge dates are younger in age
157 than the Ngatoro Rift, indicating that the ridge may have been constructed over the Ngatoro
158 Rift (and if this is correct, also the Rumble Rift), rather than being cut by rifting as previously
159 suggested (Wright et al., 1996).

160 Second, reported age data for the Havre Trough is < 1.2 Ma, and possibly < 1 Ma.
161 This is younger than, but broadly consistent with, the 1.8 Ma age of rifting suggested by
162 Malahoff et al. (1982), although that model assumed a full spreading centre, whereas more
163 recent tectonic models based on seafloor morphology suggest that the Havre Trough is
164 comprised of a number of rifts and basal plateaus (e.g. Wright, 1993; Wysoczanski et al.,
165 2010; Wysoczanski and Clark, 2012). These ages infer a c. 2.5-4 x faster extension rate for

166 the Havre Trough than the 15-20 mm yr⁻¹ rate suggested by Wright (1993). An age of 2 Ma
167 would give an average rate of c. 40-50 mm yr⁻¹. Whilst reasonably fast, this rate is not
168 unusual for extension rates in other intra-oceanic back-arc rifts, and is still significantly
169 slower than the full ocean spreading rates of > 100 mm yr⁻¹ occurring in the Lau Basin and
170 Manus Basin (e.g. Taylor and Martinez, 2003; Heuret & Lallemand, 2005; Wallace et al.,
171 2005). Notably this is similar to the extension rate of c. 40-60 mm yr⁻¹ seen at the southern
172 portion of the Lau Basin (Parson and Wright, 1996; Martinez and Taylor, 2001).

173 Third, opening of the Havre Trough is coeval with initiation of TVZ magmatism and
174 rifting at c. 2 Ma (Wilson et al., 1995) and the TVZ rift and Havre Trough are the continental
175 and oceanic expression of the same rift system (e.g. Parson and Wright, 1996). It is unclear if
176 rifting was occurring prior to c. 2 Ma onshore in New Zealand: 1.8-3.9 Ma volcanism
177 occurred along the Maungatautari-Kaimai-Tauranga alignment parallel to but northwest of
178 the TVZ, as eruptions migrated southeast from the Coromandel area (Briggs et al., 2005).
179 Given our ages for the Havre Trough, and that the youngest reported age of volcanism from
180 the Colville Ridge is 2.6 Ma (Timm et al., in review), this magmatism is more likely to be
181 related to Colville Arc magmatism rather than Havre Trough magmatism.

182 The western portion of the TVZ is the oldest part of that system (the “old TVZ” of
183 Wilson et al., 1995, and Wilson and Rowland, 2016), and rifting is now focussed more to the
184 east and along a central rift, variously defined as the “young TVZ” and “modern TVZ”
185 (Wilson et al., 1995; Wilson and Rowland, 2016), Ruaumoko Rift (Rowland and Sibson,
186 2001) and the Taupo Rift (Villamor and Berryman, 2006). Whilst young arc magmatism is
187 broadly occurring across the Havre Trough (Figure 4) we have insufficient data to identify
188 any age progression of rift-related magmatism across the Havre Trough. It remains uncertain
189 if eastern Havre Trough rift magmatism is younger than western Havre Trough rift
190 magmatism, and so akin to the old and young/modern TVZ regions, respectively.

191 The present state of extension/rifting of the Havre Trough remains uncertain. In the
192 case of the Ngatoro Rift, the ages presented here indicate prolonged magmatism over at least
193 0.4 Ma, and that the rift is not presently magmatically active at the seafloor. Importantly
194 though there is extensive shallow seismic activity (< 13 km deep) within the Ngatoro Rift (de
195 Ronde et al., 2007). Regional moment tensor analysis for recent (2003-2012) shallow (< 33
196 km) earthquakes in the southern Havre Trough show extension as well as strike slip
197 movement (Ristau, 2014). At first order the shallow extensional seismicity in the Ngatoro
198 Rift and elsewhere in the Havre Trough indicates present-day extension / rifting of the
199 trough. Magmatic rift intrusives (e.g., dykes) may also be contemporaneous, however the

200 absence of present day surficial extrusives and lack of hydrothermal activity suggests that
201 seafloor, or near seafloor, rift magmatism is not occurring at the present day.

202

203 **Conclusions**

204 New Ar-Ar ages presented here, coupled with other published radioisotopic ages from the
205 literature (Ballance et al., 1999; Mortimer et al., 2007, 2010), suggest that opening of the
206 Havre Trough initiated < c. 2 Ma, and perhaps as recently as c. 1 Ma. The oldest ages occur
207 on the margins of the basin and significant young arc magmatism occurred across the central
208 Havre Trough. The timing of initiation of magmatism is coeval with that of the TVZ. The
209 caveat to our age constraints is that all samples are surficial and there are no ages for samples
210 within c. 25 km of the Colville Ridge (Figure 4).

211 Our results show that there has been arc and rift-related magmatism across the entire
212 southern Havre Trough within the last c. 1 Ma, both within rifts (e.g., Ngatoro Rift) and
213 constructing large stratovolcano cones such as Gill and seamounts of Rumble V Ridge
214 (Wright et al., 1996; Todd et al., 2010). This, together with the >4 km water depth in the
215 deepest parts of the basin, is more consistent with distributed rifting across the basin than
216 ocean spreading. Whether there are differences in age between rift-related magmas erupted at
217 different depths, or distance across the basin, or distance northward from New Zealand, is
218 important for understanding the tectonic evolution of the basin but remains to be discovered.
219 Our experience shows that $^{40}\text{Ar}/^{39}\text{Ar}$ ages can be obtained for the challenging Havre Trough
220 samples, but that sample selection and treatment are important considerations.

221

222 **Acknowledgements**

223 The authors would like to thank Erin Todd for his internal review, and Roger Briggs and an
224 anonymous reviewer for their helpful reviews. RW was funded by the Ministry of Business,
225 Innovation and Employment (MBIE) Strategic Science Investment Fund (SSIF) programme
226 Marine Geological Processes and Resources (COPR1902). CT received funding from the
227 European Union's Horizon 2020 research and innovation programme under the Marie
228 Skłodowska-Curie grant agreement #79308.

229

230

231 **References**

- 232 Ballance PF, Ablaev AG, Pushchin IK, Pletnev SP, Biryulina MG, Itaya T, Follas HA, Gibson,
233 GW. 1999. Morphology and history of the Kermadec trench-arc-backarc basin-remnant arc
234 system at 30 to 32°S: geophysical profile, microfossil and K–Ar data. *Marine Geology*.
235 159:35–62.
236
- 237 Bassett D, Kopp H, Sutherland R, Henrys S, Watts AB, Timm C, Scherwath M, Grevemeyer
238 I, de Ronde CEJ. 2016. Crustal structure of the Kermadec arc from MANGO seismic
239 refraction profiles. *Journal of Geophysical Research*. 121:7514-7546.
240
- 241 Briggs RM, Houghton BF, McWilliams M, Wilson CJN. 2005. $^{40}\text{Ar}/^{39}\text{Ar}$ ages of silicic
242 volcanic rocks in the Tauranga-Kaimai area, New Zealand: Dating the transition between
243 volcanism in the Coromandel Arc and the Taupo Volcanic Zone. *New Zealand Journal of*
244 *Geology and Geophysics*. 48:459-469.
245
- 246 DeMets C, Gordon RG, Argus DF. 2010. Geologically current plate motions. *Geophysical*
247 *Journal International*. 181(1):1–80.
248
- 249 de Ronde CEJ, Baker ET, Massoth GJ, Lupton JE, Wright IC, Sparks RJ, Bannister SC,
250 Reyners ME, Walker SL, Greene RR, Ishibashi J, Faure K, Resing JA, Lebon GT. 2007.
251 Submarine hydrothermal activity along the mid-Kermadec Arc, New Zealand: Large-scale
252 effects on venting. *Geochemistry Geophysics Geosystems*. 8. Q07007, doi:
253 10.1029/2006GC001495.
254
- 255 Gamble JA, Wright IC, Baker JA. 1993. Seafloor geology and petrology in the oceanic to
256 continental transition zone of the Kermadec-Havre-Taupo Volcanic Zone arc system, New
257 Zealand. *New Zealand Journal of Geology and Geophysics*. 36:417-435.
258
- 259 Gamble JA, Christie HK, Wright I, Wysoczanski R. 1997. Primitive K-rich magmas from
260 Clark volcano, southern Kermadec Arc: A paradox in the K – depth relationship. *The*
261 *Canadian Mineralogist*. 35:275-290.
262
- 263 Herzer RH, Barker DHN, Roest WR, Mortimer N. 2011. Oligocene-Miocene spreading
264 history of the northern South Fiji Basin and implications for the evolution of the New

- 265 Zealand plate boundary. *Geochemistry, Geophysics, Geosystems*. 12. Q02004,
266 doi:10.1029/2010GC003291.
267
- 268 Heuret A, Lallemand S. 2005. Plate motions, slab dynamics and back-arc deformation.
269 *Physics of the Earth and Planetary Interiors*. 149:31–51
270
- 271 Malahoff A, Feden R, Fleming H. 1982. Magnetic anomalies and tectonic fabric of marginal
272 basins north of New Zealand. *Journal of Geophysical Research*. 87:4109–4125.
273
- 274 Martinez F, Taylor B. 2001. Mantle wedge control on back-arc crustal accretion. *Nature*.
275 416:417-420.
276
- 277 Martinez F, Taylor B. 2006. Modes of crustal accretion in back-arc basins: inferences from
278 the Lau Basin. In: Christie DM, Fisher CR, Lee S-M, Givens S, editors. *Back- arc Spreading*
279 *Systems: Geological, Biological, Chemical and Physical Interactions*. Geophysical
280 Monograph Series 166. American Geophysical Union. p. 5–30.
281
- 282 Mortimer N, Gans PB, Palin JM, Meffre S, Herzer RH, Skinner DNB. 2010. Location and
283 migration of Miocene–Quaternary volcanic arcs in the SW Pacific region. *Journal of*
284 *Volcanology and Geothermal Research*. 190:1–10.
285
- 286 Mortimer N, Herzer RH, Gans PB, Laporte-Magoni C, Calvert AT, Bosch D. 2007.
287 Oligocene–Miocene tectonic evolution of the South Fiji Basin and Northland Plateau, SW
288 Pacific Ocean: evidence from petrology and dating of dredged rocks. *Marine Geology*.
289 237:1–24.
290
- 291 Parson LM, Wright IC, 1996. The Lau–Havre–Taupo back-arc basin: a southward pro-
292 pagating, multi-stage, evolution from rifting to spreading. *Tectonophysics*. 263:1–22.
293
- 294 Ristau J. 2014. Update of Regional Moment Tensor Analysis for Earthquakes in New
295 Zealand and Adjacent Offshore Regions. *Bulletin of the Seismological Society of America*.
296 103:2520–2533.
297

- 298 Ruellan E, Delteil J, Wright I, Matsumoto T. 2003. From rifting to spreading in the Lau Basin
299 – Havre Trough backarc system (SW Pacific) – locking/unlocking induced by ridge
300 subduction. *Geochemistry, Geophysics, Geosystems*. 7. doi:10.1029/2001GC000261.
301
- 302 Rowland JV, Sibson RH. 2001. Extensional fault kinematics within the Taupo Volcanic
303 Zone, New Zealand: soft-linked segmentation of a continental rift system. *New Zealand
304 Journal of Geology and Geophysics*. 44:271-283.
305
- 306 Sdrolias M, Muller RD. 2006. Controls on back-arc formation. *Geochemistry, Geophysics,
307 Geosystems*. 7. Q04016, doi:10.1029/2005GC001090.
308
- 309 Skinner DNB. 1986. Neogene volcanism of the Hauraki Volcanic Region. *Royal Society of
310 New Zealand Bulletin*. 23:21-47.
311
- 312 Smith IEM, Price RC. 2006. The Tonga-Kermadec arc and Havre-Lau back-arc system: Their
313 role in the development of tectonic and magmatic models for the western Pacific. *Journal of
314 Volcanology and Geothermal Research*. 156:315-331.
315
- 316 Taylor B, Martinez F. 2003. Back-arc basin basalt systematics. *Earth and Planetary Science
317 Letters*. 210:481-497.
318
- 319 Timm C, de Ronde CEJ, Hoernle K, Cousens B, Wartho J-A, Caratori Tontini F,
320 Wysoczanski R, Hauff F, Handler M. Accepted. Colville and Kermadec Ridges, SW Pacific:
321 new radiometric age data and insights into the recent geological history and petrogenesis of
322 northern Zealandia. *Gondwana Research*.
323
- 324 Todd E, Gill JB, Wysoczanski RJ, Handler MR, Wright IC, Gamble JA. 2010. Sources of
325 constructional cross-chain volcanism in the southern Havre Trough: New insights from HFSE
326 and REE concentration and isotope systematics. *Geochemistry, Geophysics, Geosystems*. 11.
327 Q04009, doi:10.1029/2009GC002888.
328
- 329 Todd E, Gill JB, Wysoczanski RJ, Hergt J, Wright IC, Leybourne MI, Mortimer, N. 2011. Hf
330 isotopic evidence for small-scale heterogeneity in the mode of mantle wedge enrichment:

- 331 Southern Havre Trough and South Fiji Basin back-arcs. *Geochemistry, Geophysics,*
332 *Geosystems*. 11. Q09011, doi:10.1029/2011GC003683.
- 333
- 334 Villamor P, Berryman KR. 2006. Evolution of the southern termination of the Taupo Rift,
335 New Zealand. *New Zealand Journal of Geology and Geophysics*. 49:23-37.
- 336
- 337 Wallace LM, McCaffrey R, Beavan J, Ellis S. 2005. Rapid microplate rotations and backarc
338 rifting at the transition between collision and subduction. *Geology*. 33:857-860.
- 339
- 340 Wallace LM, Ellis S, Mann P. 2009. Collisional model for rapid fore-arc block rotations, arc
341 curvature, and episodic back-arc rifting in subduction settings. *Geochemistry, Geophysics,*
342 *Geosystems*. 10. Q05001, doi:10.1029/2008GC002220.
- 343
- 344 Wilson CJN, Houghton BF, McWilliams MO, Lanphere MA, Weaver SD, Briggs RM. 1995.
345 Volcanic and structural evolution of Taupo Volcanic Zone, New Zealand: a review. *Journal*
346 *of Volcanology and Geothermal Research*. 68:1-28.
- 347
- 348 Wilson CJN, Rowland JV. 2016. The volcanic, magmatic and tectonic setting of the Taupo
349 Volcanic Zone, New Zealand, reviewed from a geothermal perspective. *Geothermics*.
350 59:168–187.
- 351
- 352 Wright IC. 1993. Pre-spread rifting and heterogeneous volcanism in the southern Havre
353 Trough back-arc basin. *Marine Geology*. 113:179–200.
- 354
- 355 Wright IC, Parson LM, Gamble JA. 1996. Evolution and interaction of migrating cross-arc
356 volcanism and backarc rifting: an example from the southern Havre Trough (35°20'–37°S).
357 *Journal of Geophysical Research*. 101:22071–22086.
- 358
- 359 Wysoczanski R, Clark M. 2012. Southern Kermadec Arc – Havre Trough Geohabitats. In:
360 Harri, P, Baker E, editors. *Seafloor Geomorphology as Benthic Habitat: Geohab Atlas of*
361 *Seafloor Geomorphic Features and Benthic Habitats*. Elsevier. p. 853-867.
- 362
- 363 Wysoczanski RJ, Todd E, Wright IC, Leybourne MI, Hergt JM, Adam C, Mackay K. 2010.
364 Backarc rifting, constructional volcanism and nascent disorganised spreading in the southern

365 Havre Trough backarc rifts (SW Pacific). *Journal of Volcanology and Geothermal Research*.
366 190:39-57.

367

368 Zohrab A. 2017. The petrology, geochemistry and geochronology of back-arc stratovolcanoes
369 in the southern Kermadec Arc-Havre Trough, SW Pacific. MSc. Victoria University of
370 Wellington.

371

372 **Figures**

373

374 Figure 1: Tectonic setting of New Zealand and the SW Pacific highlighting the Kermadec
375 Arc – Havre Trough (KAHT), the Tonga-Lau subduction system, and the Taupo Volcanic
376 Zone (TVZ) of continental New Zealand (red outline). Black arrow is the relative motion of
377 the Pacific Plate to a fixed Australian Plate for the southern KAHT region (DeMets et al.,
378 2010). HP = Hikurangi Plateau, Louisville SC = Louisville Seamount Chain, NP = Northland
379 Plateau, VFR = Valu Fa Ridge. Red triangles denote oceanic volcanoes of the Kermadec Arc
380 and Havre Trough, and the offshore TVZ (southernmost volcano, Whakatane). Highlighted
381 area is that of Figure 2.

382

383 Figure 2: Bathymetric map of the southern KAHT system, bounded by the Colville Ridge to
384 the west and the Kermadec Ridge to the east. Depths on the bathymetry scale are metres
385 below sea level, with depths < 1500 m shown as 1500 m and depths > 3500 m shown as 3500
386 m. Orange triangles are volcanoes: C = Clark, G = Gill, R = Rapuhia, RIII = Rumble III, RIV
387 = Rumble IV, RV = Rumble V, T = Tangaroa. Numbers in boxes denote new Ar/Ar ages
388 (Table 1).

389

390 Figure 3: Silica content of samples analysed in this study with distance from the crest of the
391 Kermadec Ridge.

392

393 Figure 4: Ar/Ar ages of Havre Trough samples (Table 1) with distance from the Kermadec
394 Ridge crest. Error bars show 2 sigma uncertainties. Black diamonds are K/Ar ages of
395 Ballance et al. (1999) from Kermadec Ridge and Havre Trough samples at least 300 km north
396 of samples presented here. Grey square at ~80 km is an Ar/Ar preferred age for a basalt from
397 the Havre Trough (Mortimer et al., 2007). Grey square at 0 km is a U-Pb age of zircon from a

398 tonalite from Raoul volcano (Mortimer et al., 2010), 600 km to the north of the study area,
399 where the modern arc front sits on the Kermadec Ridge (Figure 1).

400

401 **Table**

402

403 Table 1: Details of samples analysed in this study. Ages are: P=plateau ages, I=Isochron
404 ages, R=Recoil age (see Supplementary File for details). Supplementary File contains plateau
405 and isochron ages and plots, experimental data including K/Ca ratio, MSWDs, number of
406 steps, and total gas age; along with an explanation of experimental methods and machine data
407 for individual heating steps within each experiment. Results have been recalculated to a
408 consistent fluence monitor age equivalent to Fish Canyon sanidine at 28.198 Ma (Menlo
409 Park) and at 28.201 Ma (NMIMT). All errors are 2σ . For four samples, X379, X690, X682,
410 and X696 the mean age is negative, so the positive fraction of the age is reported as a
411 maximum value (i.e. <xx Ma), calculated as the mean of the 2σ error. IGSN numbers are
412 given for those samples that have been assigned numbers. Reference for geochemical
413 analyses: 1, Gamble et al, 1997; 2, Wright and Gamble, unpublished data; 3, Gamble et al.,
414 1993; 4, Todd et al., 2010; 5, Zohrab, 2017; 6, Todd et al., 2011. All geochemical data are
415 reported as anhydrous, with Fe as $\text{FeO}_{\text{total}}$.

STATION	LOCATION	LATITUDE SOUTH	LONGITUDE EAST	DEPTH M	LAB	LAB NO.	IGSN	REF.	SiO ₂ WT.%	MgO WT.%	K ₂ O WT.%	AGE Ma	
C/1	Clark	36.416	177.848	2040	NMIMT	Clark #45, 6696		2	50.75	9.46	1.57	0.11 ± 0.05	P
X299	Rumble III	35.749	178.498	717	NMIMT	Rumble III #1, 6692		2	52.61	4.44	0.58	0.04 ± 0.06	P
X333	Rumble III	35.715	178.528	565	NMIMT	Rumble III #8, 6695		2	52.14	6.72	0.48	0.12 ± 0.08	P
X351	Rumble IV	36.131	178.024	1258	NMIMT	Rumble IV #9, 6703		2	66.19	1.47	1.11	0.03 ± 0.02	P
X379	Rumble V	36.153	178.161	1619	NMIMT	Rumble V#23, 6694	JBG000010	2	54.00	3.51	0.60	<0.03	P
X407	Rumble V	36.133	178.202	750	NMIMT	Rumble V #26, 6704		2	53.95	3.52	0.61	0.01 ± 0.06	P
X427/A	Tangaroa	36.311	178.004	1781	NMIMT	Tangaroa #39, 6691		2	59.26	2.63	0.67	0.06 ± 0.07	P
X153/1	Ngatoro Rift	36.260	177.300	2640	NMIMT	11574 Ngatoro Rift, 6702	JBG00001C	3	51.01	8.22	0.41	0.20 ± 0.14	P
X158/1	Ngatoro Rift	36.154	177.428	2300	NMIMT	11580 Ngatoro Rift, 6701		3	52.04	7.05	0.52	0.60 ± 0.24	P
X185/1	Ngatoro Rift	36.660	177.150	2810	NMIMT	11616 S. Ngatoro Rift, 6693	JBG000016	3	52.41	4.86	0.55	0.35 ± 0.22	P
X168/1A	Ngatoro Rift	36.258	177.573	2960	Menlo Park	10Z0107	JBG000017	3	52.84	7.38	0.60	0.68 ± 0.16	R
X690A	Cross arc	35.960	177.942	1805	Menlo Park	10Z0105	JBG000001	4	47.23	14.9	0.32	<0.11	I
X682	Cross arc	35.968	178.023	1480	Menlo Park	10Z0106	JBG000007	4	51.13	8.17	0.42	<0.03	I
X696A	Cross arc	35.886	177.843	1680	Menlo Park	10Z0104	JBG00000J	4	48.94	8.46	0.28	<0.07	I
015-04	Rapuhia Ridge	34.794	178.445	1910	Menlo Park	15Z0332		5	51.04	9.65	0.75	0.11 ± 0.03	P
016-01	Rapuhia Ridge	34.798	178.442	1800	Menlo Park	15Z0334		5	49.60	9.99	0.49	0.05 ± 0.05	P
012-01	Gill	34.623	178.379	1146	Menlo Park	15Z0319		5	47.91	9.30	0.46	1.19 ± 0.04	P
011-04	Gill	34.607	178.389	1700	Menlo Park	15Z0318		5	51.22	8.07	0.75	0.97 ± 0.03	P
011-A	Gill	34.607	178.389	1700	Wisconsin	UW93C37	JBG00001K	6	53.64	6.59	0.77	0.88 ± 0.05	P

Ages are: P=plateau ages, I=Isochron ages, R=Recoil age (see Supplementary File for details). Supplementary File contains plateau and isochron ages and plots, experimental data including K/Ca ratio, MSWDs, number of steps, and total gas age; along with an explanation of experimental methods and machine data for individual heating steps within each experiment. Results have been recalculated to a consistent fluence monitor age equivalent to Fish Canyon sanidine at 28.198 Ma (Menlo Park) and at 28.201 Ma (NMIMT). All errors are 2σ . For the four samples X379, X690, X682, and X696 the mean age is negative, so the positive fraction of the age is reported as a maximum value (i.e. <xx Ma), calculated as the mean of the 2σ error. IGSN numbers are given for those samples that have been assigned numbers.

Reference for geochemical analyses: 1, Gamble et al, 1997; 2, Wright & Gamble unpublished data; 3, Gamble et al., 1993; 4, Todd et al., 2010; 5, Zohrab, 2017; 6, Todd et al., 2011. All geochemical data is reported as anhydrous, with Fe as FeOtotal (not reported here).

For Peer Review Only




Figure 1: Tectonic setting of New Zealand and the SW Pacific highlighting the Kermadec Arc – Havre Trough (KAHT), the Tonga-Lau subduction system, and the Taupo Volcanic Zone (TVZ) of continental New Zealand (red outline). Black arrow is the relative motion of the Pacific Plate to a fixed Australian Plate for the southern KAHT region (DeMets et al., 2010). HP = Hikurangi Plateau, Louisville SC = Louisville Seamount Chain, NP = Northland Plateau, VFR = Valu Fa Ridge. Red triangles denote oceanic volcanoes of the Kermadec Arc and Havre Trough, and the offshore TVZ (southernmost volcano, Whakatane). Highlighted area is that of Figure 2.

163x225mm (300 x 300 DPI)




Figure 2: Bathymetric map of the southern KAHT system, bounded by the Colville Ridge to the west and the Kermadec Ridge to the east. Depths on the bathymetry scale are metres below sea level, with depths < 1500 m shown as 1500 m and depths > 3500 m shown as 3500 m. Orange triangles are volcanoes: C = Clark, G = Gill, R = Rapuhia, RIII = Rumble III, RIV = Rumble IV, RV = Rumble V, T = Tangaroa. Numbers in boxes denote new Ar/Ar ages (Table 1).

210x296mm (300 x 300 DPI)




Figure 3: Silica content of samples analysed in this study with distance from the crest of the Kermadec Ridge.

209x126mm (300 x 300 DPI)




Figure 4: Ar/Ar ages of Havre Trough samples (Table 1) with distance from the Kermadec Ridge crest. Error bars show 2 sigma uncertainties. Black diamonds are K/Ar ages of Ballance et al. (1999) from Kermadec Ridge and Havre Trough samples at least 300 km north of samples presented here. Grey square at ~80 km is an Ar/Ar preferred age for a basalt from the Havre Trough (Mortimer et al., 2007). Grey square at 0 km is a U-Pb age of zircon from a tonalite from Raoul volcano (Mortimer et al., 2010), 600 km to the north of the study area, where the modern arc front sits on the Kermadec Ridge (Figure 1).

209x126mm (300 x 300 DPI)