



Carbohydrate binding module of family 48 enable ferulic acid esterases action on polymeric arabinoxylan

Wilkens, Casper

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Wilkens, C. (2019). Carbohydrate binding module of family 48 enable ferulic acid esterases action on polymeric arabinoxylan. Poster session presented at 13th Carbohydrate Bioengineering Meeting (CBM), Toulouse, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Carbohydrate binding module of family 48 enable ferulic acid esterases action on polymeric arabinoxylan

Jesper Holck¹, Folmer Fredslund², Marie S. Møller³, Jesper Brask⁴, Kristian B. R. M. Krogh⁴, Birte Svensson³, Ditte H. Welner², Anne S. Meyer¹, Stefan Janeček^{5,6} and Casper Wilkens¹

¹Enzyme Technology, Dept. of Biotechnology and Biomedicine, Technical University of Denmark, Søtofts Plads, Building 224, DK-2800 Kgs. Lyngby, Denmark.

²Enzyme Engineering and Structural Biology, Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Søtofts Plads, Building 220, DK-2800 Kgs. Lyngby, Denmark.

³Enzyme and Protein Chemistry, Dept. of Biotechnology and Biomedicine, Technical University of Denmark, Søtofts Plads, Building 224, DK-2800 Kgs. Lyngby, Denmark

⁴Novozymes A/S, Krogshøjvej 36, DK-2880 Bagsværd, Denmark

⁵Laboratory of Protein Evolution, Institute of Molecular Biology, Slovak Academy of Sciences, Bratislava, Slovakia

⁶Dept. of Biology, University of SS. Cyril and Methodius, Trnava, Slovakia
E-mail: cwil@dtu.dk

Keywords: CBM48, CE1, ferulic acid esterase, structure-function relationships

Arabinoxylans (AXs) are a major component of hemicelluloses, which is widely distributed in secondary cell walls of plants. Their backbone is composed of β -1,4-linked xylopyranose residues that are single substituted with α -L-1,3-arabinofuranose (Araf) or double substituted with both α -L-1,2- and α -L-1,3-Araf, which can be further substituted with by 5-O-ferulic acid and other hydroxycinnamic acids [1]. Ferulic acid esterases (FAEs) catalyze the hydrolysis of ester bonds between hydroxycinnamic acids and Araf, however, most characterized FAEs do not display a preference for polymeric substrates [2,3].

Recently, a number of carbohydrate esterase family 1 (CE1) identified in metagenomic studies was shown to have a carbohydrate binding module of family 48 (CBM48) appended [4,5], a family associated with starch binding [6]. Our phylogenetic analysis demonstrated that these are in fact CBM48s suggesting that CBM48s is a polyspecific family since CE1s do not target starch. Adsorption assays with two CE1-CBM48 enzymes demonstrated binding to AXs, but not to starch, which was supported by a surface plasmon resonance analysis showing no binding to β -cyclodextrin or maltohexaose. Binding was detected to arabino- and xylooligosaccharides and interestingly also to maltotetraose. The two CE1-CBM48 enzymes released FA from AXs, while the CE1 domain on its own only released FA from oligosaccharides and unlike the full-length enzymes the CE1 domain was unable to bind to AXs. Crystal structures of the two CE1-CBM48s revealed two integrally folded units and multiple structurally conserved hydrogen bonds fix the CBM48's position relative to the CE1 domains. Molecular dynamics simulations confirmed that the two domains form a rigid structure. Docking studies suggest that the xylan main chain is accommodated in the cleft formed at the interface between the CE1 and CBM48 domains.

[1] Terrett O.M.; Dupree P. *Curr. Opin. Biotechnol.* **2019**, 56, 97–104

[2] Olivera D.M; Mota T.R.; Oliva B. *et al.* **2019**, *Bioresour. Technol.* 278, 408–23

[3] Ramos-de-la-Peña A.M.; Contreras-Esquivel J.C. **2016**, *J. Mol. Catal., B* 130, 74–87

[4] Wilkens C.; Busk P.K.; Pilgaard B. *et al.* **2017**, *Biotechnol Biofuels* 10, 158

[5] Wong M. T.; Wang W.; Couturier, M. *et al.* **2017**, *Front. Microbiol.* 8, 2504

[6] Janeček S.; Svensson B.; MacGregor E.A. **2011**, *Enzyme Microb. Technol.* 49, 429–440