

指 導 教 授 氏 名	指 導 役 割
久保田聡 印	
印	
印	

学 位 論 文 要 旨

岡山大学大学院医歯薬学総合研究科

専攻分野 口腔生化学	身分 大学院生	氏名 Abdellatif Elseoudi Abdellatif Abouelseoud
<p>論 文 題 名 Catabolic effects of FGF-1 on chondrocytes and its possible role in osteoarthritis.</p> <p>軟骨細胞に対するFGF-1の異化促進効果とその変形性関節症における役割</p>		
<p>論文内容の要旨（2000字程度）</p> <p>Fibroblast growth factor 1 (FGF-1) is a classical member of the FGF family and is produced by chondrocytes cultured from osteoarthritic patients. Also, this growth factor was shown to bind to CCN family protein 2 (CCN2), which regenerates damaged articular cartilage and counteracts osteoarthritis (OA) in an animal model. However, the pathophysiological role of FGF-1 in cartilage has not been well investigated. In this study, we evaluated the effects of FGF-1 in vitro and its production in vivo by use of an OA model. Treatment of human chondrocytic cells with FGF-1 resulted in marked repression of genes for cartilaginous extracellular matrix components, whereas it strongly induced MMP-13 (matrix metalloproteinase 13), representing its catabolic effects on cartilage. Interestingly, expression of the CCN2 gene was dramatically repressed by FGF-1, which repression eventually caused the reduced production of CCN2 protein from the chondrocytic cells. The results of a reporter gene assay revealed that this repression could be ascribed, at least in part, to transcriptional regulation. In contrast, the gene expression of FGF-1 was enhanced by exogenous FGF-1, indicating a positive feedback system in these cells. Of note, induction of FGF-1 was observed in the articular cartilage of a rat OA model. These results collectively indicate a pathological role of FGF-1 in OA development, which includes an insufficient cartilage regeneration response caused by CCN2 down-regulation.</p>		