

Free and Open Source Software for Geospatial (FOSS4G) Conference Proceedings

Volume 16 *Bonn, Germany*

Article 19

2016

GRASS in the Desert? Developing FOSS tools for monitoring desertification

Anna Zanchetta
University of Bologna, Italy

Follow this and additional works at: <https://scholarworks.umass.edu/foss4g>

 Part of the [Geographic Information Sciences Commons](#)

Recommended Citation

Zanchetta, Anna (2016) "GRASS in the Desert? Developing FOSS tools for monitoring desertification," *Free and Open Source Software for Geospatial (FOSS4G) Conference Proceedings*: Vol. 16, Article 19.

DOI: <https://doi.org/10.7275/R51V5C55>

Available at: <https://scholarworks.umass.edu/foss4g/vol16/iss1/19>

This Poster is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Free and Open Source Software for Geospatial (FOSS4G) Conference Proceedings by an authorized editor of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

GRASS in the Desert? Developing FOSS tools for monitoring desertification

Anna Zanchetta

University of Bologna, Italy – anna.zanchetta@unibo.it

ABSTRACT:

The use of Change Vector Analysis (CVA) combined with the Tasseled Cap transform (TCT) is a powerful remote sensing tool to monitor forests and vegetated areas, but its application to arid and semiarid environment is not straightforward.

This question is tackled through the calculation of a new set of TCT coefficients using R and GRASS-GIS for SPOT and Landsat satellites, then applied and tested in change detection analysis on a short (seasonal) and a long (decades) temporal scale.

Results show that the combined procedure is an effective method to detect changes in desert environment. Furthermore, the new TCT allows the use of this combined procedure for studies in arid and semi-arid regions, eliminating the doubts on its compatibility with the area of study.

Further development is the creation of a new GRASS-GIS module to performe CVA, thus enabling the simple usage of this technique, until now not available in most common software.

KEYWORDS: Desertification; Change detection; Grass-GIS; R

GRASS IN THE DESERT? Developing FOSS tools for monitoring desertification

A. Zanchetta^a, G. Bitelli^a, A. Karnieli^b
^aDICAM Dep. of Civil, Chemical, Environmental and Materials Engineering
 Bologna University, Viale Risorgimento 2, 40136 Bologna, Italy
^bThe Jacob Blaustein Institutes for Desert Research
 Ben Gurion University of the Negev, Sede Boqer Campus, 8490000 Israel
 anna.zanchetta@unibo.it



OBJECTIVE

The use of **Change Vector Analysis (CVA)** combined with the **Tasseled Cap Transform (TCT)** is a powerful remote sensing tool to monitor forests and vegetated areas, but its application to arid and semi-arid environment is not straightforward.
 This question is tackled through the calculation of a new set of TCT coefficients using **R** and **GRASS-GIS** for SPOT and Landsat satellites, then applied and tested in change detection analysis on a short (seasonal) and a long (decades) temporal scale.
 Further development is the creation of a new GRASS module (**l.cva**) to perform CVA, thus enabling the simple usage of this technique, until now not available in most common software.

CASE STUDY


Azraq Oasis (Jordan) is a good case study since it shows a clear seasonal pattern and was subject in the 90s to the complete drying out of the natural springs.

METHODS


1986-11-04 Landsat 5 look-like image of Fayyum Oasis (Egypt)

Selection of 20 squares of 100 pixels for three categories of surface cover, for different locations and different seasons of the year (at least 6 images, here shown as an example a location in the Sahara desert in Egypt).

Importing the selection in R through Grass, to obtain 2000 pixels for each category.



Calculating TCT through adapted Gram Schmidt orthogonalization for each image then obtaining the final transform by average on location and time.




```
> coefficients
  Band1  Band2  Band3  Band4  Band5  Band6  Band7
Brightness 0.217897 0.273443 0.328729 0.374071 0.482192 0.454707 0.438128
Greenness  0.338888 0.124758 0.080808 -0.156882 0.758528 -0.482472 -0.348724
Wetness    0.258817 0.287834 0.323058 0.429478 -0.332185 -0.486922 -0.388174
```

Creating the adapted Brightness and Greenness maps for the CVA analysis (GRASS scripts in **BASH**).


```
#!/bin/sh
set -e
# Create the adapted Brightness and Greenness maps for the CVA analysis
# Usage: ./l.cva.sh /path/to/your/area /path/to/output
```

Calculating CVA: **ANGLE** and **MAGNITUDE** maps (GRASS scripts).



CHANGE VECTOR ANALYSIS

Obtaining the change detection map by threshold.



Converting the scripts to a new GRASS module.



RESULTS

SPOT satellites

Seasonal change in Azraq Oasis (February – June 2013) detected with the existing TCT [left] and with the new desert-adapted TCT [right].

Landsat satellites

30 years change over Azraq Oasis (1984-2013) during the summer season (August) detected with the new desert-adapted TCT.

CONCLUSION

The Change Vector Analysis (CVA) applied to the new Tasseled Cap Transform (TCT) has been proven to be an effective remote sensing method to detect changes in desert environment.
 In the same time, the newly calculated TCT coefficients allow the use of this combined procedure for studies in arid and semi-arid regions eliminating the doubts on their compatibility with the area of study.
 The integration of the scripts used for the research into the GRASS environment makes this technique easily accessible for other researchers.