

Contents lists available at [ScienceDirect](http://ScienceDirect)

NFS Journal

journal homepage: [www.elsevier.com/locate/nfs](http://www.elsevier.com/locate/nfs)

Original article

## Chemical–physical characteristics, polyphenolic content and total antioxidant activity of three Italian-grown pomegranate cultivars

Roberta Passafiume<sup>a</sup>, Anna Perrone<sup>b</sup>, Giuseppe Sortino<sup>a,\*</sup>, Giuseppe Gianguzzi<sup>a</sup>, Filippo Saletta<sup>a</sup>, Carla Gentile<sup>b</sup>, Vittorio Farina<sup>a</sup><sup>a</sup> Department of Agricultural, Food and Forest Sciences (SAAF), Università degli Studi di Palermo, Viale delle Scienze, Palermo, 90128, Italy<sup>b</sup> Department of Biological, Chemical and Pharmaceutical Sciences and Technologies (STEBICEF), Università degli Studi di Palermo, Viale delle Scienze, Palermo, 90128, Italy

## ARTICLE INFO

## Keywords:

*Punica granatum*  
Fruit quality  
Antioxidant activity  
Phytochemical  
Physico-chemical characteristic  
Bioactive compound content

## ABSTRACT

**Background:** In Mediterranean countries, there is an increasing demand for pomegranate fruits due to their antioxidant properties and nutritional values. The large diffusion of new genotypes and cultivars requires the knowledge of all fruit characteristics in connection with the cultivation area, to satisfy the market demand. This study seeks to determine the fruit quality attributes and nutraceutical values of three pomegranate cultivars (Wonderful, Acco and Kamel) grown in the Mediterranean climate.

**Methods:** The fruits were evaluated for their main physico-chemical traits (weight, volume, height, width, thickness, total arils number, total arils weight, juiciness, fruit index, peel index and total soluble solids content/titratable acidity ratio), bioactive compounds (total phenolic content and total anthocyanin content) and antioxidant activities (radical scavenging activity and ferric reducing power).

**Results:** The data showed that Acco fruit is bigger, juicier and sweeter when compared with the cultivars Wonderful and Kamel. Wonderful is the smallest, least juicy and least sweet but the reddest among all the studied cultivars. Regarding the phenolic compounds and antioxidant activity (radical scavenging activity and ferric reducing power), cultivar Wonderful has the highest values and cultivar Acco contains the most anthocyanin content.

**Conclusions:** This study showed that pomegranate cultivars grown in the Mediterranean area exhibit an appreciable quality, but there are significant differences in quality properties of the arils and juice.

### 1. Introduction

The pomegranate (*Punica granatum* L.) is a tropical and subtropical evergreen shrub that belongs to the Punicaceae family. It is native to the Middle East and central Asia and has been cultivated since antiquity as a fruit crop. The fruit is spherical, with a tough calyx, and covered by a leathery pericarp derived from sepals and adhering floral tissue [1]. The rind encloses many seeds surrounded by juicy arils, which comprise the edible portion of the fruit.

Pomegranate grows well under sunlight, in dry and hot summers, and requires high summer temperatures for the fruit to ripen but, in winter, the plant thrives at minimal temperatures not lower than  $-12\text{ }^{\circ}\text{C}$  [2]. During fruit development, a temperature of  $38\text{ }^{\circ}\text{C}$  and a dry climate, result in the best quality fruit. Conversely, high relative humidity and rain are detrimental to its cultivation [3]. For these reasons, regions of the Mediterranean basin, such as Egypt, Israel, Turkey, Italy,

Spain, Portugal and Anatolia, have suitable climates for pomegranate cultivation [4,5] and show a remarkably high production and consumption of pomegranate fruits. However, pomegranate cultivation has also emerged in parts of Australia, Brazil, Argentina, South Africa and the USA [6,7]. Pomegranate fruit is consumed and sold as whole fresh fruit, juice, syrup, wine, teas and seed oil, among other products, and is valued worldwide for its health benefits [8].

Pomegranate fruit has a long history of use by humans as a food and medicine, demonstrating diverse bioactive properties and nutritional values [9–12,23], mostly associated with its antioxidant characteristics. The antioxidant activity of commercially available pomegranate juice is largely attributed to hydrolysable tannins (notably ellagic acid and derivatives and punicalagin isomers) derived from the rind [13], although other types of polyphenols, including anthocyanins, and ascorbic acid, either in combination or alone, also contribute to the antioxidant activity [16,17]. According to *in vitro* assays, commercial

\* Corresponding author.

E-mail address: [giuseppe.sortino@unipa.it](mailto:giuseppe.sortino@unipa.it) (G. Sortino).<https://doi.org/10.1016/j.nfs.2019.06.001>

Received 5 March 2019; Received in revised form 19 June 2019; Accepted 19 June 2019

Available online 20 June 2019

2352-3646/ © 2019 The Authors. Published by Elsevier GmbH on behalf of Society of Nutrition and Food Science e.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Table 1**  
Monthly mean temperature (°C) and relative humidity (%) and total rainfall (mm) during the three harvest seasons.

Parameter	2015			2016			2017		
	Aug	Sep	Oct	Aug	Sep	Oct	Aug	Sep	Oct
Mean temperature (°C)	25.9	21.6	17.9	24.4	23.1	19.3	24.9	22.8	20.1
Relative humidity (%)	26.3	36.6	43.2	44.4	39.7	48.9	39.8	41.4	45.3
Rain fall (mm)	11.0	21.0	34.0	29.6	16	36.4	12.0	40.0	39.0

pomegranate juice shows three-fold the antioxidant activity of red wine and green tea, and two- to six-fold that of other natural beverages, [14,15]. Cyanidin-3,5-*O*-diglucoside and pelargonidin-3,5-*O*-diglucoside are the most representative anthocyanins in different genotypes of pomegranate, and the predominance of anthocyanins over tannins helps to explain the high reducing activity [26–28]. Moreover, the anthocyanins impart the bright red colour of the juice [18–20], which is one of the major quality parameters appreciated by the consumer.

Clinical studies have revealed that the intake of polyphenols in pomegranate juice improves biochemical parameters, including biomarkers of oxidative stress, protecting against the onset of age-related and degenerative diseases [21,22], as well as chronic diseases related to over-production of free radicals [24,25]. All pomegranate polyphenols (for example, phenolic acids, punicalagins, gallic acid, catechin, chlorogenic acid, caffeic acid, epicatechin, ferulic acid, ellagic acid and kaempferol) show antioxidant activity associated with the indirect inhibition of inflammatory markers, such as tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) [9]. In this context, pomegranates could be considered as a potential natural drug [9].

> 500 different cultivars of pomegranate have been identified [12]. It is crucial to acknowledge all fruit characteristics to select the best cultivars to meet the market demand for quality fruits, consumer preferences, and locality, such as the upper infra-Mediterranean lower sub-humid bioclimatic belt.

Mediterranean pomegranates are based on local cultivars, and their composition can differ from those of Oriental varieties, displaying a large variety of chemical–physical traits and distinct flavour profiles. Meanwhile, modern cultivation is based on cultivars developed by breeding programs for satisfying new markets and consumer needs. The Wonderful cultivar is the most widely planted commercial pomegranate cultivar in Mediterranean countries and is the industry standard variety [29]. This cultivar was imported from the USA about 100 years ago. Its large-sized sweet-and-sour fruit contain red arils and ripen at the beginning of October. The Mediterranean basin recently introduced two additional cultivars: Kamel and Acco. Acco pomegranate is growing in importance, due to its early availability (it can be eaten from the end of summer when the Wonderful is not available), sweet flavour and deep red fruit and skin. Kamel fruit has a dark red skin that appears very early in fruit development, red arils, a sweet-sour flavour and good juice production [12].

Thus, this research reports and compares three years of data concerning the physico-chemical characteristics, antioxidant activity and the major bioactive compound content present in the cultivars Wonderful, Acco and Kamel.

**Table 2**  
Pomological characteristics of the three pomegranate cultivars Acco, Kamel and Wonderful.

Cultivar	Weight (g)	Fruit width (mm)	Fruit height (mm)	Volume (cm <sup>3</sup> )	Peel thickness (mm)	Peel index	Fruit index	Maturity index
Acco	368.3 ± 36.7 a	92.2 ± 4.3 ns	82.3 ± 3.8 a	394.0 ± 53.8 a	3.38 ± 2.1 b	0.01 ± 0.001 a	89.4 ± 5.4 a	0.82 ± 0.02a
Kamel	348.3 ± 54.6 ab	91.0 ± 4.9 ns	76.0 ± 4.6 ab	380.8 ± 57.1 a	7.93 ± 2.2 a	0.02 ± 0.001 ab	83.5 ± 3.3 ab	0.25 ± 0.08 b
Wonderful	289.5 ± 74.0 b	89.9 ± 6.9 ns	69.2 ± 7.4 b	210.8 ± 73.6 b	9.98 ± 1.6 a	0.04 ± 0.01 b	76.9 ± 5.4 b	0.29 ± 0.07 b

Values marked with different letters in the same column indicate significant differences ( $p < .05$ ).

## 2. Materials and methods

### 2.1. Vegetal material

One hundred and fifty fresh pomegranate fruits of Wonderful, Acco and Kamel cultivars (10 fruits × 5 plants × cultivar), were harvested in the years 2015, 2016 and 2017, from a commercial orchard located in Partinico (Palermo, Sicily, Italy). The plants in the research area were spaced 4.0 × 2.5 m, grown from cutting and trained to a vase shape. All fruits were hand-picked at commercial ripening, using skin colour as the maturity index. As mentioned above, Wonderful originated from Florida > 100 years ago, while Acco and Kamel are two new varieties derived from Israel and only recently introduced into the Mediterranean countries. The three studied varieties mature in succession, constituting a continuous offer for the market (Table 1).

According to the estimations provided by the weather station located closest to the research area (189 m above sea level), the average temperatures in this region hover around 18.9 °C, and the average rainfalls are close to 647 mm, with 78 rainy days [30,31]. Under the bioclimatic aspect, the station is referred to as the upper infra-Mediterranean lower sub-humid bioclimatic belt [32]. The monthly mean temperature (°C), relative humidity (%) and total rainfall (mm) during the three harvest seasons (Table 1) provide conditions highly amenable to pomegranate cultivation.

### 2.2. Physico-chemical analyses

To evaluate the pomological and physical–chemical traits of pomegranate fruits, 10 fruit × five plants × cultivar were hand-picked after 15 weeks from flower set (15th October) and analysed for fresh weight (FW), size (SZ), volume (VL), skin colour (SC), juiciness (JUI), juice colour (JC), aril size (AS), aril weight (AW), total soluble solids content (TSSC) and titratable acidity (TA). FW and AW (g) were determined using a digital scale (Gibertini EU–C 2002 RS, Novate Milanese, Italy). Fruit length, width and peel thickness (mm) were measured on the polar axis of the fruit, between the apex and stem, and perpendicular to the polar axis direction, respectively, using a TR53307 digital calliper (Turoni, Forlì, Italy). Fruit volume (cm<sup>3</sup>) was calculated by a liquid displacement method. SC, JC and aril colour were recorded in the CIELab system, using a Minolta CR-400 colorimeter (Konica Minolta Sensing, Inc., Japan). In the CIELab colour space,  $L^*$  indicates the brightness,  $a^*$  ranges from green ( $-a^*$ ) to red ( $+a^*$ ), and  $b^*$  varies from blue ( $-b^*$ ) to yellow ( $+b^*$ ). The colour characteristics of pomegranate cultivars are potential indicators of the maturity index of the fruit, for controlling the harvest and defining the commercial categories of the produce [33,34]. TSSC (°Brix) was measured using an Atago Palette PR–32 digital refractometer (Atago Co., Ltd., Tokyo, Japan). TA was determined by titrating 10 mL of juice with NaOH (0.1 N) to pH 8.1, and the results were expressed as a percentage of citric acid (in grams per litre).

In addition, the maturity index (TSSC/TA), fruit index (longitudinal diameter/transversal diameter × 100) and peel index (peel thickness/fruit weight) were calculated [35]. The peel index is a parameter applicable to sizing the fruit and indicates the typology of usable packaging. Arils from the other fruit tissues were separated, and their juice extracted using a hand-operated citrus squeezer.

**Table 3**  
Arils and juice characteristics of the three pomegranate cultivars Acco, Kamel and Wonderful.


Cultivar	Arils (number)	Arils total weight (g)	Total juice (mL)	Total soluble solid content (°Brix)	Titrateable acidity (%)	Ascorbic acid (mg/100 mL)
Acco	573.3 ± 90.4 ns	184.7 ± 41.6 a	67.3 ± 15.8 a	14.9 ± 3.5 ab	18.1 ± 4.0 c	12.4 ± 2.9 c
Kamel	619.4 ± 108.3 ns	122.9 ± 25.1 b	52.5 ± 14.2 b	12.9 ± 2.7 b	39.2 ± 2.6 b	10.5 ± 1.5 b
Wonderful	552.2 ± 197.2 ns	93.4 ± 20.9 c	38.4 ± 11.1 c	15.0 ± 2.7 a	52.7 ± 9.3 a	12.7 ± 2.3 a

Values marked with different letters in the same column indicate significant differences ( $p < .05$ ); ns = not significant.

**Table 4**  
CIELab colour values of skin, juice and arils of Acco, Kamel and Wonderful fruit.

Cultivar	Skin colour		
	$L^*$	$a^*$	$b^*$
Acco	41.4 ± 13.1 a	40.9 ± 22.1 a	23.0 ± 13.2 a
Kamel	40.0 ± 18.4 a	43.0 ± 10.2 b	11.6 ± 6.3 b
Wonderful	39.1 ± 14.6 a	46.2 ± 12.3 a	20.9 ± 8.6 ab
Cultivar	Arils colour		
	$L^*$	$a^*$	$b^*$
Acco	23.0 ± 13.5 a	15.6 ± 8.9 ab	10.1 ± 3.8 a
Kamel	39.5 ± 14.0 a	36.5 ± 8.7 b	9.03 ± 7.3 a
Wonderful	29.6 ± 9.9 a	24.1 ± 7.4 a	13.7 ± 2.2 a
Cultivar	Juice colour		
	$L^*$	$a^*$	$b^*$
Acco	25.8 ± 10.5 b	60.6 ± 5.04 ab	4.7 ± 5.3 ab
Kamel	40.8 ± 15.2 a	28.8 ± 12.4 ab	12.8 ± 5.8 a
Wonderful	22.5 ± 6.4 b	10.7 ± 6.4 a	2.5 ± 2.3 b

Values marked with different letters in the same column indicate significant differences ( $p < .05$ ).



**Fig. 1.** Colour of skin, juice and arils in Acco, Kamel and Wonderful cultivars recorded in the CIELab colour space and converted to the red–blue–green scale. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

### 2.3. Standards and chemicals

ABTS [2,2'-azinobis(3-ethylbenzothiazoline-6-sulphonic acid)] diammonium salt, Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid), gallic acid (GA), potassium persulphate, 2,4,6-tripyridyl-S-triazine (TPTZ), iron (III) chloride hexahydrate ( $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ ) and Folin–Ciocalteu's phenol reagent were purchased from Sigma Aldrich, St. Louis, <http://www.sigmaaldrich.com>. All other materials and solvents were of analytical grade unless indicated otherwise.

### 2.4. Nutraceutical analysis

The vitamin C content in the ethanol extracts was measured by enzyme colorimetric assay, with a maximum absorbance at 570 nm. The vitamin C concentrations were calculated with reference to a calibration curve constructed using ascorbic acid of increasing concentrations, and expressed as milligrams of ascorbic acid in 100 g of FW.

The total phenolic content (TPC) was determined by the reduction

of phosphotungstic-phosphomolybdic acid (Folin–Ciocalteu's reagent) to a blue pigment in alkaline solution, according to Folin and Denis [36]. Accordingly, 10  $\mu\text{l}$  of the diluted sample was mixed with 4 mL of Folin–Ciocalteu reagent (previously diluted eight-fold), and the mixture was allowed to stand at room temperature for 5 min. Next, 2 mL of 10% ( $w/v$ )  $\text{Na}_2\text{CO}_3$  solution was added, and the volume made to 10 mL with water. After incubation at room temperature in the dark for 90 min, the absorbance was read at 740 nm, using a spectrophotometer (Beckman Coulter DU-800, Fullerton, CA). GA was used to construct the calibration curve, and the results were expressed as milligrams of gallic acid equivalents (GAE) per 100 g FW. All measurements were done in triplicate.

Total anthocyanin content (TAC) of samples was determined using the pH differential method with some modifications [37]. A 20- $\mu\text{l}$  aliquot of each pomegranate juice sample was added separately to 980  $\mu\text{l}$  of KCl buffer (pH 1.0) and NaOAc buffer (pH 4.5). The absorbance was measured at 535 and 700 nm for both sets of pH 1.0 and 4.5 solutions, using 50% ethanol as a blank. The TAC was calculated using Eq. (1), and the results were expressed as milligrams of cyanidin-3-glucoside equivalents in 100 g of FW.

$$\text{Total monomeric anthocyanin} = (A \times \text{MW} \times \text{DF} \times 1000) / \epsilon \times l \quad (1)$$

where  $A$  (absorbance) =  $(A_{515\text{nm}} - A_{700\text{nm}})_{\text{pH } 1.00} - (A_{515\text{nm}} - A_{700\text{nm}})_{\text{pH } 4.5}$ ; MW (molecular weight) = 449.2;  $\epsilon$  (the molar absorptivity of cyanidin-3-glucoside) = 26,900, which was used as the standard; DF is the dilution factor; and  $l$  is the path length.

The total antioxidant activity (TAA) of each cultivar were analysed using the ABTS radical scavenging capacity assay [38] and ferric reducing antioxidant potential (FRAP) assay [39]. These methods are distinguished by their mechanism of action and complement the analysis of the antioxidant potential of the fruits. The ABTS method is based on colorimetric monitoring of the decay of the  $\text{ABTS}^{\cdot+}$  radical cation, caused by the oxidation of  $\text{ABTS}^{\cdot+}$  radicals when contacting an antioxidant.  $\text{ABTS}^{\cdot+}$  was prepared by reacting ABTS with potassium persulphate [40]. The mixture was incubated in the dark at room temperature for at least 16 h and was then diluted with ethanol to an absorbance of  $0.70 \pm 0.05$  at 734 nm. Once prepared, the  $\text{ABTS}^{\cdot+}$  solution (990 ml) was mixed with the aqueous sample (10  $\mu\text{l}$ , diluted 10–40 times), and the absorbance was recorded at 0.0 and 2.5 min. Samples were analysed in triplicate within the linearity range of the assay, as previously described [41].

The FRAP assay is based on the reduction of the  $\text{Fe}^{3+}$ -TPTZ complex to the ferrous form at low pH in the presence of antioxidant compounds [39]. Briefly, 160  $\mu\text{l}$  of FRAP reagent, prepared daily, was mixed with 30  $\mu\text{l}$  of water and 10  $\mu\text{l}$  of the sample (diluted 25–100 times) or standard. After incubation at 37 °C for 30 min, the absorbance at 595 nm was measured. All measurements were repeated three times, in both tests. The TAA evaluated by the ABTS or FRAP protocol was expressed as mmol Trolox equivalents (TE)/100 g FW.

### 2.5. Statistical analysis

Data were presented as mean  $\pm$  standard deviation, and analysed using one-way analysis of variance and Tukey's *post hoc* test at  $p < .05$ . All statistical analyses were conducted using XLSTAT software version 9.0 (Addinsoft, Paris, France). Nutraceuticals data were analysed using




Fig. 2. Total polyphenolic content (TPC, expressed as gallic acid equivalents [GAE, mg] per 100 g fresh weight [FW]) of three different pomegranate cultivars, determined by the Folin–Ciocalteu method. Each value represents the mean  $\pm$  SD ( $n = 3$ ). For each bar, different letters indicate significant differences ( $p < .0001$ ) by Tukey's multiple range test. Letter "a" indicates the highest value.




Fig. 3. Total anthocyanin content (expressed as catechin equivalents [CE, mg] per 100 g fresh weight [FW]) of three different pomegranate cultivars, determined by the pH differential method. Each value represents the mean  $\pm$  SD ( $n = 3$ ). For each bar, different letters indicate significant differences ( $p = .0356$ ) by Tukey's multiple range test. Letter "a" indicates the highest value.

GraphPad Prism v5 software (GraphPad Software Inc., La Jolla, USA).

### 3. Results and discussion

#### 3.1. Pomological characteristics of fruits

The FW varied among the genotypes, decreasing from Acco to Kamel and, finally, Wonderful, which had the lowest FW (Table 2). FW, VL and JC are important marketing attributes because of their influence on consumer preference [19]. Moreover, there is an excellent market for large fruit [12]. There were noticeable differences in the shape of the cultivars; Acco fruit had a spherical form, whereas, both Kamel and Wonderful were obloid. The shape traits showed statistical differences in height but not width.

Pomegranate fruits may be marketed in cardboard boxes and polyurethane trays, based on their size, which is defined by commercial trading practices as count, diameter or weight [43]. In such instances, the package must be labelled accordingly. By weight, the fruit of Wonderful belongs to Class D, whereas, Acco and Kamel are Class C fruit. Considering fruit diameter (width), all the fruits could be

commercialised in Class A.

We observed a greater peel thickness in Wonderful and Kamel fruits than Acco fruit, as confirmed by the peel index (Table 2). For produce of equivalent weights, the market preference is for fruits with a higher proportion of edible content [42]. Acco also presented a higher maturity index (TSSC/TA) when compared with Kamel and Wonderful cultivars (Table 2). This parameter is important because it is responsible for the taste and flavour of the pomegranate.

#### 3.2. Arils and juice characteristics

Pomegranate fruits possess seeds and juice-containing arils (Table 3). Minimally processed pomegranates (ready-to-eat arils) are attracting increasing international demand, due to their desirable characteristics, healthiness, high economic importance and the changing food consumption patterns. Internal parameters measured on pomegranate fruit arils determine the economic value of this fruit and are important factors for growers, market and industry [44]. According to the observed data, the highest number of arils was observed in Kamel (Table 3). The highest AW was observed in Acco, followed by Kamel and, lastly, Wonderful, which displayed both the lowest AW and JC (Table 3). Hence, Acco arils were juicier than those of Kamel and Wonderful. For the TSSC, Wonderful and Acco were similar to each other. Instead, Kamel fruits have a low TSSC, while the percentage of organic acids (TA) is lowest in Acco and highest in Wonderful (Table 3).

As indicated by the colour data (Table 4 and Fig. 1), Wonderful is characterised by intense red peel, juice and arils. Kamel presented a vivid red peel, and bright red juice and arils. Acco showed a dark red peel, with vivid light red arils and juice.

#### 3.3. Bioactive compounds

Acco and Wonderful contained more vitamin C relative to Kamel (Table 3). High TPC values were obtained and simultaneously comparable with other literature data [45,46]. The highest TPC belonged to Wonderful, which was significantly higher in comparison to Kamel and Acco. Moreover, the TPC of Kamel was significantly different from that of Acco (Fig. 2).

The TAC in pomegranate juices of the three varieties analysed was determined using the pH differential method with some modifications [37]. Acco exhibited the highest TAC. However, the only statistically significant difference was observed between Acco and Kamel, as reported in Fig. 3.

Concerning antioxidant activity (ABTS<sup>•+</sup> radical scavenging activity and FRAP) of pomegranate, our results showed significant variability in the nutraceutical potential of the studied cultivars. The mean ABTS<sup>•+</sup> value was 2551 TE/100 g FW. The highest ABTS<sup>•+</sup> scavenging activity was displayed by cultivar Wonderful. The ABTS<sup>•+</sup> activity displayed by cultivar Acco was significantly different from those of Kamel and Wonderful, and cultivar Kamel had a significantly greater ABTS<sup>•+</sup> radical scavenging activity than cultivar Acco (Fig. 4).

Regarding the ability of pomegranate juice to reduce Fe<sup>3+</sup> to Fe<sup>2+</sup>, the highest value was registered in cultivar Wonderful. The mean value was 14,303  $\mu$ mol TE/100 g FW. Wonderful showed a 1.875- and 1.56-fold higher mean FRAP value than those of Acco and Kamel, respectively, and cultivar Kamel registered a significantly higher FRAP value relative to Acco (Fig. 5).

Importantly, our results have shown highly significant correlations between the TAA and FRAP results, TAA and TPC, and, also, between the FRAP and TPC, although the study was conducted only on three cultivars (Table 5). The TAC and TPC were not correlated, and no correlation was observed between the ABTS<sup>•+</sup> radical scavenging activity and FRAP values. According to literature data, a good amount of vitamin C is contained in pomegranates [47], but in our study, the vitamin C content was not correlated with any of the other parameters studied, and polyphenols are fully responsible for the antioxidant






Fig. 4. Total antioxidant activity (TAA, expressed as Trolox equivalents [TE, µmol] per 100 g fresh weight [FW]) of juices of three different pomegranate cultivars, determined by the ABTS assay. Each value represents the mean  $\pm$  SD ( $n = 3$ ). For each bar, different letters indicate significant differences ( $p < .0001$ ) by Tukey's multiple range test. The letter "a" indicates the highest value.




Fig. 5. Ferric reducing antioxidant power (FRAP, expressed as Trolox equivalents [TE, µmol] per 100 g fresh weight [FW]) of juices of three different pomegranate cultivars. Each value represents the mean  $\pm$  SD ( $n = 3$ ). For each bar, different letters indicate significant differences ( $p < .0001$ ) by Tukey's multiple range test. The letter "a" indicates the highest value.

Table 5

Correlation coefficients of the most important characteristics observed in the three studied pomegranate cultivars.

Parameter	VITC	TPC	TAC	TAA	FRAP
VITC	–	–0.159	–0.134	–0.125	0.221
TPC	–0.159	–	–0.675	0.998*	0.997*
TAC	–0.134	–0.6755	–	–0.6278	–0.6162
TAA	–0.125	0.998*	–0.628	–	0.999*
FRAP	0.221	0.997*	–0.616	0.999*	–

VITC (vitamin C), TP (total polyphenolic content), TAC (total anthocyanin content), TAA (total antioxidant activity), FRAP (ferric reducing antioxidant power). Values marked with different letters in the same column indicate significant differences  $p < .0001$  (\*).

activity. To prove this hypothesis, separation and characterisation of the polyphenolic compounds in pomegranate extract by high-performance liquid chromatography–mass spectrometry, with concurrent determination of the vitamin C content are required.

#### 4. Conclusions

This study showed that pomegranate cultivars grown in the Mediterranean area exhibit an appreciable quality, but there are

significant differences in the morphology, as well as the physico-chemical properties of the arils and juice. Thus, great variability exists among these cultivars, according to marketable traits, such as SZ, peel index, JUI, acidity, TSSC, bioactive compounds and *in vitro* antioxidant activity. In particular, our results reveal that Acco produced the biggest and the juiciest fruits, characterised by dark red skin, brilliant arils and light red juice, with the best TSSC/AT ratio and TAC, but the lowest FRAP and TAA. Conversely, Wonderful fruit is smaller in size but has intense red peel, arils and juice, with a low number of arils, which are also less juicy and more acidic than those in cultivar Acco, but the fruit is also characterised by elevated TPC, TAA and FRAP activity. It is well known that the consumer is attracted to the intense colour of fruits. In this context, Wonderful fruit stands out for its intense red peel and juice. Moreover, this cultivar had the highest nutraceutical value. Regarding fruit taste only, Acco could be preferred for its sweet juice. Kamel showed intermediate characteristics between the other two cultivars and the fruit had a brilliant red peel and bright red arils and juice.

Finally, these pomegranate cultivars are selected not just for fresh consumption, but also for the food and health industries, considering the high amount of total polyphenolics, antioxidant potential (ABTS<sup>+</sup> radical scavenging activity and FRAP activity) and TAA. In this context, Wonderful can be emphasised over the cultivars Acco and Kamel. The physico-chemical properties found in this study may be useful in selecting pomegranate genotypes for cultivation, marketability and fruit processing in countries with a Mediterranean climate.

#### Declaration of Competing Interest

The authors declare no conflict of interest.

#### Acknowledgements

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

#### References

- H.Y. Wetzstein, N. Ravid, E. Wilkins, A.P. Martinelli, A morphological and histological characterization of bisexual and male flower types in pomegranate, *J. Am. Soc. Hortic. Sci.* 136 (2011) 83–92, <https://doi.org/10.21273/JASHS.136.2.83>.
- G.M. Levin, Translated from Russian by M. Hopstein, *Pomegranate Roads: A Soviet Botanist's Exile from Eden*, Floreant Press, Forestville, California, 2006.
- A.I. Özgüven, C. Yilmaz, Pomegranate growing in Turkey, *Options Mediterr. Ser. A. (CIHEAM)* 42 (2000) 41–48.
- S. Ercisli, E. Kafkas, E. Orhan, S. Kafkas, Y. Dogan, A. Esitken, Genetic characterization of pomegranate (*Punica granatum* L.) genotypes by AFLP markers, *Biol. Res.* 44 (2011) 345–350 <https://doi.org/10.4067/S0716-97602011000400005>.
- B.E. Ak, A. İkinci, H. Parlakci, A.I. Özgüven, C. Yilmaz, Some pomological traits of different pomegranate varieties grown in Sanliurfa – Turkey, *Acta Hortic.* (818) (2009) 115–119, <https://doi.org/10.17660/ActaHortic.2009.818.15>.
- M. Mars, La culture du grenadier (*Punica granatum* L.) et du figuier (*Ficus carica* L.) en Tunisie, *Cah. Options Mediterr. (CIHEAM)* 13 (1995) 85–95, <https://doi.org/10.3406/jatba.1927.4483>.
- Directory of European institutions holding crop genetic resources collections, in: E.A. Frison, J. Serwinsky (Eds.), *European Cooperative Programme for Crop Genetic Resources Networks (ECP/GR)*, fourth ed., Food and Agriculture Organization (FAO), International Plant Genetic Resources Institute, Rome, Italy, 1995.
- M.K. Sheikh, *The Pomegranate*, IBDC Publishers, Lucknow, India, 2006.
- E.P. Lansky, R.A. Newman, *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer, *J. Ethnopharmacol.* 109 (2007) 177–206, <https://doi.org/10.1016/j.jep.2006.09.006>.
- M.I. Gil, F.A. Tomás-Barberán, B. Hess-Pierce, D.M. Holcroft, A.A. Kader, Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing, *J. Agric. Food Chem.* 48 (2000) 4581–4589, <https://doi.org/10.1021/jf000404a>.
- N.P. Seeram, M. Aviram, Y. Zhang, S.M. Henning, L. Feng, M. Dreher, D. Heber, Comparison of antioxidant potency of commonly consumed polyphenol-rich beverages in the United States, *J. Agric. Food Chem.* 56 (2008) 1415–1422, <https://doi.org/10.1021/jf073035s>.
- D. Holland, K. Hatib, I. Bar-Ya'akov, Pomegranate: botany, horticulture, breeding, *Hortic. Rev.* 35 (2009) 127–191, <https://doi.org/10.1002/9780470593776.ch2>.
- P. Castilla, A. Dávalos, J.L. Teruel, F. Cerrato, M. Fernández-Lucas, J.L. Merino, C.C. Sánchez-Martín, J. Ortuño, M.A. Lasunción, Comparative effects of dietary

- supplementation with red grape juice and vitamin E on production of superoxide by circulating neutrophil NADPH oxidase in hemodialysis patients, *Am. J. Clin. Nutr.* 87 (2008) 1053–1061, <https://doi.org/10.1093/ajcn/87.4.1053>.
- [14] G. Sohrab, J. Nasrollahzadeh, H. Zand, Z. Amiri, M. Tohidi, M. Kimiagar, Effects of pomegranate juice consumption on inflammatory markers in patients with type 2 diabetes: a randomized, placebo-controlled trial, *J. Res. Med. Sci.* 19 (2014) 215–220.
- [15] P. Castilla, R. Echarri, A. Dávalos, F. Cerrato, H. Ortega, J.L. Teruel, M.F. Lucas, D. Gómez-Coronado, J. Ortuño, M.A. Lasunción, Concentrated red grape juice exerts antioxidant, hypolipidemic, and antiinflammatory effects in both hemodialysis patients and healthy subjects, *Am. J. Clin. Nutr.* 84 (2006) 252–262, <https://doi.org/10.1093/ajcn/84.1.252>.
- [16] M.I. Gil, F.A. Tomás-Barberán, B. Hess-Pierce, D.M. Holcroft, A.A. Kader, Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing, *J. Agric. Food Chem.* 48 (2000) 4581–4589, <https://doi.org/10.1021/jf000404a>.
- [17] N.J. Miller, C.A. Rice-Evans, The relative contributions of ascorbic acid and phenolic antioxidants to the total antioxidant activity of orange and apple fruit juices and blackcurrant drink, *Food Chem.* 60 (1997) 331–337, [https://doi.org/10.1016/S0308-8146\(96\)00339-1](https://doi.org/10.1016/S0308-8146(96)00339-1).
- [18] H. Alighourchi, M. Barzegar, Some physicochemical characteristics and degradation kinetic of anthocyanin of reconstituted pomegranate juice during storage, *J. Food Eng.* 90 (2009) 179–185, <https://doi.org/10.1016/j.jfoodeng.2008.06.019>.
- [19] H. Borochoy-Neori, S. Judeinstein, E. Tripler, M. Harari, A. Greenberg, I. Shomer, D. Holland, Seasonal and cultivar variations in antioxidant and sensory quality of pomegranate (*Punica granatum* L.) fruit, *J. Food Compos. Anal.* 22 (2009) 189–195, <https://doi.org/10.1016/j.jfca.2008.10.011>.
- [20] A. Patras, N.P. Brunton, C. O'Donnell, B.K. Tiwari, Effect of thermal processing on anthocyanin stability in foods; mechanisms and kinetics of degradation, *Trends Food Sci. Technol.* 21 (2010) 3–11, <https://doi.org/10.1016/j.tifs.2009.07.004>.
- [21] M.I. Kojadinovic, A.C. Arsic, J.D. Debeljak-Martacic, A.I. Konic-Ristic, N.D. Kardum, T.B. Popovic, M.D. Glibetic, Consumption of pomegranate juice decreases blood lipid peroxidation and levels of arachidonic acid in women with metabolic syndrome, *J. Sci. Food Agric.* 97 (2017) 1798–1804, <https://doi.org/10.1002/jsfa.7977>.
- [22] G. Ekhlasi, F. Shidfar, S. Agah, S. Merat, A.F. Hosseini, Effects of pomegranate and orange juice on antioxidant status in non-alcoholic fatty liver disease patients: a randomized clinical trial, *Int. J. Vitam. Nutr. Res.* 85 (2015) 292–298, <https://doi.org/10.1024/0300-9831/a000292>.
- [23] W. Elfalleh, N. Nasri, N. Marzougui, I. Thabti, A. M'rabet, Y. Yahya, B. Lachiheb, F. Guasmi, A. Ferchichi, Physico-chemical properties and DPPH-ABTS scavenging activity of some local pomegranate (*Punica granatum*) ecotypes, *Int. J. Food Sci. Nutr.* 60 (2009) 197–210, <https://doi.org/10.1080/09637480903067037>.
- [24] B. Hosseini, A. Saedisomeolia, L.G. Wood, M. Yaseri, S. Tavasoli, Effects of pomegranate extract supplementation on inflammation in overweight and obese individuals: a randomized controlled clinical trial, *Complement. Ther. Clin. Pract.* 22 (2016) 44–50, <https://doi.org/10.1016/j.ctcp.2015.12.003>.
- [25] M. Aviram, L. Dornfeld, M. Rosenblat, N. Volkova, M. Kaplan, R. Coleman, T. Hayek, D. Presser, B. Fuhrman, Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL, and platelet aggregation: studies in humans and in atherosclerotic apolipoprotein E-deficient mice, *Am. J. Clin. Nutr.* 71 (2000) 1062–1076, <https://doi.org/10.1093/ajcn/71.5.1062>.
- [26] V. Di Stefano, R. Pitozco, M.E. Novara, D. Bongiorno, S. Indelicato, C. Gentile, G. Avellone, R. Bognanni, S. Scandurra, M.G. Melilli, Antioxidant activity and phenolic composition in pomegranate (*Punica granatum* L.) genotypes from South Italy by UHPLC–Orbitrap–MS approach, *J. Sci. Food Agric.* 99 (2019) 1038–1045, <https://doi.org/10.1002/jsfa.9270>.
- [27] G. Mousavinejad, Z. Emam-Djomeh, K. Rezaei, M.H.H. Khodaparast, Identification and quantification of phenolic compounds and their effects on antioxidant activity in pomegranate juices of eight Iranian cultivars, *Food Chem.* 115 (2009) 1274–1278, <https://doi.org/10.1016/j.foodchem.2009.01.044>.
- [28] C. Kaur, R.K. Pal, A. Kar, C. Gadi, S. Sen, P. Kumar, R. Chandra, S. Jaiswal, I. Khan, Characterization of antioxidants and hypoglycemic potential of pomegranate grown in India: a preliminary investigation, *J. Food Biochem.* 38 (2014) 397–406, <https://doi.org/10.1111/jfbc.12066>.
- [29] A. Goor, J. Liberman, The pomegranate, in: J. Atsmon (Ed.), *The Pomegranate*, Bulletin of the Ministry of Agriculture, 1956, pp. 5–57. Tel Aviv, Israel.
- [30] A. Duro, V. Piccione, C. Scalia, D. Zampino, Precipitazioni e temperature medie mensili in Sicilia relative al sessantennio 1926–1985, in: A. Guerrini (Ed.), *Proceedings of the 5th Workshop Progretta Strategico: Clima, Ambiente e Territorio nel Mezzogiorno*, 28–30 April 1993, Amalfi, Italy, 1996, pp. 17–109.
- [31] A. Drago, *Climatological Atlas of Sicily – second edition*, Riv. Ital. Agrometeorol. 2 (2005) 67–83.
- [32] L. Gianguzzi, F. Papini, D. Cusimano, Phytosociological survey vegetation map of Sicily (Mediterranean region), *J. Maps* 12 (2016) 845–851, <https://doi.org/10.1080/17445647.2015.1094969>.
- [33] A.A. Kader, Increasing food availability by reducing postharvest losses of fresh produce, *Acta Hort.* (682) (2005) 2169–2176, <https://doi.org/10.17660/ActaHortic.2005.682.296>.
- [34] R.B.H. Wills, T.H. Lee, D. Graham, W.B. McGlasson, E.G. Hall, *Postharvest. An Introduction to the Physiology and Handling of Fruit and Vegetables*, Granada, London, UK (1981).
- [35] F. Hernández, P. Melgarejo, F.A. Tomás-Barberán, F. Artés, Evolution of juice anthocyanins during ripening of new selected pomegranate (*Punica granatum*) clones, *Eur. Food Res. Technol.* 210 (1999) 39–42, <https://doi.org/10.1007/s002170050529>.
- [36] V.L. Singleton, J.A. Rossi, Colorimetry of total polyphenols with phosphomolybdic-phosphotungstic acid reagents, *Am. J. Enol. Vitic.* 16 (1965) 144–158.
- [37] M.M. Giusti, P. Jing, Analysis of anthocyanins, in: C. Socaciu (Ed.), *Food Colorants: Chemical and Functional Properties*, CRC Press, Boca Raton, FL, USA, 2007, pp. 429–547.
- [38] N.J. Miller, C.A. Rice-Evans, Spectrophotometric determination of antioxidant activity, *Redox Rep.* 2 (1996) 161–171, <https://doi.org/10.1080/13510002.1996.11747044>.
- [39] I.F.F. Benzie, J.J. Strain, The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay, *Anal. Biochem.* 239 (1996) 70–76, <https://doi.org/10.1006/abio.1996.0292>.
- [40] R. Re, N. Pellegrini, A. Proteggente, A. Pannala, M. Yang, C. Rice-Evans, Antioxidant activity applying an improved ABTS radical cation decolorization assay, *Free Radic. Biol. Med.* 26 (1999) 1231–1237, [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3).
- [41] C. Gentile, A. Perrone, A. Attanzio, L. Tesoriere, M.A. Livrea, Sicilian pistachio (*Pistacia vera* L.) nut inhibits expression and release of inflammatory mediators and reverts the increase of paracellular permeability in IL-1 $\beta$ -exposed human intestinal epithelial cells, *Eur. J. Nutr.* 54 (2015) 811–821, <https://doi.org/10.1007/s00394-014-0760-6>.
- [42] B. Hess-Pierce, A.A. Kader, Responses of 'Wonderful' pomegranates to controlled atmospheres, *Acta Hort.* 600 (2003) 751–757, <https://doi.org/10.17660/ActaHortic.2003.600.115>.
- [43] *Codex Alimentarius Commission, Codex Alimentarius Commission Procedural Manual*, seventeenth ed., World Health Organization (WHO), Food and Agriculture Organization of the United Nations (FAO), Rome, Italy, 2007.
- [44] F. Hernández, P. Legua, R. Martínez, P. Melgarejo, J.J. Martínez, Fruit quality characterization of seven pomegranate accessions (*Punica granatum* L.) grown in southeast of Spain, *Sci. Hortic.* 175 (2014) 174–180, <https://doi.org/10.1016/j.scienta.2014.05.035>.
- [45] J.H.Y. Galani, J.S. Patel, N.J. Patel, J.G. Talati, Storage of fruits and vegetables in refrigerator increases their phenolic acids but decreases the total phenolics, anthocyanins and vitamin C with subsequent loss of their antioxidant capacity, *Antioxidants* 6 (2017) 59, <https://doi.org/10.3390/antiox6030059>.
- [46] B. Bazargani-Gilani, H. Tajik, J. Aliakbarlu, Physicochemical and antioxidative characteristics of Iranian pomegranate (*Punica granatum* L. cv. Rabbab-e-Neyriz) juice and comparison of its antioxidative activity with *Zataria multiflora* Boiss essential oil, *Vet. Res. Forum* 5 (2014) 313–318.
- [47] M. Dżugan, M. Wesołowska, G. Zagula, C. Puchalski, The comparison of the physicochemical parameters and antioxidant activity of homemade and commercial pomegranate juices, *Acta Sci. Pol. Technol. Aliment.* 17 (2018) 59–68, <https://doi.org/10.17306/J.AFS.2018.0529>.