

Supplementary material for the article:

Ilić, B.; Dimkić, I.; Unković, N.; Grbić, M. L.; Vukojević, J.; Vujisić, L.; Tešević, V.; Stanković, S.; Makarov, S.; Lučić, L. Millipedes vs. Pathogens: Defensive Secretions of Some Julids (Diplopoda: Julida) as Potential Antimicrobial Agents. *Journal of Applied Entomology* **2018**, *142* (8), 775–791. <https://doi.org/10.1111/jen.12526>

1 **Supporting Information**

2

3 **FIGURE S1** GC-FID profile of *C. boleti* male and female defensive secretions. Figures B and C  
4 are magnified sections of male secretions chromatogram (A) which show peaks of quinones and  
5 esters, respectively. Figures E and F are magnified sections of female secretions chromatogram  
6 (D) which show peaks of quinones and esters, respectively.

7

8 **FIGURE S2** GC-FID profile of *M. bosniense* male and female defensive secretions. Figures B  
9 and C are magnified sections of male secretions chromatogram (A) which show peaks of  
10 quinones and esters, respectively. Figures E and F are magnified sections of female secretions  
11 chromatogram (D) which show peaks of quinones and esters, respectively.

12

13 **FIGURE S3** GC-FID profile of *M. unilineatum* male and female defensive secretions. Figures B  
14 and C are magnified sections of male secretions chromatogram (A) which show peaks of  
15 quinones and esters, respectively. Figures E and F are magnified sections of female secretions  
16 chromatogram (D) which show peaks of quinones and esters, respectively.

17