

From single-particle excitations to sound waves in a box-trapped atomic Bose-Einstein condensate

Samuel J. Garratt,^{1,2} Christoph Eigen,¹ Jinyi Zhang,¹ Patrik Turzák,¹ Raphael Lopes,^{1,3} Robert P. Smith,^{1,4} Zoran Hadzibabic,¹ and Nir Navon⁵

¹*Cavendish Laboratory, University of Cambridge, J. J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

²*Theoretical Physics, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom*

³*Laboratoire Kastler Brossel, Collège de France, CNRS, ENS-PSL University, UPMC-Sorbonne Universités, 11 Place Marcelin Berthelot, F-75005 Paris, France*

⁴*Clarendon Laboratory, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom*

⁵*Department of Physics, Yale University, New Haven, Connecticut 06520, USA*



(Received 23 October 2018; published 1 February 2019)

We experimentally and theoretically investigate the lowest-lying axial excitation of an atomic Bose-Einstein condensate in a cylindrical box trap. By tuning the atomic density, we observe how the nature of the mode changes from a single-particle excitation (in the low-density limit) to a sound wave (in the high-density limit). Throughout this crossover the measured mode frequency agrees with Bogoliubov theory. Using approximate low-energy models we show that the evolution of the mode frequency is directly related to the interaction-induced shape changes of the condensate and the excitation. Finally, if we create a large-amplitude excitation, and then let the system evolve freely, we observe that the mode amplitude decays nonexponentially in time; this nonlinear behavior is indicative of interactions between the elementary excitations, but remains to be quantitatively understood.

DOI: [10.1103/PhysRevA.99.021601](https://doi.org/10.1103/PhysRevA.99.021601)

I. INTRODUCTION

Low-energy excitations play a central role in our understanding of many-body systems. They characterize a system's low-temperature thermal properties, its response to small perturbations, and its near-equilibrium transport behavior. The collective excitations of ultracold gases have been extensively studied in the traditional setting of a harmonic trap for bosons [1–7] and fermions [8–11] with contact interactions (including low-dimensional gases [12,13]), as well as for Bose-Fermi mixtures [14], spin-orbit-coupled gases [15], and gases with dipolar interactions [16,17].

The recent developments in creating quasiuniform box traps [18–21] have led to intriguing new possibilities. These traps provide a textbook setting for the study of short-wavelength excitations [22], but they also raise new questions on the nature of long-wavelength (system-size) collective modes, as highlighted by recent studies of sound propagation in three-dimensional Bose [23] and Fermi [24] gases, and two-dimensional Bose gases [25] (see also [26–28]). Due to the hard-wall boundary conditions the dynamics depend only on the interplay between kinetic and interaction energy; this is in stark contrast to harmonically trapped gases, where the lowest mode frequency is independent of interaction strength [29].

In this Rapid Communication we experimentally and theoretically study the effect of interactions on the lowest-lying axial mode of a ⁸⁷Rb Bose-Einstein condensate (BEC) confined to a cylindrical box trap. This mode was previously exploited as a route to turbulence in a continuously driven

Bose gas [23]. Here, we vary the atomic density by over two orders of magnitude to probe the near-equilibrium dynamics in both kinetic- and interaction-dominated regimes, and model our system using Bogoliubov theory to show how the mode evolves from a single-particle excitation to a sound wave. We conclude by probing the response of the mode beyond the linear regime, revealing an intriguing nonexponential decay.

II. RESONANT FREQUENCY

Our experiments start with the production of quasipure BECs of between $N = 0.9 \times 10^3$ and 137×10^3 ⁸⁷Rb atoms confined to a cylindrical optical box of length $L = 26(1)$ μm and radius $R = 16(1)$ μm (for details, see [18]). The experimental protocol used to probe the axial mode is shown schematically in Fig. 1(a). After creating the BEC we pulse an axial magnetic field gradient, corresponding to a potential difference $\Delta U = k_B \times 1$ nK over the box length, for a time $\Delta t = 20$ ms, short compared to the period of the mode. We then hold the excited cloud in-trap for a variable time τ before switching off the trap and extracting the cloud's center-of-mass velocity in time of flight. We observe an oscillation of the cloud's velocity with τ , as shown in Fig. 1(b) for $N = 13(1) \times 10^3$ atoms.

If we repeat the same kick protocol with a thermal gas just above the condensation temperature, T_c , we see the same initial velocity as for a quasipure BEC. However, as shown in the inset of Fig. 1(b), there is no subsequent collective oscillation. For a classical gas to support hydrodynamic sound waves,




FIG. 1. Probing the lowest axial mode. (a) Illustration of the experimental protocol. We prepare a BEC of ^{87}Rb in a cylindrical box trap of length L and radius R . Initially, the trapping potential V_{trap} has a flat bottom. We then pulse a magnetic field gradient corresponding to $\Delta U = k_B \times 1$ nK along the box length for $\Delta t = 20$ ms. After a time τ of in-trap evolution we switch off the trap and let the cloud evolve in free space for 140 ms before extracting its center of mass, which reflects the velocity on release from the trap, v_z . (b) $v_z(\tau)$ for a BEC of $N = 13(1) \times 10^3$ atoms. We determine the oscillation frequency ω using a decaying sinusoidal fit. Inset: $v_z(\tau)$ for a noncondensed sample just above T_c , with $N = 10(1) \times 10^3$.

local thermodynamic equilibrium needs to be established on timescales much shorter than the period of the wave [30]. With wavelength $2L$ and speed $\sim \sqrt{k_B T/m}$, where m is the atomic mass, this condition is equivalent to $L \gg 1/(na^2)$; the box length must be much greater than the mean free path. Here a is the s -wave scattering length ($a \approx 100a_0$ for ^{87}Rb , where a_0 is the Bohr radius), $n = N/V$ the atomic number density, and $V = \pi R^2 L$ the volume of the trap. For our thermal gases this condition would be fulfilled only for $N \gg 10^7$ atoms, so hydrodynamic sound waves cannot propagate even in our densest samples. The oscillations we observe in the condensed gas at high density correspond to Bogoliubov sound waves.

In Fig. 2 we summarize the measured condensate oscillation frequencies, and compare them with theories in different interaction regimes. Throughout the paper we model the trap as an infinitely deep cylindrical potential well.

At low density the gas is kinetic energy dominated and we expect ideal-gas behavior. The system is then naturally described in terms of single-particle eigenstates α_j , which are separable in cylindrical coordinates (z, r, ϕ) . The BEC wave function is simply the single-particle ground state $\alpha_0 = \varphi(r) \cos(\pi z/L)$, with $\varphi(R) = 0$, and the lowest axial mode corresponds to $\alpha_1 = \varphi(r) \sin(2\pi z/L)$. The magnetic field gradient appears in the Hamiltonian as a perturbation $\hat{H}_{\text{kick}} = (\Delta U/L)\hat{z}$ for time Δt , and this preferentially excites particles from the condensate into α_1 . The excited state is then a superposition of α_0 and α_1 , which exhibits velocity oscillations at




FIG. 2. Angular frequency of the lowest axial mode as a function of atom number, N . ω_K is the mode frequency in an ideal gas, and ω_I is the frequency of Bogoliubov sound with speed $(gn/m)^{1/2}$ and wavelength $2L$. These calculations use trap dimensions $L = 26 \mu\text{m}$ and $R = 16 \mu\text{m}$. The shaded band ω_B shows the results of numerical Bogoliubov diagonalization (see Sec. III B) accounting for the uncertainty of $\pm 1 \mu\text{m}$ in L and R .

angular frequency

$$\omega_K = \frac{1}{\hbar}(\varepsilon_1 - \varepsilon_0) = \frac{3\hbar}{2m} \left(\frac{\pi}{L} \right)^2, \quad (1)$$

where ε_0 and ε_1 are the single-particle energies, and the subscript K denotes the kinetic-dominated regime. For our trap $\omega_K/(2\pi) = 2.5(2)$ Hz.

In the interaction-dominated regime the condensate wave function is uniform away from the walls, where it decays over the healing length $\xi = (8\pi na)^{-1/2} \ll L$. The lowest axial mode is then a standing sound wave with wavelength $2L$ and speed of sound $(gn/m)^{1/2}$ [31], where $g = 4\pi\hbar^2 a/m$ is the strength of contact interactions. Consequently, at high density the axial mode frequency is

$$\omega_I = \left(\frac{gn}{m} \right)^{1/2} \frac{\pi}{L}. \quad (2)$$

Between these limiting regimes, we capture the crossover with a numerical solution to the Bogoliubov equations (see Sec. III B and [23,32]). In the next section we investigate the physics of this crossover.

III. CROSSOVER

Here we show how the condensate wave function of the interacting Bose gas, and its sound-wave excitations, emerge from the single-particle eigenstates α_j . In second-quantized form, the Hamiltonian is

$$\hat{H} = \underbrace{\sum_j \varepsilon_j \hat{a}_j^\dagger \hat{a}_j}_{\hat{K}} + \frac{g}{2V} \underbrace{\sum_{ijkl} I_{ijkl}^\alpha \hat{a}_i^\dagger \hat{a}_j^\dagger \hat{a}_k \hat{a}_l}_{\hat{I}}, \quad (3)$$

where \hat{K} and \hat{I} are the kinetic- and interaction-energy operators, respectively. \hat{a}_j^\dagger is the creation operator for single-particle eigenstate α_j , and $I_{ijkl}^\alpha/V = \int \alpha_i^* \alpha_j^* \alpha_k \alpha_l d^3r$, where V has been introduced so that I_{ijkl}^α is dimensionless.

A. Condensate

The noninteracting many-body ground state is $|\text{GS}\rangle = (\hat{a}_0^\dagger)^N |0\rangle / \sqrt{N!}$ where $|0\rangle$ is the vacuum of particles. Treating interactions perturbatively, the leading correction to the ground state, $|\delta\text{GS}\rangle$, comes from the operators of the form $\hat{a}_j^\dagger \hat{a}_0^\dagger \hat{a}_0$. At first order

$$|\delta\text{GS}\rangle = \frac{g}{V} \sum_{j \neq 0} I_{j000}^\alpha \frac{(N-1)\sqrt{N}}{\varepsilon_0 - \varepsilon_j} \frac{\hat{a}_j^\dagger \hat{a}_0}{\sqrt{N}} |\text{GS}\rangle, \quad (4)$$

where $\hat{a}_j^\dagger \hat{a}_0 |\text{GS}\rangle / \sqrt{N}$ is the many-body state with one atom excited to single-particle eigenstate α_j , and $(gI_{j000}^\alpha/V)(N-1)\sqrt{N}$ is the matrix element between this many-body state and $|\text{GS}\rangle$. The squared norm of the correction, $\langle \delta\text{GS} | \delta\text{GS} \rangle$, is of order unity for $gn\sqrt{N} = \hbar\omega_K$, at which point perturbation theory fails. Note that in our experiment $gn\sqrt{N} = \hbar\omega_K$ for as few as 10^3 atoms.

In order to develop a physical intuition for the role of the $\hat{a}_j^\dagger \hat{a}_0^\dagger \hat{a}_0$ operators, we contrast our system with a condensate with periodic boundary conditions. In the periodic case the condensate is spatially uniform regardless of interaction strength, and all I_{j000}^α (for $j \neq 0$) vanish because α_j are momentum eigenstates. However, in our case these anomalous operators couple the even-parity eigenstates, thereby changing the condensate shape.

To show this explicitly, we first write the interacting condensate wave function β_0 as a superposition of single-particle eigenstates

$$\beta_i = \sum_j U_{ij}(N) \alpha_j, \quad (5)$$

where U_{ij} is a unitary transformation with a parametric dependence on the interaction strength, in our case captured by N . The states $\beta_{i \neq 0}$, which are orthogonal to each other and to β_0 , will be used in the next section to construct the elementary excitations, but we first focus on the condensate. Working in the β basis

$$\hat{H} = \sum_{ij} \langle \beta_i | \hat{K} | \beta_j \rangle \hat{b}_i^\dagger \hat{b}_j + \frac{g}{2V} \sum_{ijkl} I_{ijkl}^\beta \hat{b}_i^\dagger \hat{b}_j^\dagger \hat{b}_k \hat{b}_l, \quad (6)$$

where \hat{b}_i^\dagger is the creation operator for β_i , and I_{ijkl}^β are the corresponding overlap integrals. Using particle number conservation $\hat{b}_0^\dagger \hat{b}_0 = N - \sum_{i \neq 0} \hat{b}_i^\dagger \hat{b}_i$ we find

$$\hat{H} = \sum_i [(\langle \beta_i | \hat{K} | \beta_0 \rangle + gnI_{i000}^\beta) \hat{b}_i^\dagger \hat{b}_0 + \text{H.c.}] + \hat{H}_2, \quad (7)$$

where we have dropped terms proportional to the identity, \hat{H}_2 is second order (and higher) in $\hat{b}_{i \neq 0}$ operators, and H.c. denotes the Hermitian conjugate. If the ground state of \hat{H} has a large condensate fraction, $\langle \hat{b}_0^\dagger \hat{b}_0 \rangle \gg \langle \hat{b}_i^\dagger \hat{b}_i \rangle$ for $i \neq 0$, we must have

$$\langle \beta_i | \hat{K} | \beta_0 \rangle + gnI_{i000}^\beta = 0, \quad (8)$$

for all $i \neq 0$; as N increases the condensate wave function follows a path in the space of single-particle eigenstates along which it is decoupled from all orthogonal states.




FIG. 3. (a) Axial density profile of the BEC, for three interaction strengths $gn/(\hbar\omega_K)$. (b) Change in the axial density profile due to the coherently occupied axial mode, scaled by $\sqrt{\omega}$ so the amplitude for $gn/(\hbar\omega_K) \gg 1$ does not depend on N .

If the set of β_i forms a complete basis, then Eq. (8) implies

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + gn|\beta_0|^2 \right) \beta_0 = \mu \beta_0, \quad (9)$$

for a constant μ , which is identified as the Hartree-Fock chemical potential. Equation (9) is then the well-known Gross-Pitaevskii (GP) equation [31], which has arisen from the sole assumption that one state, β_0 , has much greater occupation than those orthogonal to it.

In practice, we work with a truncated basis of single-particle eigenstates. Minimizing the GP energy functional (with respect to U_{0j}) within this truncated set gives the condensate wave function β_0 that is not an exact solution to the GP equation, but does satisfy Eq. (8).

In Fig. 3(a) we illustrate how the condensate shape changes in the crossover between kinetic- and interaction-energy dominated regimes.

B. Excitations

To study the evolution of excitations with interaction strength we use Bogoliubov theory. Having determined β_0 variationally, we construct the set of $\beta_{i \neq 0}$ using the Gram-Schmidt procedure, then introduce the mean field $\hat{b}_0 = \sqrt{N}$ in Eq. (6). As in conventional Bogoliubov theory, we neglect terms cubic (and higher) in $\hat{b}_{i \neq 0}$ operators, thereby arriving at an effective quadratic Hamiltonian for the near-equilibrium dynamics. This is diagonalized using a bosonic transformation [33]

$$\hat{c}_i = \sum_{j=1} (P_{ij} \hat{b}_j + Q_{ij} \hat{b}_j^\dagger), \quad (10)$$

where P and Q are chosen such that

$$\hat{H} \approx \hat{H}_B = \sum_{j=1} \hbar\omega_j \hat{c}_j^\dagger \hat{c}_j \quad (11)$$

is the Bogoliubov Hamiltonian, where we have omitted the energy of the interacting ground state and \hat{c}_j^\dagger is the creation operator for the j th normal mode. Although the Gram-Schmidt procedure does not uniquely specify $\beta_{i \neq 0}$, P_{ij} and Q_{ij} adjust accordingly to uniquely specify \hat{c}_i (up to a phase factor).

In Fig. 3(b) we show how the axial density profile of the lowest-lying axial mode changes through the crossover. Note that the wavelength of the mode in the interaction-dominated regime is double that in the ideal gas.

Before comparing our theoretical results with the experiments, it remains to be shown that throughout the crossover

the axial kick \hat{H}_{kick} leads to velocity oscillations at the frequency of the lowest axial mode. Here we show that this is indeed the case, and that the excited state has a coherent occupation of normal modes. Approximating $\hat{b}_0 = \sqrt{N}$ and assuming $N \gg 1$, we express \hat{H}_{kick} as a linear combination of normal mode operators

$$\begin{aligned}\hat{H}_{\text{kick}} &\approx \frac{\Delta U}{L} \sqrt{N} \sum_i \langle \beta_0 | \hat{z} | \beta_i \rangle \hat{b}_i + \text{H.c.} \\ &= \frac{\Delta U}{L} \sqrt{N} \sum_j Z_j \hat{c}_j + \text{H.c.},\end{aligned}\quad (12)$$

where Z_j is given by the P and Q matrices [34]. We then treat \hat{H}_{kick} as a perturbation to \hat{H}_B and find the time-evolution operator in the interaction picture,

$$\hat{U}(\Delta t) = \prod_j \mathcal{T} \exp \left[-i \int_0^{\Delta t} (\eta_j e^{-i\omega_j t'} \hat{c}_j + \text{H.c.}) dt' \right], \quad (13)$$

where $\eta_j \equiv \Delta U \sqrt{N} Z_j / (\hbar L)$ and \mathcal{T} denotes time ordering. For the low-energy modes $e^{-i\omega_j \Delta t} \approx 1$ so the time-ordering operation is trivial. To obtain the state at the end of the kick we apply $\hat{U}(\Delta t)$ to the ground state of \hat{H}_B , $|\text{GS}\rangle_B$, which yields

$$|\eta\rangle \equiv \hat{U}(\Delta t) |\text{GS}\rangle_B = \prod_j e^{-|\eta_j \Delta t|^2 / 2} e^{-i\eta_j \Delta t \hat{c}_j^\dagger} |\text{GS}\rangle_B, \quad (14)$$

a coherent occupation of normal modes. Following the kick, the in-trap time evolution is generated by \hat{H}_B . To calculate the velocity of the cloud we first write the momentum operator in terms of the normal modes. Following the same procedure as in Eq. (12),

$$\hat{p}_z \approx \hbar \sqrt{N} \sum_j D_j \hat{c}_j + \text{H.c.}, \quad (15)$$

where $\hbar D_j$ is analogous to Z_j [35]. The axial velocity is then

$$\begin{aligned}\langle v_z(\tau) \rangle &= \frac{1}{Nm} \langle \eta | e^{i\hat{H}_B \tau / \hbar} \hat{p}_z e^{-i\hat{H}_B \tau / \hbar} | \eta \rangle \\ &= 2 \frac{\Delta t}{m} \frac{\Delta U}{L} \sum_j \text{Im} \{ D_j Z_j e^{-i\omega_j \tau} \}.\end{aligned}\quad (16)$$

The contribution of the j th mode to the velocity amplitude therefore scales as $|Z_j D_j|$. We find numerically that the lowest axial mode contributes 96% of the amplitude in our most dilute samples, falling to 86% in our densest ones. Moreover, as the excitation spectrum is discrete, any significant population of higher modes would cause the velocity oscillations to be visibly nonsinusoidal, which we do not observe experimentally. All of this confirms that the oscillations that we observe [see Fig. 1(b)] arise as a result of the direct coupling to the lowest-axial mode.

C. Truncated-basis models

To understand the effect of interactions on the axial mode frequency throughout the crossover, we calculate it using (progressively larger) truncated sets of low-energy single-particle eigenstates.

First, we consider just the two lowest single-particle eigenstates (the lowest even-parity state α_0 and the lowest




FIG. 4. Crossover from single-particle excitations to sound waves. We plot ω/ω_K as a function of the interaction strength, now parametrized by $gn/(\hbar\omega_K)$. Here we assume $L = 26 \mu\text{m}$ and $R = 16 \mu\text{m}$ for calculations. The $x-y$ error bar in the bottom right corner indicates the fractional systematic uncertainties in the experimental data due to the uncertainties in the box dimensions. $\omega^{(2)}$ is determined using Bogoliubov theory within a truncated basis of just the two lowest-energy single-particle eigenstates of zero angular momentum (see text). This scheme fails even for relatively small $gn/(\hbar\omega_K)$, as it does not allow for the interaction-induced changes in the shape of the condensate or the excitation mode. $\omega^{(5)}$, based on a truncated basis of five single-particle eigenstates, is the minimal model that allows for the shape changes. This simple model already captures most of the crossover, and using progressively larger truncated bases does not qualitatively change the result, as shown by $\omega^{(15)}$, which is based on 15 single-particle eigenstates. ω_I is the sound-wave frequency, approached in the limit of large $gn/(\hbar\omega_K)$.

odd-parity state α_1). In this case the condensate wave function is $\beta_0 = \alpha_0$, the mode involves only α_1 , and neither has the freedom to change its shape with increasing $gn/(\hbar\omega_K)$. Here the bosonic transform in Eq. (10) gives $\hat{c}_1 = \cosh(\kappa)\hat{a}_1 + \sinh(\kappa)\hat{a}_1^\dagger$, where κ is chosen to diagonalize \hat{H}_B [see Eq. (11)]. The resulting mode frequency is

$$\omega^{(2)} = \omega_K \sqrt{\left(1 + \frac{\mathcal{J}gn}{3\hbar\omega_K}\right)^2 - \left(\frac{2\mathcal{J}gn}{3\hbar\omega_K}\right)^2}, \quad (17)$$

where $\mathcal{J} \approx 2.10$ is the radial factor in the overlap integral I_{0011}^α . As shown in Fig. 4, this scheme fails to describe the dynamics even for values of $gn/(\hbar\omega_K)$ well below unity.

As a minimal model that does allow for the interaction-induced shape changes of both the condensate and the excitation mode, we consider a truncated set of the five lowest-energy α_j with zero angular momentum, and calculate the corresponding $\omega^{(5)}$. In this case β_0 is a superposition of three even-parity states, and the excited mode involves two of odd parity; note that here we use five states because the third-lowest even state has a lower energy than the second-lowest odd one. We find that this simple model is sufficient to capture rather well most of the crossover, up to $gn/(\hbar\omega_K) \approx 3$ (see Fig. 4), where gn exceeds the maximum kinetic energy in the truncated set. For any $gn/(\hbar\omega_K)$ significantly larger than 3, the simple sound-wave calculation ω_I already provides a good approximation.




FIG. 5. Nonlinear damping of the lowest-lying axial mode in the interaction-dominated regime, with fixed density n_0 such that $gn_0 = k_B \times 2.1$ nK [corresponding to $gn_0/(\hbar\omega_K) \approx 17$]. (a) Velocity oscillations following kicks with $\Delta U = k_B \times 0.3$ nK (left) and $k_B \times 3.6$ nK (right). The solid lines show exponentially decaying sinusoidal functions obtained by fitting to the early-time data, $\tau < 0.25$ s. For large ΔU (right panel) the decay is clearly nonexponential. (b) Initial decay rate Γ_i as a function of the normalized kick amplitude $\Delta U/(gn_0)$. The solid line is a linear fit to data with $\Delta U/(gn_0) < 1$ [the dashed line displays its extrapolation for $\Delta U/(gn_0) > 1$]. Inset: frequency of the damped oscillation ω normalized to the low- ΔU value, ω_0 .

The agreement of the Bogoliubov calculations with the experimental data improves further as we consider ever larger basis sets, as shown by the 15-state calculation, $\omega^{(15)}$, in Fig. 4, and the full numerical result in Fig. 2; there ω_B was calculated using 80 single-particle states with kinetic energies up to $\approx 70\hbar\omega_K$. However, the success of the simple $\omega^{(5)}$ calculation highlights the key qualitative message: most of the physics of the crossover is captured by including, at lowest order, the interaction-induced shape changes that arise due to the experimentally relevant fixed boundary conditions.

IV. BEYOND LINEAR RESPONSE

We have so far neglected any coupling between the normal modes of the BEC. However, we observe at least a weak damping in all the frequency measurements summarized in Figs. 2 and 4 [see, for instance, Fig. 1(b)]. In general, the nonzero temperature of our gases will lead to Landau damping [36], but our conservative upper bound of $T < 10$ nK suggests a decay rate of $\Gamma_{\text{Landau}}/(2\pi) < 0.2$ s $^{-1}$, much smaller than observed.

Here we examine this damping for different kick amplitudes ΔU , focusing on the interaction-dominated regime with $gn/(\hbar\omega_K) \approx 17$; we fix gn to $gn_0 = k_B \times 2.1(2)$ nK by fixing the atom number to $N = 1.2(1) \times 10^5$. Figure 5(a) shows $v_z(\tau)$ following kicks with $\Delta U = k_B \times 0.3$ nK (left panel) and $k_B \times 3.6$ nK (right panel). For the weak kick only a subtle damping is observed, and an exponentially decaying sine (solid line), fit to the early-time data ($\tau < 0.25$ s), captures the data well for all τ . However, for the stronger kick we clearly see a rapid initial decay followed by a long-lived oscillation. Here the solid line, based on the same fitting to the early-time data ($\tau < 0.25$ s), clearly fails to capture the oscillations for $\tau \gtrsim 0.4$ s.

We characterize the damping using the initial velocity-decay rate, Γ_i , extracted from the early-time ($\tau < 0.25$ s) fits. In Fig. 5(b) we show Γ_i versus normalized kick amplitude, $\Delta U/(gn_0)$, and in the inset we show that the mode frequency is approximately constant across our whole range of ΔU . For relatively weak kicks ($\Delta U \lesssim gn_0$) the damping rate appears to be linear in kick amplitude, essentially vanishing (within experimental errors) as $\Delta U \rightarrow 0$. This diverging lifetime in the limit of vanishing excitation amplitude is consistent with the absence of lower-lying modes to which this mode could readily decay, and moreover it excludes (at the level of our experimental errors) damping due to nonzero temperature or technical reasons. Both the nonexponential decay and the fact that Γ_i increases with ΔU suggest that the damping occurs due to interactions between the excitations.

V. CONCLUSIONS AND OUTLOOK

We have measured the dynamics of an atomic BEC in a cylindrical box trap following an axial kick, thereby probing its lowest axial mode. By tuning the gas density we studied the evolution from single-particle to many-body dynamics. We used a simple model to elucidate the effect of interactions, and numerically evaluated the mode frequency over the whole range of densities, finding excellent agreement with the experiments. Going beyond linear response in the interaction-dominated regime, we observed a nonexponential decay of the excitation, hinting at a nonlinear many-body decay mechanism. A future challenge is to understand this decay mechanism, and in particular its dependence on the interaction strength.

ACKNOWLEDGMENTS

We are grateful to F. Werner and J. T. Chalker for comments on the manuscript. This work was supported by EPSRC (Grants No. EP/N011759/1 and No. EP/P009565/1), ERC (QBox), AFOSR, and ARO. N.N. acknowledges support from Trinity College (Cambridge) and the David and Lucile Packard Foundation. R.L. acknowledges support from the E.U. Marie-Curie program (Grant No. MSCA-IF-2015 704832) and Churchill College, Cambridge. R.P.S. acknowledges support from the Royal Society.

- [1] D. S. Jin, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Collective Excitations of a Bose-Einstein Condensate in a Dilute Gas, *Phys. Rev. Lett.* **77**, 420 (1996).
- [2] M. O. Mewes, M. R. Andrews, N. J. van Druten, D. M. Kurn, D. S. Durfee, C. G. Townsend, and W. Ketterle, Collective Excitations of a Bose-Einstein Condensate in a Magnetic Trap, *Phys. Rev. Lett.* **77**, 988 (1996).
- [3] M. R. Andrews, D. M. Kurn, H.-J. Miesner, D. S. Durfee, C. G. Townsend, S. Inouye, and W. Ketterle, Propagation of Sound in a Bose-Einstein Condensate, *Phys. Rev. Lett.* **79**, 553 (1997).
- [4] O. M. Maragò, S. A. Hopkins, J. Arlt, E. Hodby, G. Hechenblaikner, and C. J. Foot, Observation of the Scissors Mode and Evidence for Superfluidity of a Trapped Bose-Einstein Condensed Gas, *Phys. Rev. Lett.* **84**, 2056 (2000).
- [5] J. Steinhauer, R. Ozeri, N. Katz, and N. Davidson, Excitation Spectrum of a Bose-Einstein Condensate, *Phys. Rev. Lett.* **88**, 120407 (2002).
- [6] R. Meppelink, S. B. Koller, and P. van der Straten, Sound propagation in a Bose-Einstein condensate at finite temperatures, *Phys. Rev. A* **80**, 043605 (2009).
- [7] D. S. Lobser, A. E. S. Barentine, E. A. Cornell, and H. J. Lewandowski, Observation of a persistent non-equilibrium state in cold atoms, *Nat. Phys.* **11**, 1009 (2015).
- [8] M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. H. Denschlag, and R. Grimm, Collective Excitations of a Degenerate Gas at the BEC-BCS Crossover, *Phys. Rev. Lett.* **92**, 203201 (2004).
- [9] J. Joseph, B. Clancy, L. Luo, J. Kinast, A. Turlapov, and J. E. Thomas, Measurement of Sound Velocity in a Fermi Gas near a Feshbach Resonance, *Phys. Rev. Lett.* **98**, 170401 (2007).
- [10] S. Nascimbène, N. Navon, K. J. Jiang, L. Tarruell, M. Teichmann, J. McKeever, F. Chevy, and C. Salomon, Collective Oscillations of an Imbalanced Fermi Gas: Axial Compression Modes and Polaron Effective Mass, *Phys. Rev. Lett.* **103**, 170402 (2009).
- [11] L. A. Sidorenkov, M. K. Tey, R. Grimm, Y.-H. Hou, L. Pitaevskii, and S. Stringari, Second sound and the superfluid fraction in a Fermi gas with resonant interactions, *Nature (London)* **498**, 78 (2013).
- [12] H. Moritz, T. Stöferle, M. Köhl, and T. Esslinger, Exciting Collective Oscillations in a Trapped 1D Gas, *Phys. Rev. Lett.* **91**, 250402 (2003).
- [13] E. Vogt, M. Feld, B. Fröhlich, D. Pertot, M. Koschorreck, and M. Köhl, Scale Invariance and Viscosity of a Two-Dimensional Fermi Gas, *Phys. Rev. Lett.* **108**, 070404 (2012).
- [14] I. Ferrier-Barbut, M. Delehaye, S. Laurent, A. T. Grier, M. Pierce, B. S. Rem, F. Chevy, and C. Salomon, A mixture of Bose and Fermi superfluids, *Science* **345**, 1035 (2014).
- [15] J.-Y. Zhang, S.-C. Ji, Z. Chen, L. Zhang, Z.-D. Du, B. Yan, G.-S. Pan, B. Zhao, Y.-J. Deng, H. Zhai, S. Chen, and J.-W. Pan, Collective Dipole Oscillations of a Spin-Orbit Coupled Bose-Einstein Condensate, *Phys. Rev. Lett.* **109**, 115301 (2012).
- [16] G. Bismut, B. Pasquiou, E. Maréchal, P. Pedri, L. Vernac, O. Gorceix, and B. Laburthe-Tolra, Collective Excitations of a Dipolar Bose-Einstein Condensate, *Phys. Rev. Lett.* **105**, 040404 (2010).
- [17] L. Chomaz, R. M. W. van Bijnen, D. Petter, G. Faraoni, S. Baier, J. H. Becher, M. J. Mark, F. Wächtler, L. Santos, and F. Ferlaino, Observation of roton mode population in a dipolar quantum gas, *Nat. Phys.* **14**, 442 (2018).
- [18] A. L. Gaunt, T. F. Schmidutz, I. Gotlibovych, R. P. Smith, and Z. Hadzibabic, Bose-Einstein Condensation of Atoms in a Uniform Potential, *Phys. Rev. Lett.* **110**, 200406 (2013).
- [19] L. Chomaz, L. Corman, T. Bienaimé, R. Desbuquois, C. Weitenberg, S. Nascimbène, J. Beugnon, and J. Dalibard, Emergence of coherence via transverse condensation in a uniform quasi-two-dimensional Bose gas, *Nat. Commun.* **6**, 6162 (2015).
- [20] B. Mukherjee, Z. Yan, P. B. Patel, Z. Hadzibabic, T. Yefsah, J. Struck, and M. W. Zwierlein, Homogeneous Atomic Fermi Gases, *Phys. Rev. Lett.* **118**, 123401 (2017).
- [21] K. Hueck, N. Luick, L. Sobirey, J. Siegl, T. Lompe, and H. Moritz, Two-Dimensional Homogeneous Fermi Gases, *Phys. Rev. Lett.* **120**, 060402 (2018).
- [22] R. Lopes, C. Eigen, A. Barker, K. G. H. Viebahn, M. Robert-de-Saint-Vincent, N. Navon, Z. Hadzibabic, and R. P. Smith, Quasiparticle Energy in a Strongly Interacting Homogeneous Bose-Einstein Condensate, *Phys. Rev. Lett.* **118**, 210401 (2017).
- [23] N. Navon, A. L. Gaunt, R. P. Smith, and Z. Hadzibabic, Emergence of a turbulent cascade in a quantum gas, *Nature (London)* **539**, 72 (2016).
- [24] M. Zwierlein (private communication).
- [25] J. L. Ville, R. Saint-Jalm, É. Le Cerf, M. Aidelsburger, S. Nascimbène, J. Dalibard, and J. Beugnon, Sound Propagation in a Uniform Superfluid Two-Dimensional Bose Gas, *Phys. Rev. Lett.* **121**, 145301 (2018).
- [26] M. Ota and S. Stringari, Second sound in a two-dimensional Bose gas: From the weakly to the strongly interacting regime, *Phys. Rev. A* **97**, 033604 (2018).
- [27] M. Ota, F. Larcher, F. Dalfovo, L. Pitaevskii, N. P. Proukakis, and S. Stringari, Collisionless Sound in a Uniform Two-Dimensional Bose Gas, *Phys. Rev. Lett.* **121**, 145302 (2018).
- [28] A. Cappellaro, F. Toigo, and L. Salasnich, Collisionless dynamics in two-dimensional bosonic gases, *Phys. Rev. A* **98**, 043605 (2018).
- [29] W. Kohn, Cyclotron Resonance and de Haas-van Alphen Oscillations of an Interacting Electron Gas, *Phys. Rev.* **123**, 1242 (1961).
- [30] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, UK, 1995).
- [31] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superfluidity* (Oxford University Press, New York, 2016).
- [32] ω_I and ω_B are distinguishable even for our highest N because the ω_I calculation assumes a strictly uniform BEC density over the trap, which is never exact due to the nonzero healing length. The small discrepancy between the data and the Bogoliubov solution likely arises because in the experiment the trap walls are not infinitely steep [18], so the effective trap volume slightly increases with N .
- [33] J.-P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (MIT Press, Cambridge, MA, 1986).
- [34] Defining the inverse matrices P^{-1} and Q^{-1} via $\hat{b}_i = \sum_j [P_{ij}^{-1} \hat{c}_j + Q_{ij}^{-1} \hat{c}_j^\dagger]$, we have $Z_j = \sum_i [\langle \beta_0 | \hat{z} | \beta_j \rangle P_{ij}^{-1} + \langle \beta_j | \hat{z} | \beta_0 \rangle (Q_{ij}^{-1})^*]$.
- [35] Using the inverse matrices P^{-1} and Q^{-1} , we have $\hbar D_j = \sum_i [\langle \beta_0 | \hat{p}_z | \beta_j \rangle P_{ij}^{-1} + \langle \beta_j | \hat{p}_z | \beta_0 \rangle (Q_{ij}^{-1})^*]$.
- [36] L. P. Pitaevskii and S. Stringari, Landau damping in dilute Bose gases, *Phys. Lett. A* **235**, 398 (1997).