



RESEARCH ARTICLE

Distinct telomere differences within a reproductively bimodal common lizard population

Darryl McLennan^{1,2} | Hans Recknagel¹ | Kathryn R. Elmer¹ | Pat Monaghan¹

¹Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow, UK

²Department of Fish Ecology and Evolution, EAWAG, Kastanienbaum, Switzerland

Correspondence

Darryl McLennan
Email: darrylmclennan@outlook.com

Funding information

Natural Environment Research Council, Grant/Award Number: NE/N003942/1; H2020 European Research Council, Grant/Award Number: 322784

Handling Editor: Frédéric Angelier

Abstract

1. Different strategies of reproductive mode, either oviparity (egg-laying) or viviparity (live-bearing), will be associated with a range of other life-history differences that are expected to affect patterns of ageing and longevity. It is usually difficult to compare the effects of alternative reproductive modes because of evolutionary and ecological divergence. However, the very rare exemplars of reproductive bimodality, in which different modes exist within a single species, offer an opportunity for robust and controlled comparisons.
2. One trait of interest that could be associated with life history, ageing and longevity is the length of the telomeres, which form protective caps at the chromosome ends and are generally considered a good indicator of cellular health. The shortening of these telomeres has been linked to stressful conditions; therefore, it is possible that differing reproductive costs will influence patterns of telomere loss. This is important because a number of studies have linked a shorter telomere length to reduced survival.
3. Here, we have studied maternal and offspring telomere dynamics in the common lizard (*Zootoca vivipara*). Our study has focused on a population where oviparous and viviparous individuals co-occur in the same habitat and occasionally interbreed to form admixed individuals.
4. While viviparity confers many advantages for offspring, it might also incur substantial costs for the mother, for example require more energy. Therefore, we predicted that viviparous mothers would have relatively shorter telomeres than oviparous mothers, with admixed mothers having intermediate telomere lengths. There is thought to be a heritable component to telomere length; therefore, we also hypothesized that offspring would follow the same pattern as the mothers.
5. Contrary to our predictions, the viviparous mothers and offspring had the longest telomeres, and the oviparous mothers and offspring had the shortest telomeres. The differing telomere lengths may have evolved as an effect of the life-history divergence between the reproductive modes, for example due to the increased

Darryl McLennan and Hans Recknagel contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Functional Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society

growth rate that viviparous individuals may undergo to reach a similar size at reproduction.

KEYWORDS

life-history variation, oviparity, physiological state, squamate, viviparity

1 | INTRODUCTION

Differing reproductive strategies can co-occur within a species, due to both environmental and genetic differences among individuals, as well as the interaction between the two (Taborsky & Brockmann, 2010). This phenotypic variation can exist on a continuous spectrum or fall into discontinuous life-history modes (Partridge, MacManes, Knapp, & Neff, 2016; Tsubaki, Hooper, & Siva-Jothy, 1997), and strategies can be plastic or fixed within an individual's lifetime (Bailey, Gray, & Zuk, 2010; Baum, Laughton, Armstrong, & Metcalfe, 2004; Bronikowski & Arnold, 1999; Meunier et al., 2012; Zamudio & Chan, 2008).

A striking example of fixed, discontinuous reproductive modes is when a species is bimodal for both oviparity (i.e. egg-laying) and viviparity (i.e. giving birth to live young). Viviparity is evolutionarily derived from oviparity and confers many benefits to offspring, particularly a more stable and protected developmental environment in the mother's reproductive tract (Shine, 1995). However, viviparity can also restrict a mother's reproductive output and may reduce the number of clutches produced within a given year (Blackburn, 1999; Recknagel & Elmer, 2019; Sites, Reeder, & Wiens, 2011; Wourms & Lombardi, 1992). Most vertebrate species exhibit only one of these reproductive modes; however, in a small number of species, both reproductive modes occur (Murphy & Thompson, 2011). The differing maternal costs associated with each reproductive mode have been little studied, in part because so few species exhibit both modes. However, it has been suggested that viviparity may incur substantial costs for the mother. Firstly, the prolonged gestational period and presumed larger clutch mass in viviparous individuals (Horváthová et al., 2013; Qualls & Shine, 1998; Roitberg et al., 2013) could carry a greater metabolic cost for the mother. Secondly, viviparity has been linked to the evolution of larger body sizes to counteract the space constraint *in utero* (Qualls & Shine, 1995) and there can be costs associated with rapid growth and/or a larger body size (Metcalfe & Monaghan, 2003).

It is likely that such a divergence in reproductive mode will also be associated with other life-history differences, which could in turn affect patterns of ageing and longevity. One trait that could be of interest in this context is telomere length. Telomeres cap the ends of eukaryotic chromosomes and play an important role in chromosome protection (Blackburn, 1991; Monaghan, 2010). These telomere caps shorten with each round of cell division because of the 'end replication problem' (Levy, Allsopp, Futcher, Greider, & Harley, 1992). Certain species are capable of telomere elongation, mostly via the expression of the enzyme telomerase (Gomes, Shay, & Wright, 2010;

Tian et al., 2018). However, in the absence of elongation mechanisms, telomeres may shorten to such an extent that the central coding region of the chromosome becomes vulnerable. As such, a relatively short telomere length is considered to be an indicator of poor cellular and biological state, and a number of studies have linked a shorter telomere length and/or a faster rate of telomere attrition to reduced survival and/or longevity (Boonekamp, Mulder, Salomons, Dijkstra, & Verhulst, 2014; Debes, Visse, Panda, Ilmonen, & Vasemagi, 2016; Dupoué et al., 2017; Salmón, Nilsson, Nord, Bensch, & Isaksson, 2016; Wilbourn et al., 2018).

In addition to cell division, the accelerated erosion of telomeres has been linked to environmental stressors, potentially via oxidative stress pathways (Monaghan & Ozanne, 2018; Reichert & Stier, 2017, but see also Boonekamp, Bauch, Mulder, & Verhulst, 2017). Recent studies have linked telomere dynamics to various stressful conditions, both *in vitro* and *in vivo* (Barnes, Fouquerel, & Opresko, 2018; Cram, Monaghan, Gillespie, & Clutton-Brock, 2017; Debes et al., 2016; Monaghan, 2014; Olsson et al., 2018). Moreover, a recent study on Australian painted dragons (*Ctenophorus pictus*) found that telomere dynamics differed between individuals with different reproductive and life-history tactics (Rollings et al., 2017). It is therefore possible that differing reproductive costs associated with oviparity or viviparity will also influence patterns of telomere loss.

In addition to affecting maternal telomere length, differences between oviparous and viviparous life-history strategies could also affect the telomere length of the offspring that arise from these reproductive modes. A number of studies have reported a heritable component to telomere length (Bouwhuis, Verhulst, Bauch, & Vedder, 2018; Dugdale & Richardson, 2018). Moreover, offspring telomere length could also be subjected to maternal effects, for example via differences in embryonic provisioning and/or embryonic environment (McLennan et al., 2018a; Noguera, Metcalfe, Reichert, & Monaghan, 2016), as well as potential maternal effects on oocyte telomere length (Keefe, Kumar, & Kalmbach, 2015). Further, since oviparous offspring interact with the environment at an earlier life stage, it is possible that the telomeres of these offspring could then be differentially affected by environmental factors, for example via temperature and growth effects on early life development (McLennan et al., 2018b; Monaghan & Ozanne, 2018; Vedder, Verhulst, Zuidersma, & Bouwhuis, 2018).

The common lizard (*Zootoca vivipara*) is one of only a few extant vertebrate species in which both viviparity and oviparity occur (Surget-Groba et al., 2006). In this study, we examine maternal and offspring telomere dynamics in a wild common lizard population

in which different evolutionary lineages with either an oviparous or viviparous reproductive mode coexist within the same habitat. Reproductive mode in this species is known to be genetically determined and fixed between lineages (Arrayago, Bea, & Heulin, 1996; Recknagel & Elmer, 2019; Recknagel, Kamenos, & Elmer, 2018). The oviparous mothers lay calcified eggs ~33 days after copulation (i.e. oviposition) with a mean thickness of 40 μm , which are then incubated by the mother for ~28 days (Arrayago et al., 1996; Heulin, 1990; Lindtke, Mayer, & Böhme, 2010). Offspring from viviparous mothers are retained in utero for the duration of embryonic development (~57 days between copulation and birth; Arrayago et al., 1996) and are fully developed at birth (i.e. parturition). Each viviparous offspring is born individually surrounded by a thin membrane, from which they then 'hatch out' of immediately or within a few days after parturition (Recknagel & Elmer, 2019). Because this population occurs in a unique contact zone between the two reproductive modes, admixture is also possible (Lindtke et al., 2010; Recknagel, 2018). Admixed offspring are created when interbreeding occurs between the two reproductive modes. These offspring may be first-generation hybrids or result from the backcrossing of hybrids with oviparous or viviparous individuals. Offspring from admixed individuals are laid in thinner and less calcified eggs, but at a later developmental stage, compared with oviparous offspring. However, embryo mortality is also much higher in the admixed offspring, with estimates of around 40% (Lindtke et al., 2010).

Here, we have measured telomere length in viviparous, oviparous and admixed common lizards, of both mothers and their offspring. We predicted that viviparous mothers (with the presumption that they incur a higher reproductive burden) would have relatively shorter telomeres than oviparous mothers, while the telomere lengths of admixed mothers would be intermediate between their oviparous and viviparous conspecifics. Lastly, since there is known to be a heritable component to telomere length, we predicted that the patterns of telomere length in the offspring would follow those of the mothers.

2 | MATERIALS AND METHODS

2.1 | Field study

The field aspect of this study was conducted in the Gailtal valley of the Carinthian Alps, Austria (N 46.60°, E 13.14°), and under permit number HE3-NS-959/2013 (012/2016). This is currently the only known location where both viviparous and oviparous common lizards co-occur in high densities and occasionally interbreed (Cornetti et al., 2015; Lindtke et al., 2010; Recknagel, 2018). The study site covered an area of approximately 0.3 km² and an altitudinal range of 1,380–1,580 m. Wild female lizards were caught between May and July 2016; females can be distinguished from males by the absence of a hemipenial bulge at the base of the tail. Whether or not a female had recently mated was identified by the presence of a biting mark on the female's belly or flank (Lindtke et al., 2010). Immediately after capture, all lizards were weighed (to the nearest 0.001 g) and

snout–vent length (SVL) and tail length (TL) were measured (to the nearest 0.01 mm). Tail autotomy (self-amputation of the tail as a defence from predators and conspecifics) is a common occurrence in this species. Tail autotomy has previously been linked to telomere dynamics (Olsson, Pauliny, Wapstra, & Blomqvist, 2010); therefore, for this study, we decided to include only females that had a non-autotomized tail.

Females that had recently mated (and were therefore likely to be pregnant) were moved to nearby holding facilities so that their reproductive mode could be assessed. All females were individually housed in plastic terraria (56 × 39 × 28 cm) that were covered by netting on the top and on one side, to allow sufficient airflow. The terraria were housed within tents, so that each lizard was exposed to natural temperature variation. Each terrarium contained sand substrate, suitable shelter (e.g. pieces of dried wood), moisturized moss and a bowl of water. Lizards were fed ad libitum with mealworms (*Tenebrio molitor*) and crickets (*Gryllus assimilis*).

Each female was checked daily for the presence of offspring. On the same day that a female had undergone oviposition (to shelled eggs) or parturition (to live young), the female was weighed (to the nearest 0.001 g) and the number of offspring within a clutch (hereafter clutch size, CZ), relative clutch mass (RCM: clutch mass, including eggshell, amniotic fluids and yolk, divided by female mass after oviposition/parturition) and relative offspring mass (ROM: summed mass of the offspring after hatching, excluding eggshell, amniotic fluids and yolk, divided by female mass after oviposition/parturition) was measured. After oviposition/parturition, a tail clip was taken from each female for subsequent sequencing and telomere analysis. While absolute telomere lengths may differ among tissues, studies have found strong correlations in telomere length between different tissues in birds (Reichert, Criscuolo, Verinaud, Zahn, & Massemin, 2013) and lizards (Rollings et al., 2019). Moreover, a study on brown trout *Salmo trutta* by Debes et al. (2016) found trends in telomere dynamics to be similar among tissues, including highly regenerative tissues such as fin. Therefore, we were confident in using the tail clips for a non-invasive measurement of telomere length. Shortly after sampling, the females were returned and released back to the same location at which they had been caught. Unhatched offspring were incubated at 24°C in an Exo Terra Incubator. On the day of birth or hatching, each individual was weighed (to the nearest 0.001 g), SVL was measured (to the nearest 0.01 mm), and a tail clip was taken for subsequent telomere analysis. The offspring were then returned and released back to the same location at which the mothers had been caught. In total, we had complete tail samples for 69 mothers (23 oviparous, 33 viviparous and 13 admixed individuals). Offspring that did not hatch were excluded. In total, we had offspring tail clip samples from 57 of the mothers (20 oviparous, 29 viviparous and 8 admixed individuals).

2.2 | DNA extraction

The tail clip samples were transferred to the University of Glasgow, UK. DNA was extracted from all samples using the Macherey-Nagel

DNA NucleoSpin Tissue Extraction Kit, following the manufacturer's protocol. For each of the mothers, a cross-sectional sample of the tail (approximately 3 mm wide) was incubated in 180 μ l lysis buffer + 20 μ l of proteinase K solution (20 mg/ml) at 56°C overnight. Samples were then centrifuged to separate bone fragments from the tissue lysate, with the lysate being used in the subsequent DNA extraction. For the offspring, all siblings within a clutch were pooled (min 1 offspring, max 10 offspring) so that there was one averaged offspring DNA sample per mother. Again, a cross-sectional sample of tail (approximately 1 mm wide) was taken from each of the siblings and the pooled tissue was then processed in the same way as for the mothers. DNA concentration and purity was measured spectrophotometrically using a NanoDrop 8000, which confirmed that all samples met the recommended A260/280 ratio and had a DNA concentration >20 ng/ μ l.

2.3 | Identifying maternal reproductive mode by ddRADSeq

Oviparous and viviparous females are easily distinguishable by phenotype: oviparous offspring are laid in calcified shells and then require ~28 days of incubation prior to hatching, while viviparous offspring are born fully developed, but in a thin membrane that they then 'hatch out' of within a few hours to days after birth (Arrayago et al., 1996; Lindtke et al., 2010). However, because our study population occurs in a unique contact zone, there were also females that laid admixed offspring that exhibited intermediate phenotypes, such as partially calcified shells and fewer days of incubation compared to oviparous clutches. Therefore, we used a double-digest restriction site-associated sequencing (ddRADSeq) approach to genetically distinguish between the reproductive modes and admixed individuals. To do so, we followed the ddRADSeq library preparation protocol of Recknagel et al. (2018); see Appendix S1 for details.

Females were assigned a membership value (Q) to establish from which lineage of which reproductive mode they derived (Recknagel et al., 2018). Females that had a strong signature of oviparous genomic background (Q value ≤ 0.01) were assigned as oviparous, while females with a strong viviparous genomic background (Q value ≥ 0.99) were assigned as viviparous. Females that had a genomic background of admixture (Q value >0.01 and <0.99) were assigned as admixed individuals. We assigned a reproductive mode to 67 out of 69 of the females based on their genotypes. Two of the lizards were not included in the ddRADSeq library; however, their number of incubation days was clearly in the viviparous and oviparous ranges (2 and 35, respectively); therefore, we were confident in assigning their reproductive mode based on phenotype alone.

2.4 | Telomere analysis

Common lizard telomere length has been previously measured using the TeloTAGGG Telomere Length Assay (Dupoué et al., 2017), confirming that common lizard telomeres are also made up of the repetitive sequence TTAGGG. For this study, telomere length was

measured in all samples using the quantitative PCR method described by Cawthon (2002). The universal Tel1b and Tel2b primers designed by Cawthon (2002) and modified by Epel et al. (2004) were used for amplification of the telomere repeats. The recombination activating gene 1 (RAG-1) was chosen as the single-copy gene, and the *Z. vivipara* RAG-1 sequence (GenBank accession number: KY762205.1) was used to design primers. The following forward and reverse RAG-1 primers successfully amplified a single amplicon, as determined by melt curve analysis, and were subsequently used in the analysis:

LizRAG1-F 5'-GCC AAC TGC AAC AAG ATA CAC-3' and LizRAG1-R 5'-GAT ATG CTC ACA GAC CTG ACA A-3'.

A full outline of the qPCR protocol is provided in the Appendix S1. The samples (69 mothers and 57 offspring) were randomly distributed across six sets of PCR plates. qPCR data were analysed using the qBASE software for Windows (Hellemans, Mortier, Paepe, Speleman, & Vandesompele, 2007), as described in McLennan et al. (2016). For each sample, the qBASE software produced a calibrated normalized relative quantity (CNRQ). This is similar to the T/S ratio described by Cawthon (2002) but with greater control of the qPCR efficiency and inter-plate variation (see Appendix S1 for further details). Three points from the standard curve (5, 10 and 20 ng/well) were used as inter-run calibrators during qBASE analysis, to help correct for inter-run variation. The remaining three points of the standard curve (40, 2.5 and 1.25 ng/well) were used to calculate an inter-assay coefficient of variability of the CNRQs (which was 8.22). The average intra-plate variation of the Ct values was 0.94 for the telomere assay and 0.33 for the RAG-1 assay, respectively. The average inter-plate variation of the Ct values was 1.44 for the telomere assay and 0.34 for the RAG-1 assay, respectively. The efficiencies of the telomere and RAG-1 assays ranged from 95.6% to 108.5% and 91.7% to 97.9%, respectively, and were therefore within the acceptable range (85%–115%). The average quantification cycle (Ct) for the telomere and RAG-1 assays was 12.04 and 22.87, respectively. We found no significant difference in the RAG-1 Ct values among the three reproductive modes (oviparous, viviparous and admixed) (GLM $F_{2,66} = 1.08$, $p = .35$), but a highly significant difference among the telomere Ct values (GLM $F_{2,66} = 19.08$, $p < .001$).

2.5 | Statistical analysis

We used a Pearson correlation coefficient matrix to assess potential collinearity between all covariates (with a cut-off coefficient of 0.8). Mass and length were highly collinear for both mothers (Pearson $r = .91$, $p < .001$) and offspring (Pearson $r = .91$, $p < .001$); therefore, only mass was used for subsequent analyses. All variables used in the analyses are shown in Table 1.

All statistical analyses were conducted using R version 3.5.0 software. In total, we ran eight statistical models. Firstly, we ran general linear models (GLMs) to assess whether maternal mass (model 1), offspring mass (model 2), clutch size (model 3) and relative clutch mass RCM (model 4) and relative offspring mass

TABLE 1 Summary of all the variables used in statistical analyses

Variable name	Variable description
Life stage	Life stage at which a sample was taken (i.e. whether from mother or offspring). Factor
Reproductive mode	Reproductive phenotype of each mother, based on the ddRADSeq analysis. Offspring were assigned the same mode as their mother. Factor
Maternal RTL	Relative telomere length of each mother, measured at the individual level. Covariate
Mean offspring RTL	Relative offspring telomere length, measured as an average for each mother across all offspring. Covariate
Maternal mass	Somatic mass of each mother after oviposition or parturition (to the nearest 0.001 g). Covariate
Offspring mass	Somatic mass of offspring at the time of hatching, measured as an average for each mother (to the nearest 0.001 g). Covariate
RCM	Relative clutch mass: clutch mass divided by female mass after oviposition or parturition. Covariate
ROM	Relative offspring mass: sum of the offspring mass after hatching divided by female mass after oviposition/parturition. Covariate
Clutch size	Number of offspring born within a given clutch. Covariate

Note: See Section 22 for an outline of each model.

ROM (model 5) varied between the female reproductive mode. Secondly, we conducted a linear mixed model (LME) using the `lme4` and `lmerTest` functions (Bates, Maechler, Bolker, & Walker, 2015; Kuznetsova, Brockhoff, & Christensen, 2014) to assess whether RTL (relative telomere length) differed between life stage and reproductive mode, which included family ID as a random factor to control for non-independence between mother and offspring (model 6). Estimates of marginal (fixed effects) and conditional (fixed effects + random effects) R^2 values of the mixed model were calculated using the `MuMin` package. Finally, we looked at factors affecting variation in RTL at each life stage by conducting two separate GLMs. Firstly, maternal RTL was assessed in relation to reproductive mode, maternal mass, clutch size and RCM (model 7). This model was then simplified by backwards elimination, starting with the most insignificant term and continuing with insignificant main effects until the model contained only significant terms; providing that this resulted in a reduction of the AIC score. Secondly, mean offspring RTL was assessed in relation to maternal RTL, reproductive mode, offspring mass, clutch size, RCM and the interaction maternal RTL with reproductive mode to assess whether the relationship between maternal RTL and mean offspring RTL differed between the reproductive modes (model 8). As before, model 8 was then simplified by backwards elimination, starting with insignificant interactions and continuing with insignificant main effects until the model contained only significant terms. Variables were only removed from the model if this resulted in a reduction of the Akaike information criterion (AIC) score.

3 | RESULTS

3.1 | Life-history variation

We did not find a significant difference in somatic mass among the reproductive modes at the maternal stage, measured after oviposition/parturition (Table 2.1 and Figure 1a); however, there was a significant difference at the offspring stage, measured on the day of birth or

hatching, with oviparous offspring being the heaviest, viviparous offspring being the lightest and admixed offspring being the intermediate (GLM $F_{2,54} = 30.85$, $p < .001$; Table 2.2 and Figure 1b). The number of offspring within a clutch also differed significantly among the reproductive modes, with viviparous mothers producing the smallest clutches and oviparous mothers producing the largest clutches (GLM $F_{2,66} = 3.99$, $p = .023$; Table 2.3 and Figure 2c). RCM (relative clutch mass, including eggshell, amniotic fluids, etc.) did not significantly differ among the modes, suggesting that there was similar resource investment among the mothers (Table 2.4). However, we did find a significant difference in ROM (relative offspring mass, excluding eggshell, amniotic fluids, etc.) among the reproductive modes, with ROM being highest in the oviparous families and lowest in the viviparous families (GLM $F_{2,54} = 33.15$, $p < .001$; Table 2.5 and Figure 2b). This suggests that while there was a similar investment of maternal resources among the reproductive modes (in terms of embryo, eggshell, yolk and amniotic fluid production), the oviparous females had a higher net gain per clutch, supported by the larger clutch size and offspring mass.

3.2 | Telomere length variation

There was a significant difference in RTL between mothers and offspring. Mothers had relatively longer telomeres than their offspring within each of the three reproductive modes (LMM $F_{1,55.03} = 32.14$, $p < .001$; Table 3 and Figure 3). Telomere length also differed significantly between the reproductive modes. The viviparous mode had the longest telomeres, the oviparous mode had the shortest telomeres, and the admixed individuals had relatively intermediate telomeres (LMM $F_{2,59.13} = 72.55$, $p < .001$; Table 3 and Figure 3). This was true for both mothers (GLM $F_{2,66} = 37.98$, $p < .001$; Table 2.7 and Figure 3) and offspring (GLM $F_{2,54} = 63.12$, $p < .001$; Table 2.8 and Figure 3). When focusing only on the mothers, maternal mass, clutch size and RCM were not significantly associated with maternal RTL. For the offspring, maternal RTL, offspring mass, clutch size, RCM and the interaction maternal RTL X reproductive mode were not significantly associated with the relative mean telomere length.

TABLE 2 Summary of the final GLMs corresponding to models 1, 2, 3, 4, 5, 7 and 8

#		Explanatory variable	Estimate	SE	t	p
1	Maternal mass	–	–	–	–	–
2	Offspring mass	Intercept	0.204	0.010	19.67	<.001
		Repro. mode—oviparous	0.044	0.012	3.57	<.001
		Repro. mode—viviparous	–0.023	0.012	–1.97	.054
3	Clutch size	Intercept	7.154	0.553	12.94	<.001
		Repro. mode—oviparous	0.411	0.691	0.60	.554
		Repro. mode—viviparous	–1.063	0.653	–1.63	.108
4	RCM	–	–	–	–	
5	ROM	Intercept	0.414	0.034	12.11	<.001
		Repro. mode—oviparous	0.086	0.040	2.13	.037
		Repro. mode—viviparous	–0.140	0.039	–3.62	<.001
7	Maternal RTL	Intercept	0.043	0.033	1.28	.204
		Repro. mode—oviparous	–0.166	0.041	–3.98	<.001
		Repro. mode—viviparous	0.118	0.039	3.01	<.001
8	Mean offspring RTL	Intercept	–0.077	0.030	–2.52	.014
		Repro. mode—oviparous	–0.135	0.036	–3.73	<.001
		Repro. mode—viviparous	0.146	0.034	4.24	<.001

Note: See Section 22 for full definitions of the main effects and interactions initially included in each model. See Section 33 for analysis of variance test statistics.




FIGURE 1 The relationship between maternal reproductive mode and the somatic mass of (a) mothers and (b) offspring. Data plotted as individuals + mean. Somatic mass did not differ significantly between reproductive modes at the maternal stage; however, there was a significant difference at the offspring stage, measured on the day of birth or hatching ($p < .05$; see Table 2)

4 | DISCUSSION

This study has shown that different reproductive modes of the common lizard significantly differ in telomere length. The viviparous mode had the longest telomeres (for both mothers and offspring), the oviparous mode had the shortest telomeres (for both mothers and offspring), while the admixed individuals had relatively intermediate telomeres (again, for both mothers and offspring). Since the significant difference in telomere length was already evident at the offspring stage, it suggests that the reproductive modes have evolved differing telomere dynamics over time. However, it is currently unclear whether and/or how these differences in telomere length have co-evolved with other life-history traits. We also found that mothers had relatively longer

telomeres than their offspring among the reproductive ecotypes. However, we currently do not know whether this is due to physiological processes (e.g. telomere elongation mechanisms during development; Gomes et al., 2010) or due to the selective disappearance of individuals born with relatively smaller telomeres (e.g. Salmón, Nilsson, Watson, Bensch, & Isaksson, 2017).

We initially hypothesized that the viviparous mothers would have a higher reproductive investment (Lindtke et al., 2010) because of the prolonged pregnancy and presumed larger clutch mass in viviparous species (Horváthová et al., 2013; Qualls & Shine, 1998; Roitberg et al., 2013). However, the relative clutch mass (as a proxy for reproductive investment) was similar between the reproductive modes. Another potential cost to viviparity is that viviparous females may be less able to hunt and acquire resources during pregnancy, which could result in a net loss of energy. However, this was not possible




FIGURE 2 The relationship between maternal reproductive mode and (a) relative clutch mass (clutch mass, including eggshell, amniotic fluids and yolk, divided by female mass after oviposition/parturition), (b) relative offspring mass (summed mass of the offspring after hatching, excluding eggshell, amniotic fluids and yolk, divided by female mass after oviposition/parturition) and (c) the number of offspring within a clutch. Data plotted as individuals + mean. RCM did not differ between the reproductive modes; however, there was a significant reproductive mode effect on ROM and clutch size ($p < .05$; see Table 2)

TABLE 3 Summary of the linear mixed-effect model, corresponding to model 6

#	Explanatory variable	Estimate	SE	df	t	p
6	RTL Intercept	0.035	0.025	74.78	1.35	.18
	Repro. mode—oviparous	-0.155	0.031	62.04	-5.00	<.001
	Repro. mode—viviparous	0.127	0.029	62.80	4.32	<.001
	Life stage—offspring	-0.094	0.017	55.03	-4.67	<.001

Note: Family ID was included as a random factor to control for non-independence between mother and offspring. The family level variance was 0.0029 (SD 0.0542), and the residual variance was 0.0083 (SD 0.091). See Section 33 for analysis of variance test statistics. Estimates of marginal (fixed effects) and conditional (fixed effects + random effects) R^2 values were 0.62 and 0.72, respectively.

to quantify in this study, since the pregnant females were fed ad libitum. It has also been suggested that viviparous species have longer life spans, compared to oviparous species, and are therefore able to

produce more clutches over the years (Meiri, Brown, & Sibly, 2012; Tinkle, Wilbur, & Tilley, 1970). However, it is currently unknown whether the same also applies to oviparous and viviparous modes




FIGURE 3 The relationship between reproductive mode and relative telomere length (RTL). Circles correspond to mothers, while triangles correspond to the offspring. Data plotted as individuals + mean

co-occurring within the same species, and we were unable to quantify the age and past reproductive history of the mother lizards in this study.

Also in contrast with our hypothesis, the viviparous females had the longest telomeres, while the oviparous females had the shortest telomeres. In mammals, telomere length is generally phylogenetically conserved, although exceptions to this do exist (Gomes et al., 2011, 2010). It is unknown to what extent this applies to reptiles. One possibility is that differing telomere lengths in common lizards have evolved as an effect of the life-history divergence among the reproductive modes. It is also possible that we wrongly presumed oviparity to carry a lesser maternal burden. We did not find a significant difference in maternal mass among the reproductive modes, and a study by Demarco and Guillette (1992) found viviparity to carry a minimal metabolic cost. Moreover, within a reproductive season, oviparous females might invest more in their offspring compared to viviparous females (Recknagel & Elmer, 2019). We found that ROM was larger for oviparous females compared to viviparous females, indicating that oviparous females provide their offspring with more nutrients (i.e. yolk). While we could not assess this here, the shorter time of pregnancy might also allow oviparous females to produce more than one clutch per year (Heulin, Guillaume, Vogrin, Surget-Groba, & Tadic, 2000; Lindtke et al., 2010), resulting in an even larger reproductive effort per season for oviparous females.

It is worth noting that post-embryonic telomere repair mechanisms, such as telomerase expression, have been documented in several lizard species to date (Alibardi, 2015; Ujvari et al., 2017). In addition, several studies on fish have found higher telomerase expression in actively dividing cells (Peterson, Mok, & Au, 2015; Yap, Yeoh, Brenner, & Venkatesh, 2005). We found that the viviparous offspring were significantly smaller at hatching but were of equivalent size at the maternal life stage, perhaps due to an increased rate of growth. Therefore, this possible divergence in telomere length between the reproductive modes may have occurred, in part because of differences in growth rate and its possible association with telomerase expression. Longer telomeres in viviparous common lizards might therefore have co-evolved with (a) a smaller reproductive investment per season and (b) an increase in growth rate and potentially also longevity.

We do not know the age of the maternal lizards and can therefore not rule out possible age effects. It may be that viviparous common lizards differ in their longevity or their age at maturity, although there are currently no data to suggest this. Telomeres have been found to shorten in humans and other longer-lived mammals and birds (Hausmann et al., 2003), in part because many larger, longer-lived endotherm species appear to down-regulate telomere repair mechanisms in post-embryonic somatic tissues, perhaps as a tumour suppression mechanism (Gomes et al., 2010). However, the direction of the telomere–age relationship is less well established in ectothermic species, and studies have detected telomerase expression in a number of reptile, amphibian and fish species (Gomes et al., 2010; Simide, Angelier, Gaillard, & Stier, 2016; Ujvari et al., 2017). Therefore, we may have identified a difference in telomere length

between the reproductive modes, in part because age at reproduction may also differ among the modes. However, age and body size are thought to be correlated in the common lizard (Richard, Lecomte, Fraipont, & Clobert, 2005) and we included body mass (highly correlated with body size) as a covariate in analyses. In addition, the exact same pattern was found at the offspring stage, when age was similar among the reproductive modes (i.e. days since copulation).

As with the adult females, we also found that the viviparous offspring had the longest telomeres, the oviparous offspring had the shortest telomeres, and the admixed offspring had intermediate telomere lengths. Had this embryonic development occurred in the wild, we might have hypothesized that the viviparous offspring had longer telomeres because viviparity is thought to confer more stable embryonic conditions (Vedder et al., 2018). However, all of the embryos in this study developed in stable incubator conditions. While there may still be some degree of variation in embryonic conditions (e.g. higher level of respiratory exchange in the viviparous mode), it is unlikely that this caused the observed differences in telomere length. The complexity of telomere heritability is still not fully understood in lizards. It could be that offspring inherit their telomere length via the initial telomere length of the fertilized egg (Dugdale & Richardson, 2018), presuming that viviparous mothers also have a longer germline telomere length. Additionally, offspring may also be inheriting the genetic information that controls telomere elongation during embryogenesis (Kalmbach, Robinson, Wang, Liu, & Keefe, 2014; Liu et al., 2007; Schaetzlein et al., 2004). For example, it is thought that these embryonic telomere elongation programmes may restore telomeres to a set length (Schaetzlein et al., 2004); therefore, it is possible that this genetic information has also diverged between the reproductive modes.

We still know relatively little about the determinants of species-specific telomere length ranges, and how they may have evolved from species-specific trade-offs between long and short telomeres. It is often hard to disentangle such differences, in part because species differ significantly in the types of environment that they inhabit. However, by studying a unique common lizard population that exhibits both oviparous and viviparous reproduction in the same habitat, we have identified potential links between life-history divergence and telomere length, suggesting that populations such as these may prove fruitful in future studies. It would also now be interesting to examine whether the difference in telomere length between oviparous and viviparous individuals may affect long-term patterns of senescence and longevity.

ACKNOWLEDGEMENTS

We thank A. Adam, M. Capstick and W. Boner for advice and help with the laboratory analyses. We would also like to thank M. Layton, M. Sutherland, H. Leitao and Alpenferienpark Reisach for assistance in the field. All procedures were carried out under permit number HE3-NS-959/2013 (012/2016). The telomere analyses were funded by a Glasgow Natural History Society Blodwen Lloyd Binns small research grant, awarded to D.M. and H.R. During the period of data

collection, analysis and writing of this paper, D.M. was funded by ERC Advanced Grant 322784 and H.R. by a University of Glasgow Lord Kelvin-Adam Smith PhD Studentship and a NERC grant NE/N003942/1.

AUTHORS' CONTRIBUTIONS

D.M., H.R., K.R.E. and P.M. conceived of the idea for this study. H.R. led the field aspect of the study. D.M. and H.R. conducted laboratory analyses. D.M. and H.R. led data analysis and writing of the manuscript, with critical contribution from K.R.E. and P.M. All authors gave final approval for publication.

DATA AVAILABILITY STATEMENT

Data are available via the Dryad Digital Repository <https://doi.org/10.5061/dryad.712hc6m> (McLennan, Recknagel, Elmer, & Monaghan, 2019).

ORCID

Darryl McLennan  <https://orcid.org/0000-0002-8566-9437>

Hans Recknagel  <https://orcid.org/0000-0003-2728-0690>

Kathryn R. Elmer  <https://orcid.org/0000-0002-9219-7001>

REFERENCES

- Alibardi, L. (2015). Immunolocalization of the telomerase-1 component in cells of the regenerating tail, testis, and intestine of lizards. *Journal of Morphology*, 276, 748–758. <https://doi.org/10.1002/jmor.20375>
- Arrayago, M.-J., Bea, A., & Heulin, B. (1996). Hybridization experiment between oviparous and viviparous strains of *Lacerta vivipara*: A new insight into the evolution of viviparity in reptiles. *Herpetologica*, 52, 333–342.
- Bailey, N. W., Gray, B., & Zuk, M. (2010). Acoustic experience shapes alternative mating tactics and reproductive investment in male field crickets. *Current Biology*, 20, 845–849. <https://doi.org/10.1016/j.cub.2010.02.063>
- Barnes, R. P., Fouquerel, E., & Opresko, P. L. (2018). The impact of oxidative DNA damage and stress on telomere homeostasis. *Mechanisms of Ageing and Development*, 177, 37–45. <https://doi.org/10.1016/j.mad.2018.03.013>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48.
- Baum, D., Laughton, R., Armstrong, J. D., & Metcalfe, N. B. (2004). Altitudinal variation in the relationship between growth and maturation rate in salmon parr. *Journal of Animal Ecology*, 73, 253–260. <https://doi.org/10.1111/j.0021-8790.2004.00803.x>
- Blackburn, D. G. (1999). Are viviparity and egg-guarding evolutionarily labile in squamates? *Herpetologica*, 55, 556–573.
- Blackburn, E. H. (1991). Structure and function of telomeres. *Nature*, 350, 569–573. <https://doi.org/10.1038/350569a0>
- Boonekamp, J. J., Bauch, C., Mulder, E., & Verhulst, S. (2017). Does oxidative stress shorten telomeres? *Biology Letters*, 13, 20170164. <https://doi.org/10.1098/rsbl.2017.0164>
- Boonekamp, J. J., Mulder, G. A., Salomons, H. M., Dijkstra, C., & Verhulst, S. (2014). Nestling telomere shortening, but not telomere length, reflects developmental stress and predicts survival in wild birds. *Proceedings of the Royal Society B: Biological Sciences*, 281, 20133287. <https://doi.org/10.1098/rspb.2013.3287>
- Bouwhuis, S., Verhulst, S., Bauch, C., & Vedder, O. (2018). Reduced telomere length in offspring of old fathers in a long-lived seabird. *Biology Letters*, 14, 20180213. <https://doi.org/10.1098/rsbl.2018.0213>
- Bronikowski, A. M., & Arnold, S. J. (1999). The evolutionary ecology of life history variation in the garter snake *Thamnophis elegans*. *Ecology*, 80, 2314–2325.
- Cawthon, R. M. (2002). Telomere measurement by quantitative PCR. *Nucleic Acids Research*, 30(10), e47. <https://doi.org/10.1093/nar/30.10.e47>
- Cornetti, L., Belluardo, F., Ghielmi, S., Giovine, G., Ficetola, G. F., Bertorelle, G., ... Hauffe, H. C. (2015). Reproductive isolation between oviparous and viviparous lineages of the Eurasian common lizard *Zootoca vivipara* in a contact zone. *Biological Journal of the Linnean Society*, 114, 566–573.
- Cram, D. L., Monaghan, P., Gillespie, R., & Clutton-Brock, T. (2017). Effects of early-life competition and maternal nutrition on telomere lengths in wild meerkats. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20171383. <https://doi.org/10.1098/rspb.2017.1383>
- Debes, P. V., Visse, M., Panda, B., Ilmonen, P., & Vasemagi, A. (2016). Is telomere length a molecular marker of past thermal stress in wild fish? *Molecular Ecology*, 25, 5412–5424. <https://doi.org/10.1111/mec.13856>
- Demarco, V., & Guillelte, L. J. Jr (1992). Physiological cost of pregnancy in a viviparous lizard (*Sceloporus jarrovi*). *Journal of Experimental Zoology*, 262, 383–390. <https://doi.org/10.1002/jez.1402620404>
- Dugdale, H. L., & Richardson, D. S. (2018). Heritability of telomere variation: It is all about the environment!. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20160450. <https://doi.org/10.1098/rstb.2016.0450>
- Dupoué, A., Rutschmann, A., Le Galliard, J. F., Clobert, J., Angelier, F., Marciau, C., ... Meylan, S. (2017). Shorter telomeres precede population extinction in wild lizards. *Scientific Reports*, 7, 16976. <https://doi.org/10.1038/s41598-017-17323-z>
- Epel, E. S., Blackburn, E. H., Lin, J., Dhabhar, F. S., Adler, N. E., Morrow, J. D., & Cawthon, R. M. (2004). Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 17312–17315. <https://doi.org/10.1073/pnas.0407162101>
- Gomes, N. M., Ryder, O. A., Houck, M. L., Charter, S. J., Walker, W., Forsyth, N. R., ... Wright, W. E. (2011). Comparative biology of mammalian telomeres: Hypotheses on ancestral states and the roles of telomeres in longevity determination. *Aging Cell*, 10, 761–768. <https://doi.org/10.1111/j.1474-9726.2011.00718.x>
- Gomes, N. M. V., Shay, J. W., & Wright, W. E. (2010). Telomere biology in Metazoa. *FEBS Letters*, 584, 3741–3751. <https://doi.org/10.1016/j.febslet.2010.07.031>
- Hausmann, M. F., Winkler, D. W., O'Reilly, K. M., Huntington, C. E., Nisbet, I. C. T., & Vleck, C. M. (2003). Telomeres shorten more slowly in long-lived birds and mammals than in short-lived ones. *Proceedings of the Royal Society B: Biological Sciences*, 270, 1387–1392. <https://doi.org/10.1098/rspb.2003.2385>
- Hellemans, J., Mortier, G., De Paepe, A., Speleman, F., & Vandesompele, J. (2007). qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biology*, 8, R19.
- Heulin, B. (1990). Étude comparative de la membrane coquillière chez les souches ovipare et vivipare du lézard *Lacerta vivipara*. *Canadian Journal of Zoology*, 68, 1015–1019.
- Heulin, B., Guillaume, C. P., Vogrin, N., Surget-Groba, Y., & Tadic, Z. (2000). Further evidence of the existence of oviparous populations of *Lacerta (Zootoca) vivipara* in the NW of the Balkan Peninsula. *Comptes Rendus de l'Académie des Sciences - Series III - Sciences de la Vie*, 323, 461–468. [https://doi.org/10.1016/S0764-4469\(00\)00155-4](https://doi.org/10.1016/S0764-4469(00)00155-4)

- Horváthová, T., Cooney, C. R., Fitz, P. S., Oksanen, T. A., Jelić, D., Ghira, I., ... Jandzik, D. (2013). Length of activity season drives geographic variation in body size of a widely distributed lizard. *Ecology and Evolution*, 3, 2424–2442. <https://doi.org/10.1002/ece3.613>
- Kalmbach, K., Robinson, L. G., Wang, F., Liu, L., & Keefe, D. (2014). Telomere length reprogramming in embryos and stem cells. *BioMed Research International*, 2014, 925121. <https://doi.org/10.1155/2014/925121>
- Keefe, D., Kumar, M., & Kalmbach, K. (2015). Oocyte competency is the key to embryo potential. *Fertility and Sterility*, 103, 317–322. <https://doi.org/10.1016/j.fertnstert.2014.12.115>
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2014). lmerTest: Tests for random and fixed effects for linear mixed effect models (lmer objects of lme4 package). R package version 2.0-11.
- Levy, M. Z., Allsopp, R. C., Futcher, A. B., Greider, C. W., & Harley, C. B. (1992). Telomere end-replication problem and cell aging. *Journal of Molecular Biology*, 225, 951–960. [https://doi.org/10.1016/0022-2836\(92\)90096-3](https://doi.org/10.1016/0022-2836(92)90096-3)
- Lindtke, D., Mayer, W., & Böhme, W. (2010). Identification of a contact zone between oviparous and viviparous common lizards (*Zootoca vivipara*) in central Europe: Reproductive strategies and natural hybridization. *Salamandra*, 46, 73–82.
- Liu, L., Bailey, S. M., Okuka, M., Munoz, P., Li, C., Zhou, L. J., ... Keefe, D. L. (2007). Telomere lengthening early in development. *Nature Cell Biology*, 9, 1436–1441. <https://doi.org/10.1038/ncb1664>
- McLennan, D., Armstrong, J. D., Stewart, D. C., McKelvey, S., Boner, W., Monaghan, P., & Metcalfe, N. B. (2016). Interactions between parental traits, environmental harshness and growth rate in determining telomere length in wild juvenile salmon. *Molecular Ecology*, 25, 5425–5438. <https://doi.org/10.1111/mec.13857>
- McLennan, D., Armstrong, J. D., Stewart, D. C., McKelvey, S., Boner, W., Monaghan, P., & Metcalfe, N. B. (2018a). Links between parental life histories of wild salmon and the telomere lengths of their offspring. *Molecular Ecology*, 27, 804–814. <https://doi.org/10.1111/mec.14467>
- McLennan, D., Armstrong, J. D., Stewart, D. C., McKelvey, S., Boner, W., Monaghan, P., & Metcalfe, N. B. (2018b). Telomere elongation during early development is independent of environmental temperatures in Atlantic salmon. *The Journal of Experimental Biology*, 221, jeb178616. <https://doi.org/10.1242/jeb.178616>
- McLennan, D., Recknagel, H., Elmer, K. R., & Monaghan, P. (2019). Data from: Distinct telomere differences within a reproductively bimodal common lizard population. Dryad Digital Repository, <https://doi.org/10.5061/dryad.712hc6m>
- Meiri, S., Brown, J. H., & Sibly, R. M. (2012). The ecology of lizard reproductive output. *Global Ecology and Biogeography*, 21, 592–602. <https://doi.org/10.1111/j.1466-8238.2011.00700.x>
- Metcalfe, N. B., & Monaghan, P. (2003). Growth versus lifespan: Perspectives from evolutionary ecology. *Experimental Gerontology*, 38, 935–940. [https://doi.org/10.1016/S0531-5565\(03\)00159-1](https://doi.org/10.1016/S0531-5565(03)00159-1)
- Meunier, J., Wong, J. W. Y., Gómez, Y., Kuttler, S., Röllin, L., Stucki, D., & Kölliker, M. (2012). One clutch or two clutches? Fitness correlates of coexisting alternative female life-histories in the European earwig. *Evolutionary Ecology*, 26, 669–682. <https://doi.org/10.1007/s10682-011-9510-x>
- Monaghan, P. (2010). Telomeres and life histories: The long and the short of it. *Annals of the New York Academy of Sciences*, 1206, 130–142. <https://doi.org/10.1111/j.1749-6632.2010.05705.x>
- Monaghan, P. (2014). Organismal stress, telomeres and life histories. *Journal of Experimental Biology*, 217, 57–66. <https://doi.org/10.1242/jeb.090043>
- Monaghan, P., & Ozanne, S. E. (2018). Somatic growth and telomere dynamics in vertebrates: Relationships, mechanisms and consequences. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20160446.
- Murphy, B. F., & Thompson, M. B. (2011). A review of the evolution of viviparity in squamate reptiles: The past, present and future role of molecular biology and genomics. *Journal of Comparative Physiology B*, 181, 575–594. <https://doi.org/10.1007/s00360-011-0584-0>
- Noguera, J. C., Metcalfe, N. B., Reichert, S., & Monaghan, P. (2016). Embryonic and postnatal telomere length decrease with ovulation order within clutches. *Scientific Reports*, 6, 25915. <https://doi.org/10.1038/srep25915>
- Olsson, M., Friesen, C. R., Rollings, N., Sudyka, J., Lindsay, W., Whittington, C. M., & Wilson, M. (2018). Long-term effects of superoxide and DNA repair on lizard telomeres. *Molecular Ecology*, 27, 5154–5164. <https://doi.org/10.1111/mec.14913>
- Olsson, M., Pauliny, A., Wapstra, E., & Blomqvist, D. (2010). Proximate determinants of telomere length in sand lizards (*Lacerta agilis*). *Biology Letters*, 6, 651–653.
- Partridge, C. G., MacManes, M. D., Knapp, R., & Neff, B. D. (2016). Brain transcriptional profiles of male alternative reproductive tactics and females in bluegill sunfish. *PLoS ONE*, 11, e0167509. <https://doi.org/10.1371/journal.pone.0167509>
- Peterson, D. R., Mok, H. O. L., & Au, D. W. T. (2015). Modulation of telomerase activity in fish muscle by biological and environmental factors. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 178, 51–59. <https://doi.org/10.1016/j.cbpc.2015.09.004>
- Qualls, C. P., & Shine, R. (1995). Maternal body-volume as a constraint on reproductive output in lizards: Evidence from the evolution of viviparity. *Oecologia*, 103, 73–78. <https://doi.org/10.1007/BF00328427>
- Qualls, C. P., & Shine, R. (1998). Costs of reproduction in conspecific oviparous and viviparous lizards, *Lerista bougainvillii*. *Oikos*, 82, 539–551. <https://doi.org/10.2307/3546374>
- Recknagel, H. (2018). *Environmental constraints and genetic basis for the evolution of viviparity*. PhD Thesis, Doctoral, University of Glasgow.
- Recknagel, H., & Elmer, K. R. (2019). Differential reproductive investment in co-occurring oviparous and viviparous common lizards (*Zootoca vivipara*) and implications for life history trade-offs with viviparity. *Oecologia*, 190, 85–98. <https://doi.org/10.1007/s00442-019-04398-w>
- Recknagel, H., Kamenos, N. A., & Elmer, K. R. (2018). Common lizards break Dollo's law of irreversibility: Genome-wide phylogenomics support a single origin of viviparity and re-evolution of oviparity. *Molecular Phylogenetics and Evolution*, 127, 579–588. <https://doi.org/10.1016/j.ympev.2018.05.029>
- Reichert, S., Criscuolo, F., Verinaud, E., Zahn, S., & Massemin, S. (2013). Telomere length correlations among somatic tissues in adult zebra finches. *PLoS ONE*, 8, e81496. <https://doi.org/10.1371/journal.pone.0081496>
- Reichert, S., & Stier, A. (2017). Does oxidative stress shorten telomeres *in vivo*? A review. *Biology Letters*, 13, 20170463.
- Richard, M., Lecomte, J., De Fraipont, M., & Clobert, J. (2005). Age-specific mating strategies and reproductive senescence. *Molecular Ecology*, 14, 3147–3155. <https://doi.org/10.1111/j.1365-294X.2005.02662.x>
- Roitberg, E. S., Kuranova, V. N., Bulakhova, N. A., Orlova, V. F., Eplanova, G. V., Zinenko, O. I., ... Yakovlev, V. A. (2013). Variation of reproductive traits and female body size in the most widely-ranging terrestrial reptile: Testing the effects of reproductive mode, lineage, and climate. *Evolutionary Biology*, 40, 420–438. <https://doi.org/10.1007/s11692-013-9247-2>
- Rollings, N., Friesen, C. R., Sudyka, J., Whittington, C., Giraudeau, M., Wilson, M., & Olsson, M. (2017). Telomere dynamics in a lizard with morph-specific reproductive investment and self-maintenance. *Ecology and Evolution*, 7, 5163–5169. <https://doi.org/10.1002/ece3.2712>
- Rollings, N., Friesen, C. R., Whittington, C. M., Johansson, R., Shine, R., & Olsson, M. (2019). Sex- And tissue-specific differences in telomere length in a reptile. *Ecology and Evolution*, 9(11), 6211–6219. <https://doi.org/10.1002/ece3.5164>
- Salmón, P., Nilsson, J. F., Nord, A., Bensch, S., & Isaksson, C. (2016). Urban environment shortens telomere length in nestling great tits, *Parus major*. *Biology Letters*, 12, 20160155.

- Salmón, P., Nilsson, J. F., Watson, H., Bensch, S., & Isaksson, C. (2017). Selective disappearance of great tits with short telomeres in urban areas. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20171349. <https://doi.org/10.1098/rspb.2017.1349>
- Schaetzlein, S., Lucas-Hahn, A., Lemme, E., Kues, W. A., Dorsch, M., Manns, M. P., ... Rudolph, K. L. (2004). Telomere length is reset during early mammalian embryogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 8034–8038. <https://doi.org/10.1073/pnas.0402400101>
- Shine, R. (1995). A new hypothesis for the evolution of viviparity in reptiles. *The American Naturalist*, 145, 809–823. <https://doi.org/10.1086/285769>
- Simide, R., Angelier, F., Gaillard, S., & Stier, A. (2016). Age and heat stress as determinants of telomere length in a long-lived fish, the Siberian sturgeon. *Physiological and Biochemical Zoology*, 89, 441–447. <https://doi.org/10.1086/687378>
- Sites, J. W., Reeder, T. W., & Wiens, J. J. (2011). Phylogenetic insights on evolutionary novelties in lizards and snakes: Sex, birth, bodies, niches, and venom. *Annual Review of Ecology, Evolution, and Systematics*, 42, 227–244. <https://doi.org/10.1146/annurev-ecolsys-102710-145051>
- Surget-Groba, Y., Heulin, B., Guillaume, C.-P., Puky, M., Semenov, D., Orlova, V., ... Smajda, B. (2006). Multiple origins of viviparity, or reversal from viviparity to oviparity? The European common lizard (*Zootoca vivipara*, Lacertidae) and the evolution of parity. *Biological Journal of the Linnean Society*, 87, 1–11. <https://doi.org/10.1111/j.1095-8312.2006.00552.x>
- Taborsky, M., & Brockmann, H. J. (2010). Alternative reproductive tactics and life history phenotypes. In P. Kappeler (Ed.), *Animal behaviour: Evolution and mechanisms* (pp. 537–586). Berlin, Heidelberg: Springer.
- Tian, X., Doerig, K., Park, R., Can Ran Qin, A., Hwang, C., Neary, A., ... Gorbunova, V. (2018). Evolution of telomere maintenance and tumour suppressor mechanisms across mammals. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20160443. <https://doi.org/10.1098/rstb.2016.0443>
- Tinkle, D. W., Wilbur, H. M., & Tilley, S. G. (1970). Evolutionary strategies in lizard reproduction. *Evolution*, 24, 55–74. <https://doi.org/10.1111/j.1558-5646.1970.tb01740.x>
- Tsubaki, Y., Hooper, R. E., & Siva-Jothy, M. T. (1997). Differences in adult and reproductive lifespan in the two male forms of *Mnais pruinosa costalis selys* (Odonata: Calopterygidae). *Researches on Population Ecology*, 39, 149–155. <https://doi.org/10.1007/BF02765260>
- Ujvari, B., Biro, P. A., Charters, J. E., Brown, G., Heasman, K., Beckmann, C., & Madsen, T. (2017). Curvilinear telomere length dynamics in a squamate reptile. *Functional Ecology*, 31, 753–759. <https://doi.org/10.1111/1365-2435.12764>
- Vedder, O., Verhulst, S., Zuidersma, E., & Bouwhuis, S. (2018). Embryonic growth rate affects telomere attrition: An experiment in a wild bird. *The Journal of Experimental Biology*, 221, jeb181586. <https://doi.org/10.1242/jeb.181586>
- Willbourn, R. V., Moatt, J. P., Froy, H., Walling, C. A., Nussey, D. H., & Boonekamp, J. J. (2018). The relationship between telomere length and mortality risk in non-model vertebrate systems: A meta-analysis. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1741), 20160447–<https://doi.org/10.1098/rstb.2016.0447>
- Wourms, J. P., & Lombardi, J. (1992). Reflections on the evolution of piscine viviparity. *American Zoologist*, 32, 276–293. <https://doi.org/10.1093/icb/32.2.276>
- Yap, W. H., Yeoh, E., Brenner, S., & Venkatesh, B. (2005). Cloning and expression of the reverse transcriptase component of pufferfish (*Fugu rubripes*) telomerase. *Gene*, 353, 207–217. <https://doi.org/10.1016/j.gene.2005.04.038>
- Zamudio, K., & Chan, L. (2008). Alternative reproductive tactics in amphibians. In R. Oliveira, M. Taborsky, & H. Brockmann (Eds.), *Alternative reproductive tactics: An integrative approach* (pp. 300–331). Cambridge: Cambridge University Press.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: McLennan D, Recknagel H, Elmer KR, Monaghan P. Distinct telomere differences within a reproductively bimodal common lizard population. *Funct Ecol*. 2019;33:1917–1927. <https://doi.org/10.1111/1365-2435.13408>