

Ruminant Session II

Wednesday, September 18, 2013

Pre-Weaning Calf Nutrition and Subsequent Feedlot Performance

A.E. Radunz

University of Wisconsin-River Falls

Introduction

Beef cattle production in the United States is currently facing many challenges. The nation's cow herd has continued to decline since its peak in 1975 (NASS USDA, 2010). A major reason for this decline in cow numbers is increased input costs, of which feed represents the greatest portion. Therefore, investigations on how to more efficiently produce beef from conception and how environment may influence the final phenotype of the animal are critical. Phenotype of the animal is combination of genetics and environment. While genes the animal inherits from his/her parents cannot be changed, how the genes are expressed can be influenced by environmental factors, such as nutrition, from the time of conception. Most of the research in beef cattle feedlot nutrition has focused on post-weaning environmental influences on the final phenotype of the calf in regards to growth and carcass traits; however, recent research in the area of developmental programming has started to provide more insight on the impacts of early postnatal period and even gestational environment on lifetime productivity of humans and livestock species. Recent research has begun to demonstrate that maternal gestation nutrition may impact progeny postnatal growth, health, feed efficiency, and carcass composition as outlined here.

Developmental Programming

The fetal origins hypothesis was first proposed by Dr. Barker based on epidemiological studies investigating low nutrient intake by pregnant mothers during the 1944 Dutch Famine of World War II, and the resulting long-term health implication of their children. Dr. Barker reported low caloric intake during gestation induced drastic changes in developmental programming, thereby impacting future health of offspring observed by increased incidence of clinical conditions such as obesity, insulin resistance, and type 2 diabetes (Barker et al., 2002).

Similarly, in domestic animal studies, maternal nutrition during gestation has been reported to impact postnatal body composition, insulin sensitivity, and growth rate, which all have implications for production efficiency and meat quality (Ford et al., 2007; Radunz et al., 2010 and 2012). Factors such as maternal nutrition, environment, or stressors during gestation can change nutrient supply to the fetus, which can then affect growth and development of organs, skeletal muscle tissue, and adipose tissue. These changes in fetal development can alter postnatal skeletal muscle growth, fat deposition, insulin resistance, or hypertension of offspring, which can impact economically important traits such as growth rate, health, and carcass composition. To date most of the research using animals has been focused on human health implications, but recent studies have investigated beef cow gestation nutrition and management to provide evidence of its implications to offspring productivity.

One of the mechanisms by which developmental programming may be explained is epigenetics. Epigenetics encompasses changes to marks on the genome early in development that are

copied from one cell generation to the next, which may alter gene expression, but do not involve changes in primary DNA sequence. Epigenetic mechanisms such as DNA methylation and histone modifications (e.g., acetylation, methylation, and phosphorylation) can change genome activity under some environmental (nutrition or toxicants) conditions (Bollati and Baccarelli, 2010). A greater understanding of how maternal nutrition induces epigenetic modifications to adipose tissue would provide critical information in understanding pathways influencing postnatal adipose deposition.

Developmental windows of muscle and adipose tissues, economically important tissues in meat animal production, occur during gestation, which could influence production efficiency and carcass composition of the individual. Muscle hyperplasia (increasing cell number) starts in early gestation and terminates during mid-gestation. Any impact on muscle growth after this point time is achieved by hypertrophy (increasing cell size). Adipose tissue growth primarily begins during late gestation and adipose tissue hyperplasia can continue until maturity, however at a diminishing rate as the animal becomes older. Previous research in early weaning and high starch diets in beef cattle has already provided evidence of postnatal environmental influences on adiposity. In addition to these tissues, other developmental windows occur throughout gestation for the placenta, specific tissues, and organs. Therefore, not only does the type of environmental stress influence the final phenotype but also at what time in development the stress occurs. For example, nutrient restriction followed by adequate nutrition in early gestation results in larger birth weights in sheep compared to adequate nutrition, whereas nutrient restriction in late gestation results in lower birth weights (Munoz et al., 2008). Collectively, these studies and other research have provided evidence the developmental windows occur from periconception to early postnatal life of the animal (Fowden et al., 2006).

Post-weaning Growth Traits

Milk production postpartum can be influenced by prepartum nutrition of the dam, which could have development programming implications. Cows receiving an energy-deficient gestation diet had lower milk production (Corah et al., 1975). Additionally, cows allowed limited vs. ad libitum grazing access prepartum had a 9% decrease in early lactation milk production (Kearnan and Beal, 1992). Over- and under nutrition of ewes resulted in decreased IgG concentration, nutrient content, and volume of colostrum (Swanson et al., 2007). Studies have reported prepartum restriction of nutrients resulted in a decrease in colostrum IgG (Shell et al., 1995) and absorption of IgG by calves. Effective passive transfer of IgG in colostrum is vital to calf health and immunity (Perino et al., 1995). These studies indicate that late gestation nutrition impacts mammary gland development, which could impact postpartum milk production and passive immune transfer thus impacting postnatal growth and health.

Most studies investigating gestational nutrition on progeny performance have used gestating sheep as a biomedical model and have focused on the effects of global under-nutrition (McMillen et al., 2001; Ford et al., 2007) and global over-nutrition (Wallace et al., 2002; Long et al., 2010). These studies have reported an association with maternal nutrient intake and progeny's postnatal body composition and glucose metabolism as well as indicated that timing and duration of nutrient modification during gestation differentially impacts outcomes. In cattle, providing a high vs. low energy diet during late gestation in beef cattle was reported to increase calf birth weight and subsequent weaning weight (Corah et al., 1975). The impact of global nutrient restriction during early gestation (d 32 to 115) has varied. Long et al. (2008) observed that nutrient restriction of cows did not influence birth weights or postnatal growth calves. In contrast, Underwood et al. (2008) reported greater postnatal growth and feed efficiency in steers from cows' nutrient restricted during a similar period of gestation (d 31 to

120). Nutrient restriction (55% global restriction) during early to mid-gestation in sheep (d 28-78) resulted in male offspring having similar birth weights, but lighter weights at slaughter, greater amounts of internal fat, and less muscle mass (Ford et al., 2007). In these studies, nutrient restriction appeared to have provided adequate energy for fetal growth, possibly at the expense of the dam's tissue because nutrient partitioning during pregnancy favors the fetus at the expense of the dam and the placenta efficiency may be different (Barcroft, 1946).

While studies investigating maternal over- or under-nutrition are valuable, the investigation of specific diet components could provide greater insight into mechanisms of developmental programming. A few studies in beef cattle have investigated specific diet components during gestation on postnatal progeny production traits. A series of reports from the University of Nebraska demonstrated that cows supplemented protein on dormant winter range in late gestation had steer progeny with greater postnatal growth rate and intramuscular fat deposition than progeny from cows not supplemented protein (Stalker et al., 2006; Martin et al., 2007; Larson et al., 2009). Previous research focused on developmental programming affects elicited by comparing energy sources fed at isoenergetic which differ in energy substrate supply (corn, hay, and corn dried distillers grains = DDGS) during late gestation in sheep and cattle (Radunz et al., 2012). When corn (high starch) and DDGS (high fiber and fat and excess protein) energy sources versus hay (high fiber) were fed to dams during late gestation this resulted in progeny with greater birth weights. In addition research at Purdue University demonstrated that feeding diets containing excess protein during the third trimester also resulted in greater birth weights (Gunn et al., 2012). Collectively, these studies indicate maternal nutrition in late gestation influence fetal growth.

Carcass Traits

Comparable to human health research, the desired impact is to deposit less external and internal fat during growth, not only because of the consequences to efficiency of growth but also impacts on such economically important traits as carcass value and reproduction. One distinct difference in humans compared to beef cattle is concerning the intramuscular (*i.e.* marbling) adipose depot. In beef cattle, marbling has greater economic value than in other livestock species (*i.e.* pigs, sheep), because this fat depot is a major determinant in USDA Quality Grade and is used to determine carcass value. Approximately 80% of fetal adipose tissue is deposited in the final few weeks of gestation, but the development of these adipocytes starts earlier in gestation (Symonds et al., 2007). Adipose tissue growth occurs through two mechanisms: 1) preadipocyte proliferation; impacting capacity to form new mature adipocytes (hyperplasia) and 2) increased size and lipid storage capacity of mature adipocytes (hypertrophy). Adipocyte hyperplasia occurs primarily during late fetal development and early postnatal life in humans (Martin et al., 1998) and bovines (Zhu et al., 2008). This process is highly sensitive to nutritional environment and to the prevailing concentrations of insulin-like growth factors, glucose, insulin, and glucocorticoids (Martin et al., 1998). Although preadipocytes can proliferate and differentiate in adults, their capacity appears to be limited with most of the developmental work completed early in life (Martin et al., 1998). Therefore, evidence supports that the fetal and early postnatal periods are critical stages of adipose tissue programming which impacts later fat deposition.

Research from the University of Nebraska (Stalker et al., 2006; Stalker et al., 2007; and Larson et al., 2009) provided supporting evidence that late gestation is a critical period for marbling development (Table 1). Over a 3-year period, steer calves from cows grazing native range with or without protein supplementation were followed from birth to slaughter. Native range was determined not to meet the protein requirements of the cow during late gestation and protein

supplementation was provided to one group to meet those requirements. The calves born to cows supplemented with protein had carcasses with higher marbling scores, with a greater percentage of carcasses graded USDA Choice, and 60 lbs heavier hot carcass weights.

Additionally, Radunz et al. (2012) reported similar results indicating an important role of late gestation nutrition on marbling development and fetal growth. Calves born to cows fed corn had the least marbling and lower percentage of carcasses grading in the upper 2/3 of USDA Choice compared to calves born to cows fed DGGS or hay at the same fat endpoint. These results suggest that amount of marbling in the carcass may not only be determined by genetics, postnatal nutrition, and postnatal management, but also could be determined by what the cow is fed during gestation.

In a following research using the same model in sheep by Radunz, fetal adipose and *longissimus* muscle tissues collected at birth provide the first evidence maternal diet can influence expression of imprinted genes associated with adipose tissue development and growth, which could explain difference in postnatal fat deposition (Khatib et al., 2013).

Implications

One of the major challenges in gaining more knowledge in the area of developmental programming in beef cattle production is the time and resources needed to collect the data. Therefore at this time more questions may be raised than answered. The research presented here indicates maternal nutrition can impact postnatal growth and fat deposition in ruminants. More specifically, late gestation maternal nutrition may have a significant impact on intramuscular fat deposition in beef cattle. The question remains to be answered whether this is the result of changes in maternal body condition score, substrate supply to fetus, quantity and/or quality of protein supply or other dietary factors. In order to improve efficiency of beef cattle production, more research is warranted to investigate the impacts of environment, such as nutrition, during gestation and early postnatal life of cattle on lifetime productivity.

Table 1. Effects of cow protein supplementation during late-gestation on progeny feedlot performance and carcass traits

	Stalker et al. 2006		Stalker et al. 2007		Larson et al. 2009	
	NS ¹	S ¹	NS	S	NS	S
Weaning Wt., lbs	463 ^a	476 ^b	463 ^a	476 ^b	514 ^a	529 ^b
DMI, lb/d	18.7	18.8	24.6 ^a	26.6 ^b	19.8 ^x	20.3 ^y
F:G	5.41	5.46	6.97	7.19	5.37	5.38
HCW, lbs	800	814	765 ^a	805 ^b	805 ^a	822
USDA Choice, %	85	96	---	---	71 ^a	85 ^b
Marbling Score ²	467	479	449	461	445 ^a	492

^{ab}Means differ, $P \leq 0.05$

^{xy}Means differ, $P \leq 0.10$.

¹ NS = non supplemented and S = supplemented (contained = 62% dried distillers grain plus solubles, 11% wheat middlings, 9% cottonseed meal, 5% dry corn gluten feed, 5% molasses, 2% urea, 6% vitamin and trace mineral premix, and monensin)

² 400 - 499 = Low Choice; 500-599 = Average Choice

Table 2. Effects of cow late gestation dietary energy source on progeny feedlot performance and carcass traits

	Grass Hay ¹	Corn	DDGS	P-value
Birth weight, lbs	85.5 ^a	95.0 ^b	91.0 ^b	0.01
Weaning weight, lbs	548	569	564	0.09
ADG, lbs/d	3.37	3.46	3.41	0.48
DMI, lbs/d	19.3	19.6	19.6	0.78
G:F	0.174	0.178	0.176	0.43
Days on Feed, d	178	168	170	0.10
Hot carcass weight, lbs	688	688	675	0.59
12 th rib fat, in.	0.48	0.50	0.51	0.71
Ribeye area, in ²	12.0	12.0	11.9	0.79
Yield grade	2.82	2.82	2.85	0.93
Marbling score ²	549 ^a	506 ^b	536 ^{ab}	0.03

¹ Diets fed to cows during late gestation. HAY = ad libitum grass hay; CORN = limit-fed corn diet; and DDGS = limit-fed dried distillers grains

² Slight = 300 to 399, Small = 400 to 499, Modest = 500 to 599

References

- Barker, D.J.P., J.G. Eriksson, T. Forsen, and C. Osmond. 2002. Fetal origins of adult disease: Strength of effects and biological basis. *International Journal of Epidemiology* 31: 1235-1239.
- Barcroft, J. 1946. *Researches on prenatal life*. Oxford: Blackwell Scientific Publications.
- Corah, L.R., T.G., Dunn, and C.C. Kaltenbach. 1975. Influences of prepartum nutrition on the reproductive performance of beef females and the performance of their progeny. *J. Anim. Sci.* 41:819-824.
- Gunn, P.J., J.P. Schoonmaker, R.P. Lemenager, and G.A. Bridges. 2012. Meta-analysis on effects of supplementing distiller's grains to beef cows during early lactation on reproductive efficiency and pre-weaning growth. *J. Dairy Science*: 95 (Suppl. 2).
- Ford, S.P. et al. 2007. Maternal undernutrition during early to mid-gestation in the ewe results in altered growth, adiposity, and glucose tolerance in male offspring. *Journal of Animal Science* 85: 1285-1294.
- Khatib, H; X. Lan; E. Cretney; J. Kropp; K. Khateeb; M. A Berg; F. Peñagaricano; R. Magness, and A. E Radunz. 2013. Maternal diet during pregnancy induces transcriptomic and epigenomic changes in fetal tissues in sheep. *Epigenetics. Front Genet.* 2013; 4: 49.
- Kearnan, J.M., and W.E. Beal. 1992. Impact of pre-and postpartum diets on milk production, calf weaning weight, and rebreeding in beef cows. *J. Anim. Sci.* 70 (Supp. 1):277.
- Larson D.M., J.L. Martin, D.C. Adams, and R.N. Funston. 2009. Winter grazing system and supplementation during late gestation influence performance of beef cows and steer progeny. *J Anim Sci* 87: 1147-1155.
- Long, N.M., K.A. Vonnahme, B.W. Hess, P.W. Nathanielsz, and S.P. Ford. 2009. Effects of early gestational undernutrition on fetal growth, organ development, and placentomal composition in bovine. *J. Anim. Sci.* 87:1950-1959.
- Martin, J.L., K.A. Vonnahme, D.C. Adams, G.P. Lardy, and R.N. Funston. 2007. Effects of dam nutrition on growth and reproductive performance of heifer calves. *Journal of Animal Science* 85: 841-847.

- Martin, R.J., G.J. Hausman, and D.B. Hausman. 1998. Regulation of adipose cell development in utero. *Proceedings of the Society for Experimental Biology and Medicine* 219: 200-210.
- Perino, L.J., T.E. Wittum, and G.S. Ross. 1995. Effects of various risk factors on plasma protein and serum immunoglobulin concentrations of calves at postpartum hours 10 and 24. *Am. J. Vet. Res.* 56:1144-1148.
- Radunz, A, F. Fluharty, M. Day, H. Zerby, and S. Loerch. 2010. Prepartum dietary energy source fed to beef cows: I. Effects on pre- and postpartum cow performance. *J. Anim. Sci.* 88:2717-2728.
- Radunz, A, F. Fluharty, H. Zerby, and S. Loerch. 2011a. Winter-feeding systems for gestating sheep I. Effects on pre- and postpartum ewe performance and lamb progeny preweaning performance. *J. Anim. Sci.* 89:467-477.
- Radunz, A., F. Fluharty, I. Susin, T. Felix, H. Zerby, and S. Loerch. 2011b. Winter-feeding systems for gestating sheep II. Effects on feedlot performance, glucose tolerance, and carcass composition of lamb progeny. *J. Anim. Sci.* 89:478-488.
- Radunz, A.E, F.L. Fluharty, A.E. Relling, T.L. Felix, L.M. Shoup, H.N. Zerby, and S.C. Loerch. 2012. Prepartum dietary energy source fed to beef cows: II. Effects on progeny postnatal growth, glucose tolerance, and carcass composition. *J. Anim. Sci.* doi:10.2527/jas.2012-5098.
- Schell, T.M., R.J. Early, J.R. Carpenter, and B.A. Buckley. 1995. Prepartum nutrition and solar radiation in beef cattle: II. Residual effects on postpartum milk yield, immunoglobulin, and calf growth. *J. Anim. Sci.* 73: 1303-1309.
- Stalker, L.A., D.C. Adams, T.J. Klopfenstein, D.M. Feuz, and R.N. Funston. 2006. Effects of pre- and post-partum nutrition on reproduction in spring calving cows and calf feedlot performance. *J. Anim. Sci.* 84:2582-2589.
- Swanson, T.J., C.J. Hammer, J.B. Taylor, D.A. Redmer, K.A. Vonnahme, J.S. Luther, T.L. Neville, J.J. Reed, J.S. Caton, and L.P. Reynolds. 2007. Effects of plane of nutrition and selenium on colostrum quality and mammary development in ewes. *J. Anim. Sci.* 85 (Supp 1): (Abstr.).
- Symonds, M.E., T. Stephenson, D.S. Gardner, and H. Budge. 2007. Long-term effects of nutritional programming of the embryo and fetus: Mechanisms and critical windows. *Reproduction Fertility and Development* 19: 53-63.
- Underwood, K.R., J.F. Tong, J.M. Kinzey, P.L. Price, E.E. Grings, B.W. Hess, W.J. Means, and M. Du. 2008. Gestational nutrition affects growth and adipose tissue deposition in steers. *West. Sec. Am. Soc. Anim. Sci.* 59:29-32.
- Wallace, J.M., R.P. Aitken, J.S. Milne, and W.W. Hay. 2004. Nutritionally mediated placental growth restriction in the growing adolescent: Consequences for the fetus. *Biol. Reprod.* 71:1055-1062.
- Zhu, M.J. 2008. Amp-activated protein kinase signalling pathways are down regulated and skeletal muscle development impaired in fetuses of obese, over-nourished sheep. *Journal of Physiology-London* 586: 2651-2664.