

Contents lists available at [ScienceDirect](http://www.elsevier.com/locate/jtcm)

Journal of Traditional and Complementary Medicine

journal homepage: <http://www.elsevier.com/locate/jtcm>Water and ethanol extracts of *Plantago major* leaves show anti-inflammatory activity on oral epithelial cellsMuhammad Zubair ^{a, b, *}, Cecilia Widén ^{a, c}, Stefan Renvert ^c, Kimmo Rumpunen ^a^a Swedish University of Agricultural Sciences, Department of Plant Breeding and Biotechnology Balsgård, Fjälkestadvägen 459, SE-291 94 Kristianstad, Sweden^b Department of Bioinformatics and Biotechnology, Government College University (GCU), Faisalabad, Pakistan^c Kristianstad University, School of Health and Society, SE-291 88 Kristianstad, Sweden

ARTICLE INFO

Article history:

Received 15 February 2017

Received in revised form

18 July 2017

Accepted 6 September 2017

Available online 17 August 2018

Keywords:

Common plantain

Herbal medicine

NF-kB

Wound healing

ABSTRACT

Context: The leaves of *Plantago major* have been used for the treatment of wounds and inflammation in folk medicine from prehistoric times. However there is no report on the use of *P. major* to treat inflammation in oral epithelial cell lines.**Objective:** The present study was undertaken to reveal possible anti-inflammatory effects of *Plantago major* leaf extracts on oral epithelial cells *in-vitro*.**Materials and methods:** Water- and ethanol-based extracts of *P. major* leaves were prepared from freeze-dried plant material, and tested *in-vitro* using the oral epithelial cell line H400. The anti-inflammatory activity of *P. major* was tested against *E. coli* lipopolysaccharide (LPS) using the nuclear factor kappa beta (NF-kB) assay.**Results:** Both the water- and the ethanol-based extracts, as well as a combination of the two extracts, showed anti-inflammatory activity. A concentration of 0.1 mg/mL (on dry weight basis) yielded the best results for all extracts.**Discussion and conclusion:** The results show that synergistic effects of both polyphenols and water-soluble compounds (possibly polysaccharides) are responsible for anti-inflammatory activities of *P. major*.© 2017 Center for Food and Biomolecules, National Taiwan University. Production and hosting by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Plantago major leaves have been used for its wound healing and anti-inflammatory effects for centuries. Greek physicians described the traditional use of *P. major* in wound healing and for reducing inflammation already during the first century. There are many compounds in the leaves that may contribute to the anti-inflammatory effects of plant extracts. In *Plantago* species, phenolic compounds have thus been reported to have anti-inflammatory activity.^{1,2}

Numerous methods have been used to study the anti-

inflammatory activity of plant extracts. Nuclear factor kappa-B (NF-kB) assay is a commonly used *in-vitro* assay to reveal anti-inflammatory activity of both crude extracts and pure compounds.^{3,4} The NF-kB assay is based on recognition of microbial products by Toll-like receptors (TLRs), that have been shown to be present in e.g. oral epithelial cells.⁵ These receptors activate the NF-kB intracellular pathway that subsequently triggers the inflammatory and immune response processes. This assay investigates the translocation of NF-kB from cytosol to nucleus as a result of inflammation. The difference between the presence of NF-kB in nucleus and cytoplasm describes the degree of inflammation. The higher the nuclear and cytoplasmic differences, the higher is the activation of the NF-kB intracellular pathway, and the higher is also the inflammation. Studies on animal models have confirmed a correlation between NF-kB activation and inflammatory disease,^{6,7} and the assay is thus highly relevant for study of anti-inflammatory potential.

The aim of the present study was to evaluate the anti-

* Corresponding author. Department of Bioinformatics and Biotechnology, Government College University (GCU), Faisalabad, Pakistan.

E-mail addresses: muhammad.zubair1751@gmail.com, zoobi_pbg@yahoo.com (M. Zubair).

Peer review under responsibility of The Center for Food and Biomolecules, National Taiwan University.




Figure 1. Immunocytochemical analysis (nuclear cytoplasmic differences) of NF- κ B activity in OEC H400 following 47 hours treatment with different *Plantago major* extracts of dried leaves (water and ethanol-based extracts and a combination of equal volumes of the two extracts) each in three different concentrations (1.0, 0.1 and 0.01 mg/mL) and 1 hour of exposure to *E. coli* LPS 0.02 μ g/mL. PBS was used as a negative control (nc).

inflammatory activity of different extracts of *P. major* leaves on oral epithelial cells *in-vitro*.

2. Results and discussion

We found that the water- and the ethanol-based extracts as well as the combination of the two extracts, had a significant anti-inflammatory activity as revealed by the NF- κ B assay (Fig. 1). NF κ B is normally present in the cytoplasm, and translocation from the cytoplasm to the nucleus is a definitive measure of NF κ B activation. In this assay, translocation of the activated transcription factor from the cytoplasm to the nucleus is automatically detected and quantified for each cell. The ratio of intensity between the cytoplasm and nucleus is shown in Fig. 1. This confirms that this is a translocation event as it indicates a drop in cytoplasmic intensity along with an increase in nuclear intensity.

The concentration of 0.1 mg/mL (with a content of 1.24, 0.19 and 0.71 μ g plantamajoside and a content of total phenols of 1.67, 0.22 and 0.94 μ g GAE per mL for ethanol-based, water-based and the combination of both extracts, respectively) had the highest anti-inflammatory activity on the oral epithelial cells (OEC H400). The anti-inflammatory response was reduced or absent both at the lower and higher concentrations tested.

The fact that the content of plantamajoside and total phenols in the ethanol-based extract at a concentration of 0.01 mg/mL (0.12 μ g plantamajoside and a content of total phenols of 0.17 μ g GAE per mL) is comparable to the contents of plantamajoside and total phenols in the 0.1 mg/mL water-based extracts but there is significant difference between the *in-vitro* anti-inflammatory response of both extracts show that not only plantamajoside and total phenols but also other water-soluble compounds (possibly polysaccharides) are responsible for the anti-inflammatory activities and ultimately wound healing activities of *P. major*. However at a concentration of 1.0 mg/mL both the ethanol-based and the combination of the water- and ethanol-based extracts showed anti-inflammatory activity.

Our results revealing anti-inflammatory activity of *P. major* leaf extracts are in accordance with⁸ albeit with another kind of cells and another type of anti-inflammatory reaction. They have verified anti-inflammatory activities of methanol extracts of *P. major* and *P. lanceolata* leaves by the COX-1 (cyclooxygenase, expressed constitutively) and 12-LOX (lipooxygenase), enzyme assay and human platelets.

We have in a previous study found that plantamajoside is the major polyphenol in ethanol extracts of *P. major* leaves.^{9,10} It is

therefore possible that this compound is contributing to the anti-inflammatory effect noticed in the present study. Anti-inflammatory activity can be one of the ways in which extracts of *P. major* contribute to wound healing as proposed by.¹¹ Extracts of *P. major* leaves have previously shown effects on the formation of granulation tissue by stimulating the cell proliferation and migration process.¹²

The results of the present study, in combination with results from our previous studies, corroborate the wound healing effects of *P. major* as expected from its traditional use.

3. Experimental

The experimental section is available in [supplementary material](#).

4. Conclusion

We report for the first time that extracts of *P. major* leaves can protect against inflammatory mediators on human oral epithelial cell lines H400 *in-vitro*. Both ethanol-based and water extracts of leaves protected against inflammatory mediators. This makes *P. major* an interesting source of bioactive substances with anti-inflammatory activity. The authors are aware of that additional cell-lines should be tested to corroborate the conclusions. The results justify further studies to elucidate the role of specific compounds responsible for this anti-inflammatory activity. A careful optimization of the concentration of water and ethanol-based extracts and the combination of both these extracts, is also needed.

Acknowledgements

Thanks are due to Professor I.L.C. Chapple and Dr. M.R. Milward, University of Birmingham, UK, for providing the cell line and for sharing technical skills. Support from the technical staff at SLU, Balsgård, Sweden and Dr. M. Asaf Khan, Government College University, Faisalabad, Pakistan is also much appreciated. The authors would also like to acknowledge the Higher Education Commission, Pakistan (HEC), Pakistan and the Ekhaga foundation, Sweden (project number 15931100) for financial support.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jtcme.2017.09.002>.

References

1. Murai M, Tamayama Y, Nishibe S. Phenylethanoids in the herb of *Plantago lanceolata* and inhibitory effect on arachidonic acid-induced mouse ear edema. *Planta med.* 1995;Oct;61:479–480. Epub 1995/10/01.
2. Samuelsen AB. The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review. *J Ethnopharmacol.* 2000;Jul;71:1–21. Epub 2000/07/25.
3. Milward MR, Chapple IL, Wright HJ, Millard JL, Matthews JB, Cooper PR. Differential activation of NF-kappaB and gene expression in oral epithelial cells by periodontal pathogens. *Clin Exp Immunol.* 2007;May;148:307–324. Epub 2007/03/16.
4. Sharif O, Bolshakov VN, Raines S, Newham P, Perkins ND. Transcriptional profiling of the LPS induced NF-kappaB response in macrophages. *BMC Immunol.* 2007;8:1. Epub 2007/01/16.
5. Ren L, Leung WK, Darveau RP, Jin L. The expression profile of lipopolysaccharide-binding protein, membrane-bound CD14, and toll-like receptors 2 and 4 in chronic periodontitis. *J periodontol.* 2005;Nov;76:1950–1959. Epub 2005/11/09.
6. Miagkov AV, Kovalenko DV, Brown CE, et al. NF-kappaB activation provides the potential link between inflammation and hyperplasia in the arthritic joint. *Proc Natl Acad Sci U. S. A.* 1998;Nov 10(95):13859–13864. Epub 1998/11/13.
7. Poynter ME, Irvin CG, Janssen-Heininger YM. A prominent role for airway epithelial NF-kappa B activation in lipopolysaccharide-induced airway inflammation. *J Immunol.* 2003;Jun 15(170):6257–6265. Epub 2003/06/10.
8. Beara IN, Orcic DZ, Lesjak MM, Mimica-Dukic NM, Pekovic BA, Popovic MR. Liquid chromatography/tandem mass spectrometry study of anti-inflammatory activity of plantain (*Plantago* L.) species. *J Pharm Biomed anal.* 2010;Sep 5(52):701–706. Epub 2010/03/12.
9. Zubair M, Nybom H, Ahnlund M, Rumpunen K. Detection of genetic and phytochemical differences between and within populations of *Plantago major* L. (plantain). *Sci Hortic.* 2012b;136:9–16.
10. Zubair M, Nybom H, Lindholm C, Rumpunen K. Major polyphenols in aerial organs of greater plantain (*Plantago major* L.), and effects of drying temperature on polyphenol contents in the leaves. *Sci Hortic.* 2011;128:523–529.
11. Mahmood AA, Phipps ME. Wound healing activities of *Plantago major* leaf extract in rats. *Int J Trop Med.* 2006;1:33–35.
12. Zubair M, Ekholm A, Nybom H, Renvert S, Widen C, Rumpunen K. Effects of *Plantago major* L. leaf extracts on oral epithelial cells in a scratch assay. *J Ethnopharmacol.* 2012a;Jun 14(141):825–830. Epub 2012/04/03.