

広島大学学術情報リポジトリ
Hiroshima University Institutional Repository

Title	Effects of calcium salts of long-chain fatty acids and rumen-protected methionine on plasma concentrations of ghrelin, glucagon-like peptide-1 (7 to 36) amide and pancreatic hormones in lactating cows
Author(s)	Fukumori, Rika; Sugino, Toshihisa; Shingu, Hiroyuki; Moriya, Naoko; Hasegawa, Yoshihisa; Kojima, Masayasu; Kangawa, Kenji; Obitsu, Taketo; Kushibiki, Shiro; Taniguchi, Kouzo
Citation	Domestic Animal Endocrinology , 42 (2) : 74 - 82
Issue Date	2012
DOI	10.1016/j.domaniend.2011.09.005
Self DOI	
URL	http://ir.lib.hiroshima-u.ac.jp/00034813
Right	(c) 2012 Elsevier Inc. All rights reserved.
Relation	

Title: Effects of calcium salts of long-chain fatty acids and rumen-protected methionine on plasma concentrations of ghrelin, glucagon-like peptide-1 (7-36) amide and pancreatic hormones in lactating cows

Authors: R. Fukumori,^a T. Sugino,^{a,*} H. Shingu,^b N. Moriya,^b Y. Hasegawa,^c M. Kojima,^d K. Kangawa,^e T. Obitsu,^a S. Kushibiki,^b K. Taniguchi^a

Institute, address, country: ^aGraduate School of Biosphere Science, Hiroshima University, Higashi-Hiroshima, 739-8528, Japan, ^bNational Institute of Livestock and Grassland Science, Tsukuba, 305-0901, Japan, ^cSchool of Veterinary Medicine and Animal Science, Kitasato University, Towada, 034-8628, Japan, ^dInstitute of Life Science, Kurume University, Kurume, 839-0864, Japan, ^eNational Cardiovascular Center Research Institute, Osaka, 565-8565, Japan

***Correspondence:** Toshihisa Sugino, Graduate School of Biosphere Science, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8528, Japan.

Phone/Facsimile: +81-82-7956

Abstract

Our objective was to determine the effects of calcium salts of long-chain fatty acids (CLFA) and rumen-protected methionine (RPM) on plasma concentrations of ghrelin, glucagon-like peptide-1 (7-36) amide (GLP-1) and pancreatic hormones in lactating cows. Four midlactation Holstein cows were used in a 4 x 4 Latin square experiment in each 2-wk periods. Cows were fed corn silage-based diets with supplements of CLFA (1.5% added on dry matter basis), RPM (20 g/d), CLFA plus RPM and without supplement. Jugular blood samples were taken from 1 h before to 2 h after morning feeding at 10 min intervals on day 12 of each period. CLFA decreased DMI, but RPM did not affect DMI. Both supplements of CLFA and RPM did not affect metabolizable energy intake and milk yield and composition. Plasma NEFA, triglyceride (TG) and total cholesterol (T-Cho) concentrations were increased with CLFA alone, but increases of plasma TG and T-Cho concentrations were moderated by CLFA plus RPM. CLFA increased plasma ghrelin concentration and the ghrelin concentration with CLFA plus RPM was the highest among the treatments. Plasma GLP-1, glucagon and insulin concentrations were decreased with CLFA, whereas adding RPM moderated the decrease of plasma glucagon concentration by CLFA. These results suggest that the addition of Met to cows given CLFA increases plasma

ghrelin and glucagon concentrations associated with the decrease in plasma TG and T-Cho concentrations.

Key words: Dairy cow; Fatty acid; Ghrelin; Glucagon-like peptide-1 (7-36) amide; Methionine

1. Introduction

Fat supplementation in diets for milk production is valid to increase dietary energy density [1]. However, increasing the dietary fat content often depresses DMI of lactating cows [2]. The mechanism of hypophagia induced by fat supplementation is incompletely understood and assumed to be affected by various factors involving the absorption process and metabolism of fatty acids. Methionine (Met) improves lipid metabolism [3] partially by altering the activities of hormone-sensitive lipase and lipoprotein lipase [4]. In dairy cows, Met is the first limiting amino acid for milk production [5]. Supplementation of Met hydroxyl analog increases milk fat synthesis with hypertriglycemia [6]. Conversely, an insufficiency of Met during the periparturient period results in the development of hepatic lipidosis [7]. These reports

suggest that Met may affect lipid metabolism in dairy cows. In lactating ewes, supplementation of rumen-protected Met (RPM) combined with fat increased DMI, milk yield and milk fat secretion compared with the supplementation of fat only [8], but the effects of Met on plasma metabolite and hormone concentrations were unknown. Recently Relling and Reynolds [9, 10] reported changes in plasma concentrations of some gut hormones such as glucagon-like peptide-1 (7-36) amide (GLP-1) and ghrelin caused by several nutrients in lactating cows, partly because of their role in the regulation of feed intake. Additionally, these gut hormones were reported to influence insulin and glucagon secretion [11, 12]. Therefore, in this study we evaluated the effects of adding calcium salts of long-chain fatty acids (CLFA) and RPM on plasma metabolite and hormone concentrations in lactating cows.

2. Materials and Methods

The procedures used in the present study were carried out in accordance with the principles and guidelines for animal use issued by the National Institute of Livestock

and Grassland Science Animal Care Committee, and which were formulated to comply with Japanese regulations.

Table 1

Animals and Management

One primiparous and three multiparous Holstein cows (143.5 ± 3.9 d in milk, calving number: 1.75 ± 0.25 , initial body weight: 547.3 ± 14.7 kg) were fed four diets formulated to meet the nutrient requirements according to the Japanese Feeding Standard for Dairy Cattle [13]. The ingredients and composition of the diets are presented in Table 1. The cows were managed in individual tie stalls, allowed free access to water, and provided experimental diets twice daily at 0900 and 1800 h. They were milked twice daily before each feeding (0840 and 1740 h), and weighed every week.

Treatments and Experimental Design

The cows were used in a 4×4 Latin square design in each 2-wk period. Four treatments consist of basal diet only (without supplement), supplemented with CLFA made from palm and soybean oil [Megalac R (Church & Dwight Co., Inc., Princeton, NJ, USA)], with RPM [Lactet SP (Nippon Soda Co., Ltd., Tokyo, Japan)], and with

CLFA plus RPM. The CLFA containing 85% fatty acids was added to 1.5% of the diet DM, following a previous study [14]. The RPM containing 67% of D-, L-Met was supplemented at 20 g/d for RPM or CLFA plus RPM treatment. Cows were offered each diet ad libitum, allowing for 15% refusal for the first 10 d for each treatment. For the last 4 d of each period, the cows were fed 95% of ad libitum intake for estimating metabolizable energy intake (MEI). Refusals were weighed daily before each morning feeding (0800 h).

Sampling

Samples of the diets and the refusals were collected and pooled for the last 4 d of each period. Milk samples were collected for the last 4 d of each period, added with sodium azide as preservative, and stored at 4°C until analysis. Blood samples were taken at 12 d from the jugular vein catheter (Argyle 14 G CV catheter kit; Nippon Sherwood Medical Industries Ltd., Tokyo, Japan) inserted on 10 d of each period. Blood samples (8 mL) were taken at 10 min intervals from 0800 to 1100 h. Cows were milked during bleeding period (-20 and -10 min sampling time), and they were fed just after 0 min sampling. Blood samples were collected into heparinized tubes with aprotinin [500 kilo inhibitor unit (KIU) /mL of blood; Trasylol, Bayer Leverkusen,

Germany], and centrifuged at $1,500 \times g$ for 20 min at 4°C . Harvested plasma samples were stored at -80°C until assay.

Sample Analysis

The diet samples and refusals were analysed for DM, CP, NDF, crude fat and crude ash contents according to the procedures of AOAC (1990) [15]. Metabolizable energy contents in the treatment diets were calculated based on NRC (2001) [1].

Milk samples were measured for fat, protein, lactose, TS and SNF by infrared analysis (Milko-Scan 1344 A/BN.; Foss Electric Company, Inc., Hillerod, Denmark).

Plasma ghrelin, insulin and GLP-1 concentrations were measured every 10 min by time-resolved fluoro-immunoassay (TR-FIA).

Assay for bioactive ghrelin was conducted as described previously [16]. Ghrelin concentration was measured by competitive solid-phase immunoassay using europium (Eu)-labeled synthetic bovine ghrelin and polystyrene microtiter strips (Nalge Nunc Int., Tokyo, Japan) coated with anti-rabbit γ -globulin. Intra- and inter-assay coefficients of variation were 1.3 and 1.5%, respectively. Least detectable dose and 50% inhibitory concentration in this assay system were 0.025 and 0.831 ng/mL, respectively.

Insulin assay was conducted as described previously [17]. Insulin concentration

was measured by competitive solid-phase immunoassay using Eu-labeled synthetic bovine insulin and polystyrene microtiter strips coated with anti-guinea pig γ -globulin. Intra- and inter-assay coefficients of variation were 2.2 and 1.8%, respectively. Least detectable dose and 50% inhibitory concentration in this assay system were 0.016 and 1.073 ng/mL, respectively.

GLP-1 concentration was measured by competitive solid-phase immunoassay based on the method described by Sugino et al. [16] using rat GLP-1 (Peptide Institute, Inc., Osaka, Japan), Eu-labeled rat GLP-1, polystyrene microtiter strips coated with anti-rabbit γ -globulin and anti-human GLP-1 rabbit serum (1:20,000; Yanaihara Institute Inc., Shizuoka, Japan). Intra- and inter-assay of coefficients of variation were 1.7 and 4.8%, respectively. Least detectable dose and 50% inhibitory concentration in this assay system were 0.024 and 0.172 ng/mL, respectively.

Plasma glucagon levels were measured every 20 min using a commercially available RIA kit (glucagon assay kit, Daiichi Radioisotope Co. Ltd., Tokyo, Japan). Glucagon concentrations were measured in the same assay, and the intra-assay coefficient of variation was 3.4%. Least detectable dose and 50% inhibitory concentration in this assay system were 15.6 and 275 pg/mL, respectively.

Plasma glucose concentrations were determined every 10 min using a glucose

analyzer (GA-1151; Arkray Co, Ltd., Kyoto, Japan). Plasma β -hydroxy butyrate (BHBA), NEFA, triglyceride (TG), total-cholesterol (T-Cho) and urea nitrogen (UN) concentrations were determined every 20 min using an automated biochemistry analyzer (Beckman Coulter, Inc., Tokyo, Japan).

Statistics

Data for feed intake, milk yield, milk composition and plasma amino acid concentrations were analyzed using the MIXED procedure of SAS (SAS Institute, Inc., Cary, NC, USA). The mixed model included treatment as a fixed effect, and cow and period as random effects. For the statistical analysis of plasma hormone and metabolite concentration, sampling time and sampling time \times treatment were added to the model. Factorial contrasts were used to test the main effects of CLFA supplementation (CLFA vs. non-CLFA), RPM supplementation (RPM vs. non-RPM), and their interaction. Results are reported as least squares means and SEM. Significant differences were set at $P < 0.05$.

3. Results

Feed Intake, Milk Yield, and Milk Composition

DMI, milk yield and milk composition are presented in Table 2. DMI in cows fed CLFA diets compared with cows fed non-CLFA diets was lower ($P = 0.025$), whereas MEI was not affected. There was no significant effect of RPM on DMI and MEI. Milk yield and composition, except for total solid contents, were unaffected by CLFA and RPM. Milk total solid contents tended to be lower in cows fed CLFA diets compared with cows fed non-CLFA diets ($P = 0.107$).

Table 2

Plasma Concentration of Hormones and Metabolites

The changes in plasma concentration of hormones and metabolites are presented in Figures 1 and 2, respectively. Plasma ghrelin, insulin, glucose, BHBA, NEFA, TG and UN concentrations varied largely through time ($P < 0.002$, Figures 1 and 2), and no significance of the interaction time by treatment was observed ($P > 0.192$). The time effect for plasma ghrelin, glucose and NEFA concentrations was due to a postprandial decrease. Plasma insulin, BHBA, TG and UN concentrations increased after feeding.

The means of plasma ghrelin concentration were higher ($P < 0.001$, Table 3) in cows fed CLFA compared with cows not fed CLFA. Compared with cows fed

non-CLFA, plasma glucagon and insulin concentrations in cows fed CLFA were lower ($P = 0.002$ and $P = 0.012$, respectively), and plasma GLP-1 concentration tended to be lower ($P = 0.061$). Plasma insulin concentration tended to be higher ($P = 0.092$) in cows supplemented RPM than those without RPM. There were interactions between CLFA and RPM for ghrelin and glucagon ($P = 0.002$ and $P = 0.041$, respectively): CLFA plus RPM increased plasma ghrelin, glucagon, but RPM alone did not show such effects.

Compared to those without CLFA, plasma NEFA, TG, T-Cho and UN concentrations in cows fed CLFA were higher ($P < 0.001$, $P = 0.029$, $P < 0.001$, and $P < 0.001$, respectively), but with lower plasma BHBA concentration ($P = 0.006$, Table 3). In cows fed RPM diets, plasma glucose concentration was higher ($P = 0.001$), plasma T-Cho concentration was lower ($P = 0.009$), and plasma NEFA and UN concentration tended to be lower ($P = 0.15$ and $P = 0.121$, respectively). Interactions between CLFA and RPM were observed for TG and T-Cho ($P = 0.049$ and $P = 0.003$, respectively): RPM decreased plasma TG and T-Cho concentrations in cows fed CLFA, but did not affect the cows not fed with CLFA.

Table 3
Figure 1
Figure 2

4. Discussion

High fat inclusion in diets reduces fiber digestion, increases fatty acid absorption, inhibits abomasal motility [18] and increases gut hormone secretion [19, 20]. Consequently, DMI tends to be depressed. CLFA could prevent such negative effects on ruminal fermentation and fiber digestibility in lactating cows [21, 22]. However, DMI was decreased by CLFA in this study consistent with other reports [23, 24]. RPM did not improve such DMI depression by CLFA contrary to results found by Goulas et al. [8] who used lactating ewes. Additionally, Chillard and Doreau [25] observed no improvement effect of RPM on DMI of cows fed a fish oil supplemented diet during midlactation, although such non-protected fat including high polyunsaturated fatty acids could largely inhibit dietary fiber digestion and depress DMI.

The increase in plasma glucose concentration of cows fed RPM was consistent with Berthiaume et al. [26], while inconsistent with Bertics and Grummer [27]. Such discrepancies might be explained by the differences in the physiological state of cows (mid vs. early lactation) and energy balance. Plasma NEFA, T-Cho and TG concentrations increased by CLFA in the current study has been previously observed in cows [19, 28]. On the other hand, RPM in cows fed CLFA decreased plasma T-Cho and TG concentrations and tended to decrease plasma NEFA concentration. Met is a

methyl group donor for phosphatidylcholine in dairy cows [29] to enhance plasma lipoprotein. In laboratory animals, casein (a Met-rich protein) or L-Met induced hypercholesterolemia in rabbits and rats [30, 31]. In calves, however, high fat diet supplemented with L-Met did not increase plasma VLDL concentration [32], and Met hydroxyl analog did not affect hepatic TG accumulation in cows [27]. Whether Met enhanced VLDL that was not measured in this study is unknown. On the other hand, because Met is converted to taurine in dairy cows, plasma taurine concentration increased linearly with an increase of postprandial Met infusion [33]. Taurine conjugates with bile acids to become taurocholate in the liver, and promotes lipid absorption and cholesterol consumption [34]. Yagasaki et al. [3] showed that dietary Met and Gly reduced serum T-Chol and increased fecal sterol excretion in rats. Furthermore, Met, as a sulfur amino acid, suppressed serum NEFA and TG concentration through the control of hormone-sensitive lipase activity and the restoration of lipoprotein lipase activity in peripheral tissues of hepatoma-bearing rats [4]. In this study, therefore, suppression of plasma T-Chol, NEFA and TG with CLFA plus Met might be induced by enhanced uptake and use of lipids by various tissues.

The increase in plasma ghrelin concentration of cows fed CLFA is consistent with that of rats that ingested some fatty acids [35]. In addition, RPM decreased plasma

ghrelin concentration in cows not fed CLFA, whereas CLFA plus RPM compared with CLFA alone tended to increase plasma ghrelin concentration. Thus, only a simultaneous inclusion of CLFA and RPM may enhance ghrelin secretion via the changes in lipid metabolism. Ghrelin enhances food intake in both nonruminants [36] and ruminants [37, 38]. As a result, we could not find the relationship between plasma ghrelin concentration and feed intake in this study. Since reports on ghrelin responses to lipid metabolism in ruminants are limited, further research is desirable.

Many researchers have reported that high fat supplementation increased GLP-1 concentration [20, 39, 40]. Litherland et al. [40] demonstrated that abomasal infusion of fat increased plasma GLP-1 concentration dependent on dosage. Compared with others, the decrease of plasma GLP-1 concentration with CLFA in this study could be attributed to a lesser fat ingestion level of 280 g/d. GLP-1 plays a role in food intake [41] and Relling and Reynolds [20] have reported that dietary fats increased GLP-1 in dairy cows. Additionally, Relling et al. [42] observed that intrajugular infusion of GLP-1 tended to decrease DMI in growing wethers. In this study however, plasma GLP-1 concentration was decreased by CLFA, suggesting that GLP-1 did not mediate DMI.

Plasma insulin concentration decreased in cows fed CLFA with decreasing DMI

consistent with previous reports [23, 43]. We consider that insulin depression might be due to lessor volatile fatty acids (VFA) production caused by decreased DMI. Although we did not determine plasma VFA concentrations, CLFA decreased plasma BHBA concentration, derived from butyrate by rumen fermentation. And, we also observed that plasma GLP-1 concentration tended to decrease at the same time, suggesting decreased GLP-1, which stimulates insulin secretion, may be related to insulin depression in cows fed CLFA.

CLFA also decreased plasma glucagon concentration. Our result is consistent with that of Cummins and Sartin [43], but inconsistent with that of Khorasani and Kennelly [44]. In several species, elevated NEFA decreases plasma glucagon concentration [45, 46]. However, the present results showed that the decrease of plasma glucagon with CLFA was attenuated by RPM. In our previous study, glucagon secretion was more strongly enhanced when amino acids and ghrelin were simultaneously administered compared with the administration of ghrelin alone in lactating cows [11]. In addition, RPM tended to depress plasma NEFA concentration of cows fed CLFA. Therefore, we consider that increased Met absorption, higher plasma ghrelin concentration and a decreased plasma NEFA concentration might modulate plasma glucagon depression in CLFA-fed cows.

The present study did not show any favorable effects of RPM on feed intake and milk production in cows fed with CLFA although plasma ghrelin concentration increased. Further studies are desirable to elucidate the effects of lactating stage and Met supplemental level with CLFA on DMI and milk production.

In conclusion, responses to CLFA are associated with increases in plasma concentration of ghrelin and decreases of insulin, glucagon and GLP-1. RPM plus CLFA modulated plasma lipid concentrations and concomitantly elevated plasma ghrelin concentration. These metabolic and endocrine changes were induced by absorbed long-chain fatty acid and methionine and not by energy intake because MEI was not different among treatments.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research No. 18780202 from the Japan Society for the Promotion of Science. Rumen-protected methionine and calcium salts of long-chain fatty acids used in this study were donated by Nippon Soda Co., Ltd. (Tokyo, Japan) and Nisso Shoji Co., Ltd. (Tokyo, Japan), respectively. We thank Dr. Lawrence M. Liao, Graduate School of Biosphere Science, Hiroshima

University (Higashi-Hiroshima, Japan), for critical reading of the manuscript.

References

- [1] National Research Council. Nutrient Requirements of Dairy Cattle. 7th Rev. ed. Washington, DC: Natl. Acad. Press; 2001.
- [2] Allen MS. Effects of diet on short-term regulation of feed intake by lactating dairy cattle. *J Dairy Sci* 2000;83:1598-1624.
- [3] Yagasaki M, Machida M, Funabiki R. Effects of dietary methionine and glycine on serum lipoprotein profiles and fecal sterol excretion in normal and hepatoma-bearing rats. *J Nutr Sci Vitaminol* 1989;36:45-54.
- [4] Kawasaki M, Miura Y, Funabiki R, Yagasaki M. Comparison of the effects on lipid metabolism of dietary methionine and cystine between hepatoma-bearing and normal rats. *Biosci Biotechnol Biochem* 2010;74:158-167.
- [5] Schwab CG, Satter LD, Clay AB. Response of lactating cows to abomasal infusion of amino acids. *J Dairy Sci* 1976;59:1254-1270.
- [6] Huber JT, Eney RS, Bergen WG, Liesman JS, Kung L Jr., King KJ, Gardner RW, Checketts M. Influences of methionine hydroxyl analog on milk and milk fat

production, blood serum lipids, and plasma amino acids. *J Dairy Sci* 1984;67:2525-2531.

[7] Shibano K, Kawamura S. Serum free amino acid concentration in hepatic lipidosis of dairy cows in the periparturient period. *J Vet Med Sci* 2005;68:393-396.

[8] Goulas C, Zervas G, Papadopoulos G. Effect of dietary animal fat and methionine on dairy ewes milk yield and milk composition. *Anim Feed Sci Technol* 2003;105:43-54.

[9] Relling AE, Reynolds CK. Abomasal infusion of casein, starch and soybean oil differentially affect plasma concentrations of gut peptides and feed intake in lactating cows. *Domest Anim Endocrinol* 2008;35:35-45.

[10] Relling AE, Reynolds CK. Plasma ghrelin and oxyntomodulin concentrations in lactating cows receiving abomasal soybean oil, corn starch, and casein infusions. *Domest Anim Endocrinol* 2010;38:284-288.

[11] Fukumori R, Yokotani A, Sugino T, Itoh F, Kushibiki S, Shingu H, Moriya N, Hasegawa Y, Kojima M, Kangawa K, Obitsu T, Taniguchi K. Effects of amino acids infused into the vein on ghrelin induced GH, insulin and glucagon secretion in lactating cows. *Anim Sci J* 2011;82:267-273.

[12] Hare KJ, Vilsbøll T, Asmar M, Deacon CF, Knop FK, Holst JJ. The glucagonostatic

and insulinotropic effects of glucagon-like peptide-1 contribute equally to its glucose-lowering action. *Diabetes* 2010;59:1765-1770.

[13] Agriculture, Forestry, and Fisheries Research Council Secretariat. Japanese Feeding Standards for Dairy Cattle, Tokyo, Japan, 2006.

[14] Bilby TR, Block J, do Amaral BC, Sa Filho O, Silvestre FT, Hansen PJ, Staples CR, Thatcher WW. Effects of dietary fatty acids on oocyte quality and follicular development in lactating dairy cow in summer. *J Dairy Sci* 2006;89:3891-3903.

[15] AOAC. Official methods of analysis. 15th ed. Arlington, VA: Association of Official Analytical Chemists; 1990.

[16] Sugino T, Hasegawa Y, Kurose Y, Kojima M, Kangawa K, Terashima Y. Effects of ghrelin on food intake and neuroendocrine function in sheep. *Anim Reprod Sci* 2004;82-83:183-194.

[17] Takahashi H, Kurose Y, Kobayashi S, Sugino T, Kojima M, Kangawa K, Hasegawa Y, Terashima Y. Ghrelin enhances glucose-induced insulin secretion in scheduled meal-fed sheep. *J Endocrinol* 2006;189:67-75.

[18] Nicholson T, Omer SA. The inhibitory effect of intestinal infusions of unsaturated long-chain fatty acids on forestomach motility of sheep. *Brit J Nutr* 1983;50:141-149.

- [19] Choi BR, Palmquist DL, Allen MS. Cholecystokinin mediates depression of feed intake in dairy cattle fed high fat diet. *Domest Anim Endocrinol* 2000;19:159-175.
- [20] Relling AE, Reynolds CK. Feeding rumen-inert fats differing in their degree of saturation decreases intake and increases plasma concentrations of gut peptides in lactating cows. *J Dairy Sci* 2007;90:1506-1515.
- [21] Schauff DL, Clark JH. Effects of prilled fatty acids and calcium salts of fatty acids and calcium salts of fatty acids on rumen fermentation, nutrient digestibilities, milk production, and milk composition. *J Dairy Sci* 1989;72:917-927.
- [22] Schneider PL, Sklan D, Chalupa W, Kronfeld DS. Feeding calcium salts of fatty acids to lactating cows. *J Dairy Sci* 1988;71:2143-2150.
- [23] Choi BR, Palmquist DL. High fat diets increase plasma cholecystokinin and pancreatic polypeptide, and decrease plasma insulin and feed intake in lactating cows. *J Nutr* 1996;126:2913-2919.
- [24] Schauff DJ, Clark JH. Effects of feeding diets containing calcium salts of long-chain fatty acids to lactating dairy cows. *J Dairy Sci* 1992;75:2990-3002.
- [25] Chilliard Y, Doreau M. Influence of supplementary fish oil and rumen-protected methionine on milk yield and composition in dairy cows. *J Dairy Res*

1997;64:173-179.

- [26] Berthiaume R, Dubreuil P, Stevenson M, McBride BW, Lapierre H. Intestinal disappearance and mesenteric and portal appearance of amino acids in dairy cows fed ruminally protected methionine. *J Dairy Sci* 2001;84:194-203.
- [27] Bertics SJ, Grummer RR. Effects of fat and methionine hydroxyl analog prevention or alleviation of fatty liver induced by feed restriction. *J Dairy Sci* 1999;82:2731-2736.
- [28] Gobert M, Martin B, Ferlay A, Chilliard Y, Graulet B, Pradel P, Bauchart D, Durand D. Plant polyphenols associated with vitamin E can reduce plasma lipoperoxidation in dairy cows given n-3 polyunsaturated fatty acids. *J Dairy Sci* 2009;92:6095-6104.
- [29] Pinotti L, Baldi A, Di'Orto V. Comparative mammalian choline metabolism with emphasis on high-yielding dairy cows. *Nutr Res Rev* 2002;15:315-331.
- [30] Khosla P, Samman S, Carroll KK, Huff MW. Turnover of ¹²⁵I-VLDL and ¹³¹I-LDL apolipoprotein B in rabbits fed diets containing casein or soy protein. *Biochim Biophys Acta* 1989;1002:157-163.
- [31] Oda H, Fukui H, Hitomi Y, Yoshida A. Alteration of serum lipoprotein metabolism by polychlorinated biphenyl and methionine in rats fed a soybean protein diet. *J*

Nutr 1991;121:925-933.

- [32] Auboiron S, Durand D, Bauchart D, Robert J, Chapman MJ. Lipoprotein metabolism in the preruminant calf: Effect of a high fat diet supplemented with L-methionine. *J Dairy Sci* 1993;77:1870-1881.
- [33] Pisulewski H, Rulquin H, Peyraud JL, Verite R. Lactational and systemic responses of dairy cows to postprandial infusions of increasing amounts of methionine. *J Dairy Sci* 1996;79:1781-1791.
- [34] Yokogoshi H, Oda H. Dietary taurine enhances cholesterol degradation and reduce serum and liver cholesterol concentrations in rats fed a high-cholesterol diet. *Amino Acids* 2002;23:433-439.
- [35] Nishi Y, Hiejima H, Hosoda H, Kaiya H, Mori K, Fukue Y, Yanase T, Nawata H, Kangawa K, and Kojima M. Ingested medium-chain fatty acids are directly utilized for the acyl modification of ghrelin. *Endocrinology* 2005;146:2255-2264.
- [36] Nakazato M, Murakami N, Date Y, Kojima M, Matsuo H, Kangawa K, Matsukura S. A role for ghrelin in the central regulation of feeding. *Nature* 2001;409:194-198.
- [37] Harrison JL, Miller DW, Findlay PA, Adam CL. Photoperiod influences the central effects of ghrelin on food intake, GH and LH secretion in sheep. *Neuroendocrinology* 2008;87:182-192.

- [38] Wertz-Lutz AE, Knight TJ, Pritchard RH, Daniel JA, Clapper JA, Smart AJ, Trenkle A, Beitz DC. Circulating ghrelin concentrations fluctuate relative to nutritional status and influence feeding behavior in cattle. *J Anim Sci* 2006;84:3285-3300.
- [39] Benson JA, Reynolds CK. Effects of abomasal infusion of long-chain fatty acids on splanchnic metabolism of pancreatic and gut hormones in lactating dairy cows. *J Dairy Sci* 2001;84:1488-1500.
- [40] Litherland NB, Thire S, Beaulieu AD, Reynolds CK. Dry matter intake is decreased more by abomasal infusion of unsaturated free acids than by unsaturated tryglycerides. *J Dairy Sci* 2005;88:632-643.
- [41] Turton MD, Shea D, Gunn I, Beak SA, Edwards CMB, Wilding KJPH, Smith DM, Ghatei MA, Herbert J, Bloom SR. A role for glucagon-like peptide-1 in the central regulation of feeding. *Nature* 1995;379:69-72.
- [42] Relling AE, Reynolds CK, Loerch SC. Effects of feeding fat or intrajugular infusion of glucagon-like peptide-1 and cholecystokinin on dry matter intake, digestibility, and digesta rate of passage in growing wethers. *J Anim Sci* 2011;89:168-178.
- [43] Cummins KA, Sartin JL. Response of insulin, glucagon, and growth hormone to

intravenous glucose challenge in cows fed high fat diets. J Dairy Sci 1987;70:277-283.

[44] Khorasani G., Kennelly JJ. Effects of added fat on performance, rumen characteristics, and plasma metabolites of midlactation cows. J Dairy Sci 1998;81:2459-2468.

[45] Gerich JE, Charles MA, Grodsky GM. Regulation of pancreatic insulin and glucagon secretion. Ann Rev Physiol 1976;38:353-388.





[46] Lavau M, Fried SK, Susini C, Freychet P. Mechanism of insulin resistance in adipocytes of rats fed a high-fat diet. J Lipid Res 1979;20:8-16.

Figure captions:

Figure 1. Plasma hormone (ghrelin: A, GLP-1: B, glucagon: C and insulin: D) concentrations in cows fed each treatment diet: non-CLFA + non-RPM (○ with a solid line), non-CLFA + RPM (● with a dotted line), CLFA + non-RPM (△ with a solid line), and CLFA + RPM (▲ with a dotted line). Values are expressed as least square of means (n=4). *P*-value for effects of treatment, time and the interaction between treatment and time. The horizontal bar and arrow show the milking period and feeding

time, respectively.

Figure 2. Plasma metabolite (glucose: A, BHBA: B, NEFA: C, TG: D, T-Cho: E and UN: F) concentrations in cows fed each treatment diet: non-CLFA + non-RPM (○ with a solid line), non-CLFA + RPM (● with a dotted line), CLFA + non-RPM (△ with a solid line), and CLFA + RPM (▲ with a dotted line). Values are expressed as least square of means (n=4). *P*-value for effects of treatment, time and the interaction between treatment and time. The horizontal bar and arrow show the milking period and feeding time, respectively.











Table 1. Ingredient and chemical composition of the diets

Item	Non-CLFA	CLFA ^a
Ingredient (DM basis)		
Corn silage (%)	42.2	41.6
Alfalfa hay cubes (%)	7.60	7.48
Sudan grass hay (%)	5.68	5.60
Concentrate ^b (%)	43.9	43.2
Calcium salts of fatty acids ^c (%)	0.00	1.50
Calcium Carbonate (%)	0.60	0.60
Salt (%)	0.01	0.01
Vitamin premix (%)	0.01	0.01
Chemical composition (DM basis)		
OM (%)	93.0	92.7
CP (%)	14.1	13.9
NDF (%)	32.2	31.7
Crude fat (%)	3.24	4.45
ME ^d (Mcal/kg)	2.42	2.48

^aLCFA: Calcium salts of long-chain fatty acids.

^bConcentrate contained 39% corn grain, 17% corn gluten feed, 16% beet pulp pe, 10% soybean meal, 6% canola meal, 5% wheat middlings, 5% wheat bran, 2% m

^cMegalac R: Declared fatty acids contained 26% palmitic acid, 4% stearic acid, 33% oleic acid, 32% linoleic acid, and 5% linolenic acid.

^dMetabolizable energy: Estimated value from NRC (2001).

Table 2. Feed intake, milk production and composition during experimental period in lactating cows

Item	Treatment				SEM	Contrast ^c (<i>P</i> -value)		
	non-CLFA		CLFA ^a			CLFA	RPM	CLFA × RPM
	non-RPM	RPM	non-RPM	RPM ^b				
DMI (kg/d)	22.4	22.2	21.3	21.7	0.675	0.025	0.803	0.322
MEI ^d (Mcal/d)	56.7	56.2	55.4	56.4	1.72	0.463	0.794	0.325
Milk yield (kg/d)	27.2	27.1	27.3	27.5	1.83	0.508	0.982	0.655
4% FCM ^e (kg/d)	29.9	28.8	30.0	30.3	1.63	0.505	0.766	0.574
Milk composition								
Fat (%)	4.46	4.35	4.48	4.42	0.44	0.825	0.685	0.909
Protein (%)	3.51	3.56	3.47	3.45	0.132	0.177	0.729	0.508
Lactose (%)	4.62	4.59	4.61	4.61	0.059	0.736	0.743	0.662
Total solid (%)	13.6	13.5	12.7	12.5	1.01	0.107	0.759	0.966
Solid non-fat (%)	9.13	9.11	9.08	9.11	0.124	0.736	0.973	0.710
Body weight changes (kg)	14.3	18.5	13.5	12.7	4.60	0.359	0.612	0.474

Data are shown by least squares means (LSM) and SEM.

^aCLFA: Calcium salts of long-chain fatty acids.

^bRPM: Rumen-protected methionine.

^c*P*-value for factorial contrasts: CLFA, RPM and the interaction between CLFA and RPM.

^dMEI: Metabolizable energy intake.

^e4% FCM: 4% fat corrected milk (kg/d) = 0.4 × Milk yield (kg/d) + 15 × Milk yield (kg/d) × Milk fat (%).

Table 3. Plasma hormone and metabolite concentrations during feeding period in lactating cows

Item	Treatment				SEM	Contrast ^c (<i>P</i> -value)		
	non-CLFA		CLFA ^a			CLFA	RPM	CLFA × RPM
	non-RPM	RPM	non-RPM	RPM ^b				
Hormones								
Ghrelin (ng/mL)	0.106	0.087	0.116	0.126	0.027	<0.001	0.301	0.002
GLP-1 ^d (ng/mL)	0.519	0.464	0.452	0.448	0.128	0.061	0.185	0.258
Glucagon (pg/mL)	99.1	93.6	60.3	85.5	46.2	0.002	0.187	0.041
Insulin (ng/mL)	2.51	2.86	2.36	2.41	0.128	0.012	0.092	0.202
Metabolites								
Glucose (mg/dL)	62.9	63.4	62.1	63.6	1.91	0.872	0.001	0.344
BHBA ^e (μmol/L)	595	579	514	517	43.4	0.006	0.796	0.729
NEFA (μEq/L)	95.4	90.2	116	110	7.13	<0.001	0.150	0.842
TG ^f (mg/dL)	7.57	8.29	9.86	8.41	0.82	0.029	0.501	0.049
T-Cho ^g (mg/dL)	197	200	247	211	14.9	<0.001	0.009	0.003
UN ^h (mgN/dL)	8.79	8.13	9.26	9.23	0.70	<0.001	0.121	0.152

Data are shown by least squares means (LSM) and SEM.

^aCLFA: Calcium salts of long-chain fatty acids.

^bRPM: Rumen-protected methionine.

^c*P*-value for factorial contrasts: CLFA, RPM and the interaction between CLFA and RPM.

^dGLP-1: Glucagon-like peptide-1 (7-36) amide.

^eBHBA: β-hydroxy butyrate.

^fTG: Triglyceride

^gT-Cho: Total-cholesterol.

^hUN = Urea nitrogen.