

麻酔と蘇生 第55巻第1号
 平成31年3月1日発行 昭和47年4月19日第4種学術刊行物認可
 広島麻酔医学会発行 略名：麻と蘇 Anesth Resus

ISSN 0385-1664
 (国際標準逐次刊行物番号)

麻酔と蘇生

Anesthesia and Resuscitation

休刊最終号

Volume 55

Final edition for publication suspension

Number 1

March 2019

目次

臨床研究

頸椎症性脊髄症患者を対象とした挿管用デバイスの前向き比較研究 —エアウェイスコープ™ vs スタイレットスコープ™—	平野 洋子, 他	1
麻酔科開設以来50年間の麻酔管理と使用麻酔薬の変遷	福田 秀樹, 他	5
無線LANにより医療機器からの自動記録が可能になった オープンMRI手術室での25症例の麻酔経験	石井 友美, 他	13
救急救命士が病院到着前に測定した患者の血糖値とその病態の検討	檜崎 壮志, 他	17
小児先天性心疾患手術に伴う肺コンプライアンスの変動に関する調査	北川麻紀子, 他	21
体位変換に伴う循環動態の腹臥位と膝胸位間での比較	三好 寛二, 他	25

症例報告

重症筋無力症患者に対してロクロニウムとスガマデクスを使用した 2症例	大野 麻紀, 他	31
脊髄幹麻酔に起因する脊髄髄節性ミオクロヌスが疑われた2症例	田嶋 実, 他	35
脳深部刺激療法を留置したパーキンソン病患者の脊髄くも膜下麻酔中に ウェアリングオフ様症状と自律神経反射亢進症状を合併した1症例	田嶋 実, 他	39
抗生剤によるビタミンK欠乏性凝固異常のために硬膜外カテーテル 抜去に難渋した3症例	平田 友里, 他	45
胸部硬膜外麻酔によりHorner症候群を呈した1例	横田真優子, 他	49

紹介

第64回 広島麻酔医学会抄録集		53
-----------------	--	----

English Article

CLINICAL ARTICLE

Influence of Maternal Hypotension on Umbilical Artery pH in Parturients Undergoing Cesarean Section	Kana FUKUTOKU, et al	61
Relationship Between Age and Frequency of Side Effects Associated with Postoperative Analgesia	Hiroshi HAMADA, et al	67
Tactile Hypoesthesia Associated with Myofascial Trigger Points in Patients with Persistent Post-Mastectomy Pain —A Close Observation Study in A Case Series—	Katsuyuki MORIWAKI et al	71
Evaluation of Hemodynamics During Posture Change to Knee-Chest Position by FloTrac™	Hirotsugu MIYOSHI, et al	75
Precise Prediction of Right Atrium Position within Expiratory Phase Thorax	Hirotsugu MIYOSHI, et al	79



Evaluation of Hemodynamics During Posture Change to Knee-Chest Position by FloTrac™

Hirotsugu MIYOSHI*¹, Ryuji NAKAMURA*¹, Takashi KONDO*¹, Takahiro KATO*¹,
Toshimichi YASUDA*¹, Hiroyuki TANAKA*², Hiroshi HAMADA*¹
and Masashi KAWAMOTO*¹

Summary : In order to evaluate changes in hemodynamics, a FloTrac™ system was used during posture change from supine to knee-chest position. Thirty-five patients undergoing lumbar surgery participated in this study. Anesthesia was performed with total intravenous anesthesia using propofol and remifentanyl. Cardiac index (CI), stroke volume index (SVI), and stroke volume variation (SVV) were measured by using FloTrac™ in addition with heart rate (HR), systolic blood pressure (SBP), diastolic blood pressure (DBP) and mean blood pressure (MBP). The values before and after postural change were compared. SVV increased immediately after posture change and remained high. With posture change to the knee-chest position, HR, DBP and SVV increased and SBP, CI and SVI decreased. An increase in SVV from immediately after postural change indicated that SVV did not reflect circulating blood volume.

Key words : FloTrac™, knee-chest position, postural change, total intravenous anesthesia

The spine surgery is performed in the prone position. It is known that increase of the intra-abdominal pressure is caused by pressing the abdomen, by which the increased pressure is transmitted to the epidural venous plexus in the prone position. As a result, blood loss will be increased in the prone position.¹⁾ To avoid the pressure to the abdomen, various type of prone positions were invented by using special operating tables and frames, such as Mohammedan praying position,²⁾ modification type of the Tuck position,³⁾ which have the feature of bending the thighs and the knees. Since the body weight was mainly supported by chest and knees, the pressure of the abdomen can be decreased. While knee-chest position has the advantage of decrease of abdominal pressure, we often experience in anesthetized patients whose blood pressure decrease by postural changes to the knee-chest position from the supine position.

The variation of hemodynamics with postural changes from supine to prone have been evaluated by various methods.⁴⁻⁸⁾ However, there is no report to which the evaluation of hemodynamics by changing position from supine position to knee-chest position by using FloTrac™. In this study, we evaluated hemodynamics during the

postural change by using FloTrac™ in patients with lumbar spine surgery.

Materials and Methods

This study was approved by the ethical committee of Asa city hospital, and written informed consent was obtained from each patient. From April 2010 to March 2011, the subjects of the study were adults graded as American Society of Anesthesiologists physical status 1 or 2 who were planned to be operated on the lumbar spine surgery in the knee-chest position according to Mohammedan praying position. Patients with known diabetic neuropathy, uncontrolled hypertension, obesity more than 30 of body mass index and cardiac disease such as atrial fibrillation and aortic regurgitation were excluded.

Anesthetic management protocol

General anesthesia was induced with remifentanyl 0.2–0.5 µg/kg/min, target controlled infusion (TCI) system (TCI pump TE-371, Terumo Corporation, Japan) of propofol 2.0–3.0 µg/ml, and rocuronium 0.6 mg/kg intravenously, and was maintained with remifentanyl 0.2–0.3 µg/kg/min. TCI system of propofol was used to keep concentration within 1.5–2.5 µg/ml. We used Entropy sensor™ system (GE Healthcare, Finland, Helsinki) and kept the response entropy and the state entropy within the range of 40 to 60.

After endotracheal intubation, ventilation was controlled mechanically at 8–10 ml/kg of tidal volume and 10–12 /min

*¹ Department of Anesthesiology and Critical Care, Hiroshima University Hospital, Hiroshima, Japan

*² Department of Anesthesiology and Critical Care, Hiroshima City Asa Citizens Hospital, Hiroshima, Japan

of respiratory rate to maintain end-tidal carbon dioxide in the range of 30 to 40 mmHg using capnography. Tidal volume and respiratory rate were not changed during postural change. Postural change was performed 10 minutes after endotracheal intubation. Acetated Ringer's solution was given at the rate of 10 ml/kg/hour during the study period.

Monitoring

A 22-gauge catheter was inserted into left radial artery during anesthetic induction, and CI, stroke volume index (SVI), stroke volume variation (SVV) were measured using a FloTrac™ system (Edwards Lifesciences, Irvine, CA, USA) and analyzed by the Version.3.02 of Vigileo™ monitor (Edwards Lifesciences, Irvine, CA, USA). Heart rate (HR) and direct measurement of arterial pressure of systolic blood pressure (SBP), diastolic blood pressure (DBP) and mean blood pressure (MBP) were also measured.

Statistical analysis

We measured each value at before postural change and at 3, 5, and 10 minutes after postural change. The values were analyzed by Analysis of variance (ANOVA), and post-hoc test was by Fisher's PLSD. A p-value less than 0.05 were considered to be statistically significant.

Results

Thirty-five patients were participated in this study. The patient's backgrounds are shown in Table 1. The sequential values obtained at each time in HR, SBP, DBP, MBP, CI, SVI, and SVV are shown in Table 2.

Discussion

There are several reports on hemodynamics associated with postural change.⁴⁻⁸ The most universal finding is decreases in CI. This is considered to be decrease in the left ventricular compliance. Increase in the intrathoracic

pressure depresses the venous return according to increased intra-abdominal pressure, and the decrease in venous return is produced because position of the extremities were placed lower than the heart.⁴⁾ The same mechanisms of hemodynamics are predicted in the postural change of the knee-chest position. However, the hemodynamic changes are probably emphasized in the knee-chest position because the limbs placed lower compared with that of prone position. It is reported that the central venous pressure of the knee-chest position is lower than that of the prone position, and this difference is attributable to the position of the lower limb.⁷⁾

In this study, we evaluated the hemodynamic changes of the postural change to the knee-chest position from the supine position by using a FloTrac™ system. In the results of our study, HR, DBP and SVV increased, and SBP, CI and SVI decreased after postural change. We considered that these changes were caused by activated sympathetic nervous system and inhibited parasympathetic activity.⁹⁾ It was also considered that these autonomic nerve reactions took place as the venous return decreased due to postural changes. However, in the previous reports the increase of HR does not occur in postural changes to the prone position.^{4,5,8)} In the knee-chest position, the extremities are placed lower than in the prone position, and changes in hemodynamics are probably emphasized. In the prone

Table. 1

The patient's backgrounds	
Number of patients	35
Surgery	
Lumbar disc hernia	7
Lumbar canal stenosis	28
ASA PS Class 1 : 2	8 : 27
Age (years)	65.3 ± 13.1
Gender man : woman	21 : 14
Height (cm)	159 ± 9.9
Weight (kg)	60.7 ± 9.6
BMI (kg/m ²)	23.7 ± 3.0

Mean ± SD

Table. 2

	Before the postural change	After the postural change			
		3 min	5 min	7 min	10 min
HR (beats/min)	71.8 ± 13.6	78.2 ± 14.1*	78.0 ± 14.0*	75.3 ± 12.4*	73.3 ± 11.8
SBP (mmHg)	115.8 ± 22.8	111.8 ± 23.1	101.9 ± 21.1*	96.0 ± 16.7*	93.2 ± 14.3*
DBP (mmHg)	53.6 ± 11.7	59.6 ± 11.4*	56.8 ± 9.8*	55.1 ± 8.8	54.2 ± 8.4
MBP (mmHg)	74.3 ± 13.8	77.0 ± 13.7	71.9 ± 12.6	68.7 ± 10.5*	67.2 ± 9.8*
CI (L/min/m ²)	2.6 ± 0.5	2.5 ± 0.6	2.2 ± 0.5*	2.1 ± 0.5*	2.0 ± 0.3*
SVI (ml/m ²)	35.9 ± 8.2	32.0 ± 6.9*	28.4 ± 6.1*	26.6 ± 5.3*	27.4 ± 4.5*
SVV (%)	13.4 ± 5.8	21.2 ± 6.5*	20.9 ± 6.7*	19.7 ± 5.8*	17.9 ± 4.9*

Note: Values are given as means ± SD (n = 35)

*p < 0.05 vs. Before the postural change

position, it is reported that the decrease of CI are caused by the decrease in venous return and the decrease in left ventricular compliance due to increased pleural pressure.¹⁾ Similar mechanism seems to be produced in the postural change of the knee-chest position.

In this study, CI decreased with postural change. Even though CI decreased and SVV increased, SVI did not change immediately after postural change. This seemed to be due to SVV not only reflecting changes in the circulating blood volume at the knee chest position. An increase in intrathoracic pressure would be considered as one of the reasons of increases in SVV. In the prone position, it is reported that SVV is able to predict fluid responsiveness in prone during spine surgery.⁸⁾ Because the influence of postural change was apparent in the knee-chest position than that in prone position, the response threshold of SVV for infusion seemed to be higher.

With posture change to the knee-chest position, HR, DBP and SVV increased and SBP, CI and SVI decreased. An increase in SVV from immediately after postural change indicated that SVV seemed to be not reflecting only circulating blood volume.

The abstract was presented at annual meeting of 57th Japanese society of anesthesiologists.

References

- 1) Edgcombe H, Carter K, Yarrow S: Anaesthesia in the prone position. *Br J Anaesth*, 100: 165–183, 2008
- 2) Lipton S: Anaesthesia in the surgery of retropulsed vertebral discs. *Anaesthesia*, 5: 208–212, 1950
- 3) Ecker A: Kneeling position for operations on the lumbar spine; especially for protruded intervertebral disc. *Surgery*, 25: 112, 1949
- 4) Toyota S, Amaki Y: Hemodynamic evaluation of the prone position by transesophageal echocardiography. *J Clin Anesth* 10: 32–35, 1998
- 5) Yokoyama M, Ueda W, Hirakawa M, et al: Hemodynamic effect of the prone position during anesthesia. *Acta Anaesthesiol Scand*, 35: 741–744, 1991
- 6) Hatada T, Kusunoki M, Sakiyama T, et al: Hemodynamics in the prone jackknife position during surgery. *Am J Surg*, 162: 55–58, 1991
- 7) McNulty SE, Weiss J, Azad SS, et al: The effect of the prone position on venous pressure and blood loss during lumbar laminectomy. *J Clin Anesth*, 4: 220–225, 1992
- 8) Biais M, Bernard O, Ha JC, et al: Abilities of pulse pressure variations and stroke volume variations to predict fluid responsiveness in prone position during scoliosis surgery. *Br J Anaesth*, 104: 407–413, 2010
- 9) Mizumaki K, Fujiki A, Tani M, et al: Left ventricular dimensions and autonomic balance during head-up tilt differ between patients with isoproterenol-dependent and isoproterenol-independent neurally mediated syncope. *J Am Coll Cardiol*, 26: 164–173, 1995

Accepted for Publication, January 4, 2019