

Proceedings of the Iowa Academy of Science

Volume 91 | Number

Article 9

1984

Additional Iowa Pteridophyte References

James H. Peck
University of Arkansas - Little Rock

Copyright ©1984 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Peck, James H. (1984) "Additional Iowa Pteridophyte References," *Proceedings of the Iowa Academy of Science*, 91(2), 82-84.

Available at: <https://scholarworks.uni.edu/pias/vol91/iss2/9>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Additional Iowa Pteridophyte References

JAMES H. PECK

Department of Biology, University of Arkansas-Little Rock, Little Rock, AR 72204

Fifty-two annotated references are added to the bibliography of references on Iowa pteridophytes. The bibliography was compiled to include references which treated the taxonomic, ecologic, physiologic or phytogeographic aspects of Iowa's ferns, horsetails, quillworts, clubmosses, and spikemosses. The total bibliography now contains 218 references. The status of the Iowa pteridophyte floristics is compared with that in other states in the Upper Midwest and the Great Plains. The Iowa pteridophyte flora consists of 57 species and 8 hybrids, including 47 ferns, 9 horsetails, 1 quillwort, 6 clubmosses, and 2 spikemosses.

INDEX DESCRIPTORS: Clubmosses, Ferns, Horsetails, Iowa Flora, Pteridophytes, Quillworts, Spikemosses, Bibliography.

The Iowa pteridophyte (ferns and fern allies) flora has been the subject of a field, herbarium, and literature investigation for the past decade (1972-1982). These efforts have resulted in the preparation of a new summation of floristic collections (Peck, 1976b), along with an annotated bibliography to references on Iowa pteridophytes (Peck, 1976a). Since then, continued field and herbarium study has resulted in additional floristic information being reported (Peck, 1980; 1983). Similar studies have been conducted in neighboring states over the past decade as part of their state floristic surveys, endangered species programs, or natural heritage programs. States in the Upper Midwest with recent pteridophyte floras include Illinois (Mohlenbrock, 1967; 1970; Mohlenbrock and Ladd, 1978; Mohlenbrock and Ladd, in press), Minnesota (Tyron, 1980), Missouri (Steyermark, 1963; Key, 1982), and Wisconsin (Peck and Taylor, 1980a; 1980b; 1981). States in the Great Plains to the west of Iowa which have had recent floras include North Dakota (Pelvitt and Barker, 1975) and South Dakota (Van Bruggen, 1976). Two physiographic regions adjacent to Iowa have also been studied, resulted in reports covering the entire Great Plains (Great Plains Flora Association, 1977; Petrik-Ott, 1975; 1979) and the Driftless Area of northeastern Iowa and parts of Illinois, Minnesota, and Wisconsin (Hartley, 1966; Peck, 1982). Maps of pteridophyte distributions in the Upper Midwest (Illinois, Iowa, Minnesota, Missouri, and Wisconsin) were included in the recent

revision of the Driftless Area pteridophyte flora (Peck, 1982). These state and regional treatments, along with the manuals of Wherry (1961) and Mickel (1979) and the new world treatment of pteridophytes by Tyron and Tyron (1982), provide pteridologists and Iowa's botanists with much improved floristic and distributional data. Consequently, studies of the floristic affinities and vicariance geography of Iowa's pteridophytes can now examine historical and ecological influences on species' abundance and patterns of distribution.

Since publication of the Iowa pteridophyte bibliography (Peck, 1976a), 52 additional references to Iowa pteridophytes have been located, including 35 published since 1975 and 17 older references. As before, each reference was annotated to clarify the title, contents, and significance. To date, a total of 218 references to Iowa pteridophytes have been located and annotated. Based upon these references, the Iowa pteridophyte flora consists of 65 pteridophytes (58 species plus 8 hybrids), including 47 ferns, 9 horsetails, 1 quillwort, 6 clubmosses, and 2 spikemosses. The fern *Botrychium dissectum* Spreng. and the quillwort *Isoetes melanopoda* Gay & Dur. occur in Iowa as two distinct forms, making the final flora of Iowa pteridophytes a total of 67 (species, hybrids, plus noteworthy forms). The purpose of this report is to provide a current statement on Iowa pteridophyte literature by updating a previous report (Peck, 1976a) for workers in Iowa, as well as workers in other states, who use Iowa data for floristic and ecologic investigations.

ANNOTATED BIBLIOGRAPHY

- ALLEN, W. H., B. G. GOETTSCHE, and R. K. KENNEDY. 1970. Surficial geology, soils, and plant zonation of Ross Biological Area. Proc. Iowa Acad. Sci. 77:233-248. (*Equisetum hyemale* was reported as a good indicator of high soil moisture, particularly at the foot of slopes, where it was strongly zoned and correlated with high soil moisture).
- ANDERSON, W. A. 1946. Development of prairie at Iowa Lakeside Laboratory. Amer. Midl. Nat. 36:431-455. (Reports on the presence of *Equisetum laevigatum* in the northern portion of the field station for the years 1936, 1944, and 1945 as a component of the prairie flora).
- BUTTERS, F. K. 1917. *Pellaea atropurpurea* (L.) Link and *Pellaea glabella* Mett. ex Kuhn. Amer. Fern J. 7:77-87. (Presents geographic and morphologic differences between these species; also of historical interest, since Butters incorrectly referred to all *P. atropurpurea* in the Driftless Area as *P. glabella*).
- CARR, M. D. 1981. Iowa ferns. Linnaean Fern Bull. 1(3):6-7. (Brief note on the abundance of 13 ferns located in the vicinity of West Union, Fayette County, Iowa).
- CARVEY, K. 1977. The autecology and genetic load of disjunct populations of *Asplenium platyneuron* on coal spoils in Marion County, Iowa. M. S. thesis, Iowa State University, Ames. (Comparisons of isolated, disjunct plants in Iowa with populations in Illinois and Missouri; isolated plants formed from hermaphroditic gametophytes which readily selfed and rapidly matured; coal spoils provided unique habitats, scattered across southcentral to central Iowa).
- CHRISTIANSEN, P., H. HADOW, AND E. HIMMAN. 1980. Natural resources inventory of Pikes Peak/Point Ann State Park, Clayton County, Iowa. Report to Iowa Conservation Commission, Des Moines. 170 pp. (Reports 14 pteridophytes: 1 clubmoss, 2 horsetails, and 11 ferns; community associations).
- CLUTE, W. N. 1911. *Pellaea gracilis* on sandstone. Linnaean Fern Bull. 19:84. (Reports that the first observation of *Cryptogramma stelleri* on sandstone was from a collection at Waukon, Iowa).
- ENGELMANN, G. 1882. The genus *Isoetes* in North America. Trans. St. Louis Acad. Sci. 4:358-390. (Reports that collections of *Isoetes* in Iowa by Vasey in 1863 were the 20th observation of *Isoetes* in North America and the second collection of *Isoetes melanopoda* for North America, preceded only by its discovery and description from specimens collected in 1853 from southern Illinois).
- FARRAR, D. R. 1976. Spore retention and release from overwintering fern fronds. Amer. Fern J. 66:49-52. (Observations on natural populations of ferns which note that spore release and retention varies with respect to species; some species released all spores promptly, while others released spores slowly, and some released spores over a 9 month interval).
- CRIST, K. C., and D. R. FARRAR. 1983. Genetic load and long distance dispersal in *Asplenium platyneuron*. Can. J. Bot. 61:1809-1814. (Iowa populations exhibited remarkable genetic and ecologic adaptations for successful colonization by long distance dispersal, by their possession of low genetic load, regular sporophyte production from isolated gametophytes, and the ability of such plants to release variability through homeologous chromosome pairing in subsequent generation; includes distribution map).
- FINK, B. 1894. Blights, orchids, and ferns of Fayette, Iowa. Bull. Upper Iowa Univ. (Cited in Wilson (1910) on page 48; efforts to locate this document in Iowa libraries, including Upper Iowa University, have not been successful).
- GILMAN, J. C., and W. A. ARCHER. 1929. The fungi of Iowa parasitic on plants. Iowa St. Coll. J. Sci. 3:299-507. (Reports 2 fungi on horsetails and 4 fungi on ferns).
- GLENN-LEWIN, D. C. 1977. The vegetation of Stinson Prairie, Kossuth County, Iowa. Proc. Iowa Acad. Sci. 83:88-93. (Notes 2 horsetails).
- GRAVES, E. W. 1927. A new colony of *Botrychium obliquum* in Iowa. Amer. Fern J. 17:113-116. (Reports on discovery of this species in Van Buren County).

- JENNINGS, V. M., and R. S. FAWCETT. 1977. Weed control: Horsetail (*Equisetum arvense* L.). Cooperative Extension Serv., Iowa State University, Ames. PM-777. (Descriptions, symptoms, treatment, and control measures; erroneously uses *E. laevigatum* to depict fertile stems of *E. arvense*).
- KNUTSON, R. M. 1979. Natural resource study of Volga River State Recreation Area, Fayette County, Iowa: assessment of vegetation. Report to Iowa Conservation Commission, Des Moines. (Reports 13 pteridophytes: 1 clubmoss, 3 horsetails, and 9 ferns).
- LAMMERS, T. G. 1980. The vascular flora of Starr's Cave State Preserve. Proc. Iowa Acad. Sci. 87:148-158. (Reports 2 horsetails and 11 ferns at a state preserve in Des Moines County; provides floristic comparison with other preserves in Iowa).
- LAMMERS, T. G. 1981. The vascular flora of Des Moines County, Iowa. M.A. thesis, University of Northern Iowa, Cedar Falls, IA. (Reports 23 species, including 6 horsetails and 17 ferns).
- LAMMERS, T. G. 1983. The vascular flora of Des Moines County, Iowa. Proc. Iowa Acad. Sci. 90:55-71. (Reports 23 pteridophytes, including 6 horsetails and 17 ferns).
- LAMMERS, T. G., and A. G. VAN DER VALK. 1977. A checklist of the aquatic and wetland vascular plants of Iowa: I. Ferns, fern allies, and dicotyledons. Proc. Iowa Acad. Sci. 84:41-88. (Reports habitats and distribution maps for 16 pteridophytes: 1 spikemoss, 1 quillwort, 4 horsetails, and 10 ferns).
- LAMMERS, T. G., and A. G. VAN DER VALK. 1979. A checklist of the aquatic and wetland vascular plants of Iowa: II. Monocotyledons, plus a summary of the geographic and habitat distributions of all aquatic and wetland species in Iowa. Proc. Iowa Acad. Sci. 85:121-163. (Reports community and habitat preferences of 16 pteridophytes).
- MCGINNIS, M. R. 1969. Selected aspects of the biology of *Hyalospora polypodii* (Pers.) Magn. on *Cystopteris fragilis* (L.) Bernh. Ph. D. dissertation, Iowa State University, Ames. (Phenetic investigations on viability and germination requirements of uredospores using various storage conditions; fungus is a parasite on fern fronds, perpetuating itself in Iowa by asexual reproduction).
- PAMMEL, L. H., and C. M. KING. 1926. The weed flora of Iowa. Iowa Geol. Surv. Bull. 4:1-715. (Lists *Equisetum arvense* as a weed; description, distribution, and suggestions for eradication).
- PECK, J. H. 1976a. An annotated bibliography to the literature on pteridophytes in Iowa. Proc. Iowa Acad. Sci. 82:203-208. (Includes 166 references by 84 authors on Iowa pteridophytes).
- PECK, J. H. 1976b. The pteridophyte flora of Iowa. Proc. Iowa Acad. Sci. 83:143-160. (Reports 57 species and 3 hybrids; habitats, abundance, and distributional data; commentary on history of Iowa pteridophyte study).
- PECK, J. H. 1978. The location and distribution of threatened, endangered, and possibly extirpated Iowa pteridophytes. Report to Iowa Preserves Advisory Board, Iowa Conservation Commission, Des Moines. 90 pp. (Specimen citations, Iowa distribution map, and north-central states distribution maps for 26 pteridophytes).
- PECK, J. H. 1979. Compilatory list of the rare, threatened, and endangered plants of the five states of the Upper Mississippi River: Illinois, Iowa, Minnesota, and Wisconsin. Contr. Herbarium University Wisconsin - La Crosse 26:1-48. (Lists 6 horsetails, 2 quillworts, 7 clubmosses, 3 spikemosses, and 52 ferns of biological concern in the Upper Midwest; Iowa lists 4 horsetails, 1 quillwort, 5 clubmosses, 2 spikemosses, and 14 ferns; discusses idiosyncratic approaches taken by each state in designating and listing species).
- PECK, J. H. 1980a. Additions to the pteridophyte flora of Iowa. Proc. Iowa Acad. Sci. 87:39-40. (Reports two state records and 85 county records; *Dryopteris X triplioidea* Wherry and *Equisetum X litorale* Kuhlw. were reported for first time).
- PECK, J. H. 1980b. *Equisetum X litorale* in Illinois, Iowa, Minnesota, and Wisconsin. Amer. Fern J. 70:33-38. (Reports field survey which discovered the first and second Iowa stations, the second Illinois station, the second, third and fourth Minnesota stations, and eight new localities in Wisconsin; habitat, stand dynamics, and reproductive ecology were investigated; roles of flooding and draw-downs on populations were identified as critical to population occurrence, persistence, and dispersal).
- PECK, J. H. 1980c. The life history and reproductive biology of ferns at Woodman Hollow, Iowa. Ph. D. dissertation, Iowa State University, Ames. (Phytogeographic and microclimate investigation of Woodman Hollow State Preserve, Webster County; field and laboratory study of 14 species of ferns, including sporophyte, gametophyte, spore, and sporeling ecology).
- PECK, J. H. 1982. Ferns and fern allies of the Driftless Area of Illinois, Iowa, Minnesota, and Wisconsin. Milwaukee Public Museum Contr. Bio. Geol. 53:1-140. (Reports on distribution and provides keys to 73 species, 13 hybrids, and 6 infraspecific taxa which comprise the flora of this phytogeographically interesting area; distribution maps of pteridophytes within the Driftless Area and in the north central United States show county occurrence).
- PECK, J. H. 1983. Additions to the pteridophyte flora of Iowa — II. Proc. Iowa Acad. Sci. 90:28-31. (Reports *Dryopteris X bootii* and *Dryopteris X uliginosa* as state records, along with 118 new county records; observations on field status of populations of the 12 rarest species were summarized; reports on the rediscovery of *Isoetes melanopoda* in Iowa which had not been observed since its discovery in 1863).
- PECK, J. H., and W. R. BUCK. 1978. The *Selaginella apoda* complex in Iowa. Amer. Fern J. 68:29. (The single station of a wetland spikemoss in Iowa was determined not to have been extirpated as previously reported; the species is known as *S. elipes* Buck, being found from the St. Lawrence River westward through the Great Lakes states and south to Missouri; *S. apoda* is found to the south of *S. elipes*).
- PECK, J. H., L. J. EILERS, and D. M. ROOSA. 1978. The vascular flora of Fremont County, Iowa. Iowa Birdlife 48:3-18. (Lists 4 horsetails and 5 ferns).
- PECK, J. H., T. G. LAMMERS, B. HAGLAN, D. M. ROOSA, and L. J. EILERS. 1981. A checklist of the vascular flora of Lee County, Iowa. Proc. Iowa Acad. Sci. 88:159-171. (Reports 22 pteridophytes, including 4 horsetails and 18 ferns).
- PECK, J. H., and D. M. ROOSA. 1983. Bibliography of Iowa aquatic and wetland plant literature. Proc. Iowa Acad. Sci. 90:72-77. (Lists 350 references by 210 authors, including references on aquatic and wetland pteridophytes).
- PECK, J. H., D. M. ROOSA, and L. J. EILERS. 1980. A checklist of the vascular plants of Allamakee County, Iowa. Proc. Iowa Acad. Sci. 87:62-75. (Reports 52 pteridophytes, including 1 spikemoss, 5 clubmosses, 8 horsetails, and 38 ferns; habitat observations are reported on the first discovery of *Dryopteris X triplioidea* and *Equisetum X litorale* in Iowa).
- PETRIK-OTT, A. J. 1979. The pteridophytes of Kansas, Nebraska, South and North Dakota, U.S.A. Nov. Hedwigia Beih. 61:1-332. (Manual with keys, descriptions, illustrations, and maps of the Great Plains and North American distributions; erroneously excludes *Cystopteris X tennesseensis* and *Equisetum scirpoides* from the Iowa flora and incorrectly includes *Cheilanthes lanosa*, *Asplenium trichomanes*, *Polypodium hesperium*, *P. polypodioides*, and *Lycopodium obscurum* as occurring in Iowa).
- PLOUFFE, M. E. 1977. An autecological study of *Betula papyrifera* in the Iowa River greenbelt in Hardin County, with notes on the vegetation. M. S. thesis, Iowa State University, Ames. (Investigation on the silvics and phytogeography of peripheral and disjunct populations; lists 29 pteridophytes: 1 clubmoss, 5 horsetails, and 23 ferns).
- ROOSA, D. M., and L. J. EILERS. 1978. Endangered and threatened Iowa vascular plants. Special Report State Preserves Advisory Board 5:1-93. Lists 25 pteridophytes: 17 endangered, 5 threatened, and 3 presumed extirpated).
- RUTZ, L. M. 1982. Biology of the genus *Equisetum*, Subgenus *Hippochaete*, in central Iowa. M. S. thesis, Iowa State University, Ames, IA. (An investigation of the field demographics and experimental analysis of sexual and asexual reproductive biology of *Equisetum hyemale*, *E. laevigatum*, and *E. X ferrissii* to assess their frequency, abundance, and persistence).
- SARVELA, J., D. M. BRITTON, and K. PRYER. 1981. Studies on the *Gymnocarpium robertianum* complex in North America. Rhodora 83:421-431. (Cytomorphotypes of *G. robertianum* and *G. dryopteris* are discussed; map of *G. robertianum* range includes Allamakee Co., Iowa, localities).
- SYLVESTER, E. P. 1965. Horsetail (*Equisetum arvense* L.). Iowa State University Cooperative Extension Service, Weed Control Series, WC-37 Revised Aug. 1965. (Description, symptoms, treatment, and control measures; sterile stems of *Equisetum laevigatum* were erroneously used to represent fertile stems of *Equisetum arvense*).
- TRYON, R. M. 1948. Some Woodsias from the North Shore of Lake Superior. Amer. Fern J. 38:159-170. (Discusses the Iowa collections of *Woodsia oregana* with respect to the taxonomic significance of *W. catbartiana* in the Upper Midwest).
- TRYON, R. M., and A. F. TRYON. 1982. Ferns and allied plants with special reference to Tropical America. Springer-Verlag, New York. 857 pp. (Provides the Neotropical distribution of the genera found in the Western Hemisphere; excellent compendium on fern systematics with general distribution maps of value in interpreting the range of Iowa's pteridophyte genera).
- VANDER ZEE, D. 1977. Gitchee Manitou Preserve — flora, ecology, and management. M.S. thesis, Iowa State University, Ames. (Lists 5 pteridophytes: 1 spikemoss, 2 horsetails, and 2 ferns).
- VANDER ZEE, D. 1980. Vascular flora of Gitchee Manitou State Preserve, Lyon County, Iowa. Proc. Iowa Acad. Sci. 86:66-75. (Lists 5 pteridophytes: 1 spikemoss, 2 horsetails, and 2 ferns).
- VINJE, J. M., and M. M. VINJE. 1955. Preliminary aerial survey of microbiota in the vicinity of Davenport, Iowa. Amer. Midl. Nat. 54:418-432. (Reports trapping a fern sporangium at 3500 feet on an exposed glass slide suspended from an aircraft).
- WAGNER, W. H. 1959. American grapeferns resembling *Botrychium ternatum*: a preliminary report. Amer. Fern J. 49:97-103. (Indicates that *Botrychium oneidense* occurs in Iowa; probably based on specimens that were collected by E. W. Graves and deposited in a non-Iowa herbarium; efforts to locate these vouchers have been unsuccessful).
- WAGNER, W. H., and D. M. JOHNSON. 1981. Natural history of the ebony spleenwort fern, *Asplenium platyneuron* (Aspleniaceae) in the Upper Great Lakes area. Can. Field-Nat. 95:156-166. (Reports on the distribution and ecology of *Asplenium platyneuron* in North America, including Iowa populations; suggests that northern populations are recent and transient in occurrence).
- WEATHERBY, C. A. 1936. E. W. Graves. Amer. Fern J. 26:75-76. (Obituary for E. W. Graves, an amateur pteridologist who retired to Bentonsport, Iowa, and began a study of the genus *Botrychium* in the Upper Midwest, which resulted in the first reports for *B. dissectum* and *B. obliquum* in Iowa, Minnesota, Wisconsin, and Illinois; see references by Graves reported here and in Peck (1976a); he profoundly changed the perception of the occur-

rence of *Botrychium* in the Upper Midwest).
 WILSON, G. W. 1910. Preliminary list of the parasitic fungi of Fayette County, Iowa. *Proc. Iowa Acad. Sci.* 17:47-49. (Reports *Hyalospora* parasitic on *Cystopteris fragilis*).

REFERENCES

- BROOKS, R. E. 1980. Additions to the pteridophyte flora of the Great Plains. *Amer. Fern J.* 70:91-92.
- COOPERRIDER, T. S. 1959. The ferns and other pteridophytes of Iowa. *St. Univ. Iowa Stud. Nat. Hist.* 20(1):1-66.
- GREAT PLAINS FLORA ASSOCIATION. 1977. Atlas of the flora of the Great Plains. Iowa State University Press, Ames, IA.
- HARTLEY, T. G. 1966. The flora of the Driftless Area. *Univ. Iowa Stud. Nat. Hist.* 21(1):1-174.
- KEY, J. S. 1982. Field guide to Missouri ferns. Missouri Dept. Conservation, Jefferson City, MO.
- MICKEL, J. T. 1979. How to know the ferns and fern allies. W. C. Brown, Dubuque, IA.
- MOHLENBROCK, R. H. 1967. The illustrated flora of Illinois: ferns. Southern Illinois University Press, Carbondale, IL.
- MOHLENBROCK, R. H. 1970. Additions to the illustrated flora of Illinois. I. *Trans. Illinois St. Acad. Sci.* 63:42-50.
- MOHLENBROCK, R. H., and D. M. LADD. 1978. Distribution of Illinois vascular plants. Southern Illinois University Press, Carbondale, IL.
- MOHLENBROCK, R. H., and D. M. LADD. In press. Additions to the illustrated flora of Illinois. II. Ferns. *Trans. Illinois St. Acad. Sci.*
- PECK, J. H. 1976a. An annotated bibliography to the literature on pteridophytes in Iowa. *Proc. Iowa Acad. Sci.* 82:203-208.
- PECK, J. H. 1976b. The pteridophyte flora of Iowa. *Proc. Iowa Acad. Sci.* 83:140-160.
- PECK, J. H. 1980. Additions to the pteridophyte flora of Iowa. *Proc. Iowa Acad. Sci.* 87:39-40.
- PECK, J. H. 1982. Ferns and fern allies of the Driftless Area of Illinois, Iowa, Minnesota, and Wisconsin. Milwaukee Public Museum Contr. Biol. Geol. 53:1-140.
- PECK, J. H. 1983. Additions to the pteridophyte flora of Iowa — II. *Proc. Iowa Acad. Sci.* 90:28-31.
- PECK, J. H., and W. C. TAYLOR. 1980a. Checklist and distributions of Wisconsin ferns and fern allies. *Mich. Bot.* 19:251-268.
- PECK, J. H., and W. C. TAYLOR. 1980b. Wisconsin pteridophyte literature. *Bull. Wisconsin Bot. Club* 12(1):2-4.
- PECK, J. H., and W. C. TAYLOR. 1981. Wisconsin pteridophyte literature II. *Bull. Wisconsin Bot. Club* 13:19-23.
- PELVITT, J., and W. T. BARKER. 1975. North Dakota ferns and fern allies. Tri-College Center for Environmental Stud. Pub. Ser. No. 1. North Dakota St. Univ., Fargo, ND.
- PETRIK-OTT, A. J. 1975. A county checklist of the ferns and fern allies of Kansas, Nebraska, South Dakota, and North Dakota. *Rhodora* 77:478-511.
- PETRIK-OTT, A. J. 1979. The pteridophytes of Kansas, Nebraska, South Dakota, and North Dakota. *Nov. Hedwigia Beih.* 61:1-322.
- STEYERMARK, J. A. 1963. Flora of Missouri. Iowa State University Press, Ames, IA.
- TRYON, R. M. 1980. Ferns of Minnesota. 2nd ed. University of Minnesota Press, Minneapolis, MN.
- TRYON, R. M., N. C. FASSETT, D. W. DUNLOP, and M. E. DIEMER. 1953. The ferns and fern allies of Wisconsin. Univ. Wisconsin Press, Madison, WI.
- TRYON, R. M., and A. F. TRYON. 1982. Ferns and allied plants with special reference to Tropical America. Springer-Verlag, New York. 857 pp.
- VAN BRUGGEN, T. 1976. The vascular plants of South Dakota. Iowa State University Press, Ames, IA.
- WHERRY, E. T. 1961. The fern guide. Doubleday and Co., New York, NY.

IN MEMORIAM

Dr. Robert H. Chapman

Dr. Robert H. Chapman, Assistant Professor of Botany, Iowa State University, died February 3, 1984 in Ames, Iowa. He was born in Atlanta, Georgia and received a B.A. degree from Emory University in 1969. Following three years in the Army Security Agency, including 15 months as an interpreter in Viet Nam, he enrolled at the University of Georgia and received an M.S. (1974) and Ph.D. (1977) in Botany. He was a post-doctoral fellow at the University of Rochester but left early to accept an appointment at Iowa State in fall, 1977.

His teaching responsibilities included general biology, biological evolution, and plant population biology. He became especially interested in evolution and, in addition to teaching in the formal evolution course, often spoke to student groups about evolution. He became a spokesman for biologists on the question of what is science and what is

not science in the field of evolutionary biology.

Dr. Chapman was known beyond the campus for his efforts to raise awareness of the impact of the evolution controversy on science. He wrote a chapter for a book on the subject and presented papers about it at meetings. He was a member of the Controversial Issues Panel of the Iowa Academy of Science which dealt with this matter.

His research interests were in evolutionary biology, specifically in the application of electrophoresis techniques to problems in plant population biology. He had published papers in this discipline. For a short period he was Associate Editor for Botany for the *Proceedings of the Iowa Academy of Science*.

Bob Chapman was an amiable and intelligent teacher and scientist. The early end to his career is an unfortunate loss to both Iowa State University and the state of Iowa.