

# Proceedings of the Iowa Academy of Science

---

Volume 78 | Number 3-4

Article 3

---

1972

## Effects of DDT on the Density and Diversity of Tardigrades

Gary W. Barrett  
*Miami University*

Ronald G. Kimmel

Copyright ©1972 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Barrett, Gary W. and Kimmel, Ronald G. (1972) "Effects of DDT on the Density and Diversity of Tardigrades," *Proceedings of the Iowa Academy of Science*, 78(3-4), 41-42.

Available at: <https://scholarworks.uni.edu/pias/vol78/iss3/3>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

Effects of DDT on the Density and Diversity of Tardigrades<sup>1</sup>GARY W. BARRETT<sup>2</sup> and RONALD G. KIMMEL<sup>3</sup>

GARY W. BARRETT and RONALD G. KIMMEL. Effects of DDT on the Density and Diversity of Tardigrades. *Proc. Iowa Acad. Sci.*, 78(3-4):41-42, 1972.

SYNOPSIS. Twenty lichen-bearing tree bark samples were collected from DDT sprayed American elm trees (*Ulmus americana* L.) in central Iowa. Twenty comparable samples were collected from an adjacent nontreated habitat. A tardigrade density of 4 individuals collected from the treated habitat was found to differ significantly ( $P < .01$ ) from a density of 97 individuals collected from the non-treated area. Margalef's diversity index ( $D = S - 1 / \ln N$ ) determinations were found to be 0.00 and 0.44 for the treated and

nontreated habitats, respectively. Tardigrade species organization within the nontreated habitat did not differ significantly from that as predicted by MacArthur's broken-stick model, suggesting that tardigrade species occupy contiguous, non-overlapping niches within a lichen-bearing tree bark ecosystem. It is suggested that measurements depicting microfauna-lichen-tree bark relationships might serve as useful criteria in evaluating pesticide stress effects on total forest ecosystems.

INDEX DESCRIPTORS: Tardigrades, Density, Diversity, Effects of DDT.

The ecological effects of DDT when used in the control of Dutch elm disease have been studied by several investigators (Benton, 1951; Blagbrough, 1952; Barker, 1958; Hunt, 1960; and others). Several studies have also been concerned with the effects of DDT on natural invertebrate populations (Davis, 1952; Menhinick, 1962; Odum, Woodwell, and Wurster, 1969; and others). Many of these findings have been summarized in review articles by Ripper (1956) and Moore (1967). However, no studies have been concerned with the effects of DDT on natural populations of tardigrades. This investigation was designed to measure and evaluate the effects of DDT on the density, diversity, and species organization of tardigrades collected from sprayed American elm trees (*Ulmus americana* L.) in central Iowa.

## MATERIALS AND METHODS

Twenty lichen-bearing tree bark samples, 25 cm<sup>2</sup> each, were collected from DDT sprayed American elm trees in Polk County, Iowa. Twenty comparable samples were collected from an adjacent nontreated habitat. All samples contained lichens belonging to the genera *Physcia*. All samples were collected on 29 September 1967, approximately one meter above ground level on the east side of randomly selected trees. The twenty treated samples were collected from separate trees that had been aerially sprayed with DDT on 5 November 1966 at the rate of 0.5 lb. per acre (Mr. Harold E. Smith, Executive Administrator, Public Works Division, City of Des Moines, Iowa, personal communication).

Tardigrades were removed from the samples by a water flooding technique as described by Kimmel and Meglitsch (1969). Specimens from each sample were fixed mounted in Hoyer's medium and identified as to species.

The  $D = S - 1 / \ln N$  index of Margalef (1957) and the broken-stick model of MacArthur (1957) were computed as indications of tardigrade species diversity and group organization, respectively.

## RESULTS

Only one species (*Milnesium tardigradum* Doyere), repre-

sented by four individuals, was collected from the DDT sprayed area, whereas three species (*Milnesium tardigradum* Doyere, *Macrobiotus islandicus* Richters, and *Macrobiotus areolatus* Murray), represented by a total of 97 individuals, were found to inhabit samples collected from the nontreated habitat. The three species in the nontreated area were represented as follows: *M. tardigradum*, 53 individuals; *M. islandicus*, 35 individuals; and *M. areolatus*, 9 individuals. A highly significant difference ( $P < .01$ ) was found between the densities of the two areas when tested with a Mann-Whitney U test.

Margalef's diversity index ( $D = S - 1 / \ln N$ ), where  $S$  equals the total number of species and  $N$  equals the total number of individuals, was computed and found to be 0.00 and 0.44 for the treated and nontreated samples, respectively (Margalef, 1957).

Tardigrade species organization within the nontreated samples did not differ significantly ( $X^2 = 3.34$ , 2 df) from that as predicted by MacArthur's broken-stick model for contiguous, non-overlapping niches (MacArthur, 1957). This model generates the expected abundance of species according to their rank in rarity and is based upon the observed total number of species and individuals. It may be summarized as follows:

$r^{\text{th}}$  rarest species =

$$\frac{S}{N} \sum_{i=1}^r \frac{1}{S-i+1}$$

where  $S$  equals total number of species,  $N$  equals total number of individuals, and  $i$  equals rank in rarity of species.

## DISCUSSION

The present study was conducted in an attempt to measure and evaluate the effects of a pesticide stress (DDT) on the density, diversity and species organization of tardigrades located within a lichen-bearing tree bark ecosystem. One might, perhaps, view the trunk of an American elm tree as a multichanneled system composed of numerous potential water pathways; these channels being water fed during rainy periods by an overhead deliquescent type of branching tributaries. Further, lichens adhering to this trunk system often tend to serve as "sponges" for the rapid uptake of water due to a system of air-filled spaces which appear between the

<sup>1</sup> Contribution No. 42 from the Dept. of Biology, Drake University.

<sup>2</sup> Department of Zoology, Miami University, Oxford, Ohio.

<sup>3</sup> 202 Kellogg Place, Wheaton, Illinois.

cortical hyphae during dry conditions (Stocker, 1927). Also, lichens have been shown to adsorb and retain numerous organic compounds (Smith, 1960). Thus, an intensive analysis of this lichen-bearing tree bark ecosystem, with its associated microfauna, should provide an excellent site in which to monitor the effects of a pesticide following its application to a forest community. Such was the rationale behind the present study.

Tardigrade density was significantly reduced in samples collected from the sprayed area. Of greater importance, perhaps, was an indication of a reduction in species diversity in the treated area. Once again, as has been found by other investigators (Elton, 1958; Menhinick, 1962; Barrett, 1968; and others), a pesticide stress tends to reduce the diversity of a system, thereby making it more susceptible to additional external perturbations (Odum, 1969).

Of ecological interest was the organization of tardigrade species within the nontreated habitat. Species organization within this relatively simple system appeared to fit MacArthur's broken-stick model for contiguous, non-overlapping niches (MacArthur, 1957). Should further studies confirm this species arrangement, then such a criterion would serve as a useful tool in evaluating the effects of an environmental stress (e.g., pesticides, fire, radiation, etc.) on the structure of an intact, functional forest ecosystem.

The functional parameters of a lichen-bearing tree bark ecosystem (i.e., energy flow or mineral recycling) have not been investigated. We suggest that this relatively simple and accessible system should lend itself well to such future investigations.

#### ACKNOWLEDGMENT

The authors wish to express their thanks to Dr. Paul A. Meglitsch, Department of Biology, Drake University, for his constructive criticism of this manuscript.

#### LITERATURE CITED

- BARKER, R. J. 1958. Notes on some ecological effects of DDT sprayed on elms. *J. Wildlife Mgmt.* 22:269-274.
- BARRETT, G. W. 1968. The effects of an acute insecticide stress on a semi-enclosed grassland ecosystem. *Ecology* 49:1019-1035.
- BENTON, A. H. 1951. Effects on wildlife of DDT used for control of Dutch elm disease. *J. Wildlife Mgmt.* 15:20-27.
- BLACBROUGH, H. P. 1952. Reducing wildlife hazards in Dutch elm disease control. *J. Forest.* 50:468-469.
- DAVIS, D. W. 1952. Some effects of DDT on spider mites. *J. Econ. Ent.* 45:1011-1019.
- ELTON, C. S. 1958. *The ecology of invasions by animals and plants.* Oxford Univ. Press, London. 181pp.
- HUNT, L. B. 1960. Songbird breeding populations in DDT-sprayed Dutch elm disease communities. *J. Wildlife Mgmt.* 24:139-146.
- KIMMEL, R. G. & P. A. MEGLITSCH. 1969. Notes on Iowa tardigrades. *Proc. Iowa Acad. Sci.* 76:454-462.
- MACARTHUR, R. H. 1957. On the relative abundance of bird species. *Proc. Nat. Acad. Sci., U.S.* 43:293-295.
- MARGALEF, R. 1957. La teoria de la informacion en ecologia. *Mem. Real Acad. Ciencias by Artes de Barcelona* 32:373-449.
- MENHINICK, E. F. 1962. Comparison of invertebrate populations of soil and litter of mowed grasslands in areas treated and untreated with pesticides. *Ecology* 43:556-561.
- MOORE, N. W. 1967. A synopsis of the pesticide problem. pp. 75-129. In J. B. Cragg (ed.), *Advances in ecological research*, Vol. 4.
- ODUM, E. P. 1969. The strategy of ecosystem development. *Science* 164:262-270.
- ODUM, W. E., G. M. WOODWELL & C. F. WURSTER. 1969. DDT residues absorbed from organic detritus by fiddler crabs. *Science* 164:576-577.
- RIPPER, W. E. 1956. Effects of pesticides on balance of arthropod populations. *Ann. Rev. Entomol.* 1:403-438.
- SMITH, D. C. 1960. Absorption and utilization of some simple organic nitrogen compounds by *Peltigera polydactyla*. *Ann. Botany* 24:172-185.
- STOCKER, O. 1927. Physiologische und ökologische untersuchungen an laubund strauchflechten. *Flora* 121:334-415.