

Proceedings of the Iowa Academy of Science

Volume 49 | Annual Issue

Article 56

1942

Dissociation Pressure of Some Potassium Polyhalides

Jacob Cornog
University of Iowa

Eldon Bauer
University of Iowa

Copyright © Copyright 1942 by the Iowa Academy of Science, Inc.
Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Cornog, Jacob and Bauer, Eldon (1942) "Dissociation Pressure of Some Potassium Polyhalides," *Proceedings of the Iowa Academy of Science*: Vol. 49: No. 1 , Article 56.
Available at: <https://scholarworks.uni.edu/pias/vol49/iss1/56>

This Research is brought to you for free and open access by UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

DISSOCIATION PRESSURES OF SOME POTASSIUM POLYHALIDES

JACOB CORNOG AND ELDON BAUER

Potassium dichloro iodide (KLCL_2 or KCL-ICL , m. p. 195°) and potassium trichloro di-iodide (KCL_2ICL , m. p. 45°) have been prepared and their dissociation pressures measured.

Both of these are new compounds. The potassium dichloro iodide (m. p. 60°) described by Wells and Wheeler (also by Ephraim) has been found to have the formula $\text{KLCL}_2\cdot\text{H}_2\text{O}$.

CHEMISTRY DEPARTMENT,
UNIVERSITY OF IOWA,
IOWA CITY, IOWA

A MEASUREMENT OF THE MAGNITUDE OF THE ELECTROKINETIC CURRENT IN LIQUID FLOW THROUGH A SINGLE CAPILLARY.

W. G. EVERSELE AND W. W. BOARDMAN

The potential difference, E , between the two ends of a capillary, through which a steady flow of liquid was maintained, was measured by means of unpolarizable electrodes connected to a potentiometer circuit, and at the same time shunted through a known resistance, R . The value of R was varied from 1 to 98000 megohms. For each value of R there was a corresponding value of E , and a current, I ($=E/R$), through the resistance. The electrokinetic current, I_e , was obtained by extrapolating the plot of E/R versus R to the limit $R=0$.

The value of the electrokinetic potential, ζ , can be calculated from I_e by means of the Helmholtz-Smoluchowski equation,

$$\zeta = \frac{4 \eta l}{r^2 DP} I_e,$$

without the use of conductivity data.

The specific surface conductivity can also be evaluated from the same data if the specific conductivity of the liquid in bulk is known.

DEPARTMENT OF CHEMISTRY,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA