

Proceedings of the Iowa Academy of Science

Volume 48 | Annual Issue

Article 62

1941

Vapor-Pressure Lowering of Sulfur Dioxide Solutions of Potassium Thiocyanate from 15 to 25 C. (Abstract)

W. G. Eversole
State University of Iowa

G. H. Wagner
State University of Iowa

Copyright ©1941 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Eversole, W. G. and Wagner, G. H. (1941) "Vapor-Pressure Lowering of Sulfur Dioxide Solutions of Potassium Thiocyanate from 15 to 25 C. (Abstract)," *Proceedings of the Iowa Academy of Science*, 48(1), 275-275.

Available at: <https://scholarworks.uni.edu/pias/vol48/iss1/62>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE THICKNESS OF A RIGID WATER LAYER ON
QUARTZ FROM MEASUREMENTS OF NEWTON'S
RINGS

(ABSTRACT)

W. G. EVERSOLE AND PAUL H. LAHR

Newton's rings produced by a quartz plate and lens in a slit source of light were observed in a low power traveling microscope. The square of the radius of the rings (r) plotted against the order of the ring (n) from the center yields a straight line whose intercept on the axis where $n = 0$ is shown to be a function of the vertical separation of the lens and plate but not a function of the index of refraction of the medium between them nor the angle of incidence of the light. The ring radii were measured with air and with water between the lens and plate. The graphs for the two cases have different intercepts which is interpreted as indicating a rigid water multilayer between the quartz surfaces of the order of 200 $\text{A}\text{\AA}$ in thickness.

DIVISION OF PHYSICAL CHEMISTRY,
STATE UNIVERSITY OF IOWA.

VAPOR-PRESSURE LOWERING OF SULFUR DIOXIDE
SOLUTIONS OF POTASSIUM THIOCYANATE
FROM 15° TO 25° C.

(ABSTRACT)

W. G. EVERSOLE AND G. H. WAGNER

The vapor-pressure lowering of sulfur dioxide by potassium thiocyanate at 15°, 20° and 25° C. was measured by means of a differential mercury manometer connected to an all glass apparatus which employed no stopcocks. Readings were made to 0.001 cm. with a cathetometer. All concentrations were corrected for the amount of sulfur dioxide in the vapor phase. The molal lowering was approximately one-half the theoretical value at 0.005 molal and decreased with increasing concentration up to 1 molal, the highest concentration studied.

DIVISION OF PHYSICAL CHEMISTRY,
THE STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.