

Proceedings of the Iowa Academy of Science

Volume 46 | Annual Issue

Article 43

1939

X-Ray Diffraction Studies on Nerve

R. S. Bear
Iowa State College

Copyright ©1939 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Bear, R. S. (1939) "X-Ray Diffraction Studies on Nerve," *Proceedings of the Iowa Academy of Science*, 46(1), 217-217.

Available at: <https://scholarworks.uni.edu/pias/vol46/iss1/43>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE USE OF THE SPECTROSCOPE IN CHEMISTRY

F. H. SPEDDING

A discussion was given of the types of spectroscopes with particular attention to the limitations and applications of certain instruments for absorption and emission spectra studies. The difficulties of manipulation in various regions of the spectra were itemized with a discussion of present technics for such regions.

DEPARTMENT OF CHEMISTRY,
IOWA STATE COLLEGE,
AMES, IOWA.

X-RAY DIFFRACTION STUDIES ON NERVE

R. S. BEAR

A summary of the present knowledge of the composition and molecular structure of nerve tissue with particular reference to the application of X-ray technic to the problem.

DEPARTMENT OF CHEMISTRY,
IOWA STATE COLLEGE,
AMES, IOWA.

RING COMPOUNDS FROM CHLOROAMINES AND AMIDES

GEORGE H. COLEMAN

Substituted pyrrolidines have been prepared in good yields by heating N-chlorine derivatives of secondary amines with a mixture of sulfuric acid and water. Amines such as methyl-n-butylamine, ethyl-n-butylamine, methyl-n-amylamine, and di-n-butylamine have been used. The optimal conditions for ring closure are not the same for all amines. Varying amounts of secondary amines are usually recovered. In this reaction an intermediate compound