

Proceedings of the Iowa Academy of Science

Volume 46 | Annual Issue

Article 92

1939

Relative Intensities of Fluorescence at Low Temperatures

J. E. Dinger
Iowa State College

Let us know how access to this document benefits you

Copyright ©1939 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Dinger, J. E. (1939) "Relative Intensities of Fluorescence at Low Temperatures," *Proceedings of the Iowa Academy of Science*, 46(1), 272-272.

Available at: <https://scholarworks.uni.edu/pias/vol46/iss1/92>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

constants. The frequencies of vibration of higher harmonics of finite plates, divided by the order of the harmonic, should approach very close to the frequency of the theoretical infinite plate.

Plates of quartz were accurately oriented and cut from well-developed, flaw-free quartz crystals and the frequencies of higher-harmonic piezo-electric vibrations were accurately determined. With this information, the six adiabatic elastic constants of quartz were evaluated.

DEPARTMENT OF PHYSICS,
IOWA STATE COLLEGE,
AMES, IOWA.

ELASTICITY OF LEAD CRYSTALS

A. F. DEMING

Additional measurements have been made on the Young's Modulus of lead single crystals. The value of the modulus depends on the orientation of the length of the specimen relative to the crystallographic axis. This relationship will be shown.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

RELATIVE INTENSITIES OF FLUORESCENCE AT LOW TEMPERATURES

J. E. DINGER

The change in intensity of fluorescence of ZnSiO_3 , CaWO_4 , CdB_2O_5 , was measured at various temperatures between room temperature and the temperature of liquid oxygen. The fluorescence was excited by means of the 2537 Å line of a mercury arc. Microphotometric measurements were made on photographs of the fluorescent spectra. The results indicate a decrease in intensity with decrease in temperature of the CdB_2O_5 and the fluorescence of ZnSiO_3 passes through a maximum of intensity between room temperature and liquid oxygen temperature.

DEPARTMENT OF PHYSICS,
IOWA STATE COLLEGE,
AMES, IOWA.