

# Proceedings of the Iowa Academy of Science

---

Volume 44 | Annual Issue

Article 26

---

1937

## Formation of Heterocyclic Compounds From Haloamines

George H. Coleman  
*State University of Iowa*

Gilbert E. Goheen  
*State University of Iowa*

Copyright ©1937 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Coleman, George H. and Goheen, Gilbert E. (1937) "Formation of Heterocyclic Compounds From Haloamines," *Proceedings of the Iowa Academy of Science*, 44(1), 123-123.

Available at: <https://scholarworks.uni.edu/pias/vol44/iss1/26>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## FORMATION OF HETEROCYCLIC COMPOUNDS FROM HALOAMINES

GEORGE H. COLEMAN AND GILBERT E. GOHEEN

N-chloro-N-methyl-n-butylamine has been prepared and found to be a relatively stable substance. Ring closure was effected by heating this compound with concentrated sulfuric acid, N-methylpyrrolidine being formed in 45 per cent yields.

The corresponding bromoamine was prepared by the method of Loffler<sup>1</sup> and ring closure effected by heating with concentrated sulfuric acid. The yield of N-methylpyrrolidine was, however, considerably less than with the chloroamine.

Several reagents other than sulfuric acid were tried in an attempt to bring about the same ring closure. However, positive results were obtained only with 85 per cent phosphoric acid.

DEPARTMENT OF CHEMISTRY,  
LABORATORY OF ORGANIC CHEMISTRY,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

## HIGH MOLECULAR WEIGHT FATTY ACID DERIVATIVES

GROVER M. FORD AND HENRY GILMAN

A series of derivatives for identifying lauric, myristic, palmitic, stearic and oleic acids have been prepared.

The ten best series of derivatives may be listed in the following order of decreasing importance: (1) N-acylcarbazole derivatives; (2) N-acyl-*p*-toluenesulfonamides; (3) *p*-phenyl-phenacyl esters; (4) N-acylphenothiazine derivatives; (5) N-acyl-2-nitro-*p*-toluidine derivatives; (6) N-acylsaccharin derivatives; (7) 2, 4-dinitrophenylhydrazides; (8) *p*-nitro-anilides; (9) phenylmercuric salts; (10) *p*-xenylamides.

The N-acylcarbazole derivatives show a difference in melting points of adjacent members of from 3-5°, and a depression of

<sup>1</sup> Loffler, Ber., 42: 3427 (1909).