

# Proceedings of the Iowa Academy of Science

---

Volume 42 | Annual Issue

Article 41

---

1935

## The Reaction of Potassium Amide in Liquid Ammonia with Diarylbromethenes

George H. Coleman  
*State University of Iowa*

William H. Host  
*State University of Iowa*

Copyright ©1935 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Coleman, George H. and Host, William H. (1935) "The Reaction of Potassium Amide in Liquid Ammonia with Diarylbromethenes," *Proceedings of the Iowa Academy of Science*, 42(1), 119-119.

Available at: <https://scholarworks.uni.edu/pias/vol42/iss1/41>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## THE REACTION OF POTASSIUM AMIDE IN LIQUID AMMONIA WITH DIARYLBROMOETHENES

GEORGE H. COLEMAN AND WILLIAM H. HOST

The behavior of certain diarylchloroethenes with potassium amide in liquid ammonia has been reported previously.<sup>1</sup> A rapid reaction occurs and the ethenes are changed with the loss of hydrochloric acid to tolanes. Thus 1, 1-di-p-tolyl-2-chloroethene forms p. p'-dimethyltolane. A series of such para substituted compounds was studied and as anticipated only symmetrical tolanes were obtained.

In the present work the following diarylbromoethenes with substituents in the ortho and in the meta positions were used:

1,1-di-o-tolyl-2-bromoethene;  
1,1-di-m-tolyl-2-bromoethene;  
1,1-di-o-methoxyphenyl-2-bromoethene  
1,1-di-m-methoxyphenyl-2-bromoethene

In each case the corresponding symmetrical tolane was obtained. Therefore whatever the exact mechanism of this rearrangement may be a given substituent in the aryl radical of the ethene is found in the same relative position in the tolane.

DEPARTMENT OF CHEMISTRY,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

## HYDROXYFURANS AND THEIR BIOLOGICAL SIGNIFICANCE

W. F. HOEHN, C. H. WERKMAN AND H. GILMAN

Hydroxyfurans, particularly the mono- and di- $\beta$ -substituted types (prepared from furans and by ring closure procedures) have been examined biologically, particularly for their growth **stimulation** effects on bacteria, yeasts and various higher plants. A discussion is included of the synthetic procedures and the proof of structure of the variously substituted hydroxyfurans.

<sup>1</sup> Coleman and Maxwell, J. Am. Chem. Soc., 56, 132 (1934).

DEPARTMENT OF CHEMISTRY,  
IOWA STATE COLLEGE,  
AMES, IOWA.