

Proceedings of the Iowa Academy of Science

Volume 39 | Annual Issue

Article 42

1932

The Preparation and Properties of Furyl Phenyl Ketimine

J. B. Culbertson
Cornell College

Ben Davis
Cornell College

Copyright © Copyright 1932 by the Iowa Academy of Science, Inc.
Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Culbertson, J. B. and Davis, Ben (1932) "The Preparation and Properties of Furyl Phenyl Ketimine," *Proceedings of the Iowa Academy of Science*, 39(1), 177-177.
Available at: <https://scholarworks.uni.edu/pias/vol39/iss1/42>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE STRENGTHS OF PHENOLIC KETIMINES AND THEIR METHYL ETHERS AS BASES

J. B. CULBERTSON, PAUL BIEBER AND ADOLPH ZAVODSKY

The ionization constants of the monohydroxy-diphenyl ketimines and their methyl ethers are calculated from measurements of the hydrogen ion concentration of aqueous solutions of their hydrochlorides through the use of the quinhydrone electrode. This data has been collected as a part of the information expected to throw light on the varied stability toward hydrolysis shown by different ketimines.

DEPARTMENT OF CHEMISTRY,
CORNELL COLLEGE,
MT. VERNON, IOWA.

THE PREPARATION AND PROPERTIES OF FURYL PHENYL KETIMINE

J. B. CULBERTSON AND BEN DAVIS

This ketimine has been prepared through the condensation of furonitrile with magnesium phenyl bromide, followed by treatment with ice and ammonium chloride at about -15 degrees C. The hydrochloride, a white solid, is rather quickly hydrolyzed to the corresponding ketone, benzoyl furane. The velocity of this hydrolysis and the basic strength of the free ketimine base are determined.

DEPARTMENT OF CHEMISTRY,
CORNELL COLLEGE,
MT. VERNON, IOWA.