

1932

Motion of K⁺ Ions in Mercury Vapor

Carl Frische
State University of Iowa

Let us know how access to this document benefits you

Copyright ©1932 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Frische, Carl (1932) "Motion of K⁺ Ions in Mercury Vapor," *Proceedings of the Iowa Academy of Science*, 39(1), 220-220.

Available at: <https://scholarworks.uni.edu/pias/vol39/iss1/62>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

ABSORPTION SPECTRA CHARACTERISTIC OF VITAMIN A IN ANIMAL AND VEGETABLE OILS

J. B. PHILIPSON AND JAY W. WOODROW

A continuation of our previous work on the ultra-violet absorption spectra of substances showing vitamin A activity, tends to further substantiate the evidence that the absorption band in the regions 325-330 m. μ ., which Morton and Heilbron have shown to be characteristic of vitamin A in animal fats, also characterizes vegetable compounds containing the growth promoting factor.

Using the same photoelectric spectrophotometer with which we showed this selective absorption in juices and oils extracted from carrots and from spinach, we have found that fresh orange juice, shown by Osborne and Mandel to contain the growth promoting factor, also shows a distinct absorption band at about 328 m. μ .

These results form the basis for our conclusions that, since they show the same selective absorption, the growth promoting factor in vegetables and the vitamin A in animal oils, if not identically the same compound, at least have very closely related chemical structure.

DEPARTMENT OF PHYSICS,
IOWA STATE COLLEGE,
AMES, IOWA.

MOTION OF K⁺ IONS IN MERCURY VAPOR

CARL FRISCHE

Ionization by positive ion impact was studied in A, Ne, N₂, CO, He, at voltages up to 4000 volts and the results seem to check with those of Sutton (Phys. Rev. 33, 363, 1929) and Beeck (Ann. d. Phys. 6, 8, 1001, 1930) wherever a check was possible.

When the work was extended to an element heavier than K⁺ i.e., Hg. the results are more complicated. The evidence seems to indicate; (a) reflection of K⁺ ions at all voltages, (b) an increase in mean free path with increase in voltage, (c) no ionization below 250 volts, but above 250-350 volts ionization increases with increase in voltage.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.