

# Proceedings of the Iowa Academy of Science

---

Volume 37 | Annual Issue

Article 40

---

1930

## Catalytic Hydrogenation of Furfural

Ralph L. Van Peursesem

*Let us know how access to this document benefits you*

Copyright ©1930 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Van Peursesem, Ralph L. (1930) "Catalytic Hydrogenation of Furfural," *Proceedings of the Iowa Academy of Science*, 37(1), 225-225.

Available at: <https://scholarworks.uni.edu/pias/vol37/iss1/40>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## CATALYTIC HYDROGENATION OF FURFURAL

RALPH L. VAN PEURSEM

A study was made of the possibility of producing the reduction products of furfural by passing its vapor, with an excess of hydrogen, over finely divided catalyst.

In work which had previously been done,<sup>1</sup> the chief product obtained was furfuryl alcohol. With this in mind, a study was made of the identification of this compound. Several of its esters were prepared and identified. Among these was the mono-furfuryl ester of phthalic acid.

In the reduction work, the catalyst first used was freshly reduced nickel. Runs were made at various temperatures, keeping the other conditions constant. The maximum yield was obtained when the catalyst was kept at a temperature of 200°C.

Another similar set of runs was made using finely divided copper as catalyst. The maximum reduction took place at 215°C. The yield of furfuryl alcohol was larger in the case of nickel. The highest yield obtained was 5% of furfuryl alcohol.

---

SOME PHYSICAL PROPERTIES OF CONCENTRATED  
AQUEOUS SALT SOLUTIONS

J. N. PEARCE AND MILDRED A. HOOPER

The densities and refractive indices were determined at 25° for solutions of the halides of some of the alkali metals, over a concentration range from 0.1 molal to almost saturation. The partial molal volumes were calculated by differentiating the equations of the volume-molality curves. From the refractive index the specific refractivity of the solution and of the solute were calculated. The specific refractivity of the solute was found to be constant throughout the concentrations studied. Using an equation derived by Hückel, involving the refractive index, the dielectric constants of the solutions were calculated.

---

<sup>1</sup> Kaufmann and Adams, *J. Am. Chem. Soc.*, 45, 3029-44 (1923).