

# Proceedings of the Iowa Academy of Science

---

Volume 37 | Annual Issue

Article 81

---

1930

## The Disruption of Potentials within a Raw Potato under Constant Applied Voltage

E. C. McCracken  
*Iowa State College*

Copyright ©1930 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

McCracken, E. C. (1930) "The Disruption of Potentials within a Raw Potato under Constant Applied Voltage," *Proceedings of the Iowa Academy of Science*, 37(1), 313-313.

Available at: <https://scholarworks.uni.edu/pias/vol37/iss1/81>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

for Iron and Cobalt films (Phys. Rev. 30, 681 (1927); 35 292 (1930)). Films about 130  $\mu$  thick attain a magnetization of about 380 c. g. s. units in a field of 200 gauss, a value about equal to that for bulk nickel. As in Iron and Cobalt the coercive force is high, but the remanence is somewhat less than was found for iron and cobalt.

STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

### THE DISTRIBUTION OF POTENTIALS WITHIN A RAW POTATO UNDER CONSTANT APPLIED VOLTAGE

E. C. McCRACKEN

Uniformly shaped sections of raw potatoes were placed between parallel plane electrodes and subjected to a constant potential. The potential drops existing between one electrode and various points in the potatoes were measured. The equi-potential surfaces were found to be planes parallel to the surface of the terminal electrodes. The potentials of these planes were found to vary with time and previous passage of current. The relative size of the probing electrodes used influenced the shape of the potential distribution curves.

IOWA STATE COLLEGE,  
AMES, IOWA.

---

### FACTORS INFLUENCING THE ELECTRICAL RESIST- ANCE OF THE POTATO

E. C. McCRACKEN

Extensive experimentation upon the electrical resistance of raw potatoes has given the following results: (1) the resistance of a uniformly shaped section varies inversely as its cross-sectional area; (2) the resistance is independent of the applied voltage; (3) the resistance of sections taken from a relatively homogeneous part of the tuber varies directly with the length; (4) the cortical layer lengthwise of the potato has the least resistance, the resistances of internal medullary area, the external medullary area, and the outer skin following in the order named. No investigation