

Proceedings of the Iowa Academy of Science

Volume 38 | Annual Issue

Article 113

1931

Experiments with Dry Dust Cultures for the Inoculation of Legume Seeds

R. H. Walker
Iowa State College

G. B. Killinger
Iowa State College

Let us know how access to this document benefits you

Copyright ©1931 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Walker, R. H. and Killinger, G. B. (1931) "Experiments with Dry Dust Cultures for the Inoculation of Legume Seeds," *Proceedings of the Iowa Academy of Science*, 38(1), 333-333.

Available at: <https://scholarworks.uni.edu/pias/vol38/iss1/113>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

EXPERIMENTS WITH DRY DUST CULTURES FOR THE INOCULATION OF LEGUME SEEDS

R. H. WALKER AND G. B. KILLINGER

Within the past few years a number of commercial concerns have produced a dry dust type of culture for the inoculation of legume seeds. Field experiments have been conducted to test the value of these cultures for the inoculation of soybeans. The results showed that the dry dust cultures did not give satisfactory inoculation.

Greenhouse experiments have also shown that the inoculation of legume seeds sometime in advance of the planting date, as is recommended by the producers of the dry dust manufacturers, is undesirable.

IOWA STATE COLLEGE,
AMES, IOWA.

COMPARATIVE GROWTH RATES OF DIFFERENT STRAINS OF *RHIZOBIUM MELILOTI*

R. H. WALKER AND DEAN A. ANDERSON

The growth rates of two strains of *Rhizobium meliloti* were measured when the organisms were grown in a nitrogen-free medium and in a medium containing nitrate nitrogen. The strains used differed widely in their ability to aid the host plant in symbiotic nitrogen fixation. They also differed somewhat in cultural characteristics.

The strain which was more efficient in nitrogen fixation in symbiosis with the host plant grew faster in the medium without nitrogen, than it did in the medium containing nitrate nitrogen. On the other hand, the strain of less efficient organisms grew faster in the medium containing nitrate nitrogen than in the nitrogen-free medium.

The results seem to indicate that the organisms which are less efficient in fixing nitrogen grow faster in the presence of com-