

Proceedings of the Iowa Academy of Science

Volume 30 | Annual Issue

Article 18

1923

The Distribution of Intensity in the Broadened Balmer Lines of Hydrogen

E. O. Hulburt

Copyright © Copyright 1923 by the Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Hulburt, E. O. (1923) "The Distribution of Intensity in the Broadened Balmer Lines of Hydrogen," *Proceedings of the Iowa Academy of Science*, 30(1), 77-77.

Available at: <https://scholarworks.uni.edu/pias/vol30/iss1/18>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE DISTRIBUTION OF INTENSITY IN THE BROADENED BALMER LINES OF HYDROGEN

E. O. HULBURT

(ABSTRACT)

The *intensity distribution across the broadened Balmer lines of hydrogen* H_{β} , H_{γ} and H_{δ} excited by condensed discharges for a pressure range from 48 to 250 mm. of mercury was determined by photographing the spectra through a neutral wedge. The broadening was symmetrical and the general form of the intensity curve was convex to the axis of wave-lengths; the curves for each line gave evidences of structure characteristic of the line.

A *theoretical analysis* yielded an expression which agreed with the general form of the experimental intensity curve, namely, the intensity I_{λ} at wave-length λ in the broadened line whose centre was at λ_0 is

$$I_{\lambda} = \frac{a}{\lambda_0 - \lambda} e^{-\frac{b}{\lambda_0 - \lambda}}$$

where a and b are quantities which vary with the line and with the pressure; b is small. In the derivation of this formula the luminous atoms of the gas were considered to be subjected to electric fields caused by the charged particles of the gas, the electric fields being allotted according to a probability law. The electric field resolved the line into the Stark components, the displacement of the components from the parent line being given by a general relation due to Sommerfeld.

As a result of the theory the *action of the condensed discharge* in producing widening of the line is attributed to the high value of the current intensity through the gas in the early portion of the discharges, which may perhaps rise to hundreds of amperes. The slight structure in the broadened line may be explained in terms of the observed intensities and positions of the components of the electrically resolved line.