

THE THIRD HOMOTOPY GROUP AS A π_1 -MODULE

HANS-JOACHIM BAUES AND BEATRICE BLEILE

ABSTRACT. It is well-known how to compute the structure of the second homotopy group of a space, X , as a module over the fundamental group, $\pi_1 X$, using the homology of the universal cover and the Hurewicz isomorphism. We describe a new method to compute the third homotopy group, $\pi_3 X$, as a module over $\pi_1 X$. Moreover, we determine $\pi_3 X$ as an extension of $\pi_1 X$ -modules derived from Whitehead's Certain Exact Sequence. Our method is based on the theory of quadratic modules. Explicit computations are carried out for pseudo-projective 3-spaces $X = S^1 \cup e^2 \cup e^3$ consisting of exactly one cell in each dimension ≤ 3 .

1. INTRODUCTION

Given a connected 3-dimensional CW-complex, X , with universal cover, \widehat{X} , Whitehead's Certain Exact Sequence [W2] yields the short exact sequence

$$(1.1) \quad \Gamma\pi_2 X \twoheadrightarrow \pi_3 X \twoheadrightarrow H_3 \widehat{X}$$

of π_1 -modules, where $\pi_1 = \pi_1(X)$. As a group, the homology $H_3 \widehat{X}$ is a subgroup of the free abelian group of cellular 3-chains of \widehat{X} , and thus itself free abelian. Hence the sequence splits as a sequence of abelian groups. This raises the question whether (1.1) splits as a sequence of π_1 -modules – there are no examples known in the literature.

It is well-known how to compute $\pi_2(X) \cong H_2 \widehat{X}$ as a π_1 -module, using the Hurewicz isomorphism, and how to compute $H_3 \widehat{X}$ using the cellular chains of the universal cover. In this paper we compute $\pi_3(X)$ as π_1 -module and (1.1) as an extension over π_1 . We answer the question above by providing an infinite family of examples where (1.1) does not split over π_1 , as well as an infinite family of examples where it does split over π_1 . As a first surprising example we obtain

Theorem 1.1. *There is a connected 3-dimensional CW-complex X with fundamental group $\pi_1 = \pi_1 X = \mathbb{Z}/2\mathbb{Z}$, such that π_1 acts trivially on both $\Gamma\pi_2 X$ and $H_3 \widehat{X}$, but non-trivially on $\pi_3 X$. Hence*

$$\Gamma\pi_2 X \twoheadrightarrow \pi_3 X \twoheadrightarrow H_3 \widehat{X}$$

does not split as a sequence of π_1 -modules.

Below we describe examples for all finite cyclic fundamental groups, π_1 , of even order, where (1.1) does not split over π_1 . The examples we consider are CW-complexes,

$$X = S^1 \cup e^2 \cup e^3,$$

with precisely one cell, e^i , in every dimension $i = 0, 1, 2, 3$. In general, we obtain such a CW-complex, X , by first attaching the 2-cell e_2 to S^1 via $f \in \pi_1 S^1 = \mathbb{Z}$. We assume $f > 0$. This yields the 2-skeleton of X , $X^2 = P_f$, which is a pseudo-projective plane, see [O]. Then $\pi_1 = \pi_1 X = \pi_1 P_f = \mathbb{Z}/f\mathbb{Z}$ is a cyclic group of order f . We write $R = \mathbb{Z}[\pi_1]$ for the integral group ring of π_1 and K for the kernel of the augmentation $\varepsilon : R \rightarrow \mathbb{Z}$. Then the pseudo-projective 3-space, $X = P_{f,x}$, is determined by the pair, (f, x) , of attaching maps, where $x \in \pi_2 P_f = K$ is the attaching map of the 3-cell e_3 . In this case

$$\pi_2(X) = H_2(\widehat{X}) = K/xR,$$

and

$$H_3 \widehat{X} = \ker(d_x : R \rightarrow R, x \mapsto xy),$$

where xy is the product of $x, y \in R$.

A splitting function u for the exact sequence (1.1) is a function between sets, $u : H_3\widehat{X} \rightarrow \pi_3 X$, such that $u(0) = 0$ and the composite of u and the projection $\pi_3 X \rightarrow H_3\widehat{X}$ is the identity. Such a splitting function determines maps

$$A = A_u : H_3\widehat{X} \times H_3\widehat{X} \rightarrow \Gamma(\pi_2 X) \quad \text{and} \quad B = B_u : H_3\widehat{X} \rightarrow \Gamma(\pi_2 X),$$

by the cross-effect formulæ

$$A(y, z) = u(y + z) - (u(y) + u(z)) \quad \text{and} \quad B(y) = (u(y))^1 - u(y^1).$$

Here B is determined by the action of the generator 1 in the cyclic group π_1 , denoted by $y \mapsto y^1$.

Remark 1.2. The functions A and B determine $\pi_3 X$ as a π_1 -module. In fact, the bijection $H_3\widehat{X} \times \Gamma(\pi_2 X) = \pi_3(P_{f,x})$, which assigns to (y, v) the element $u(y) + v$ is an isomorphism of π_1 -modules, where the left hand side is an abelian group by

$$(y, v) + (z, w) = (y + z, v + w + A(y, z))$$

and a π_1 -module by

$$(y, v)^1 = (y^1, v^1 + B(y)).$$

The cross-effect of B satisfies

$$B(y + z) - (B(y) + B(z)) = (A(y, z))^1 - A(y^1, z^1),$$

such that B is a homomorphism of abelian groups if $A = 0$.

In this paper we describe a method to determine a splitting function $u = u_x$, which, a priori, is not a homomorphism of abelian groups. We investigate the corresponding functions A and B and compute them for a family of examples.

Theorem 1.3. *Let $X = P_{f,x}$ be a pseudo-projective 3-space with $x = \tilde{x}([\overline{1}] - [\overline{0}]) \in K$, $\tilde{x} \in \mathbb{Z}$, $\tilde{x} \neq 0$ and $f > 1$. Let $N = \sum_{i=0}^{f-1} [\tilde{v}]$ be the norm element in R . Then*

$$H_3(\widehat{P}_{f,x}) = \{\tilde{y}N \mid \tilde{y} \in \mathbb{Z}\} \cong \mathbb{Z}$$

is a π_1 -module with trivial action of π_1 , and

$$\pi_2(P_{f,x}) = (\mathbb{Z}/\tilde{x}\mathbb{Z}) \otimes_{\mathbb{Z}} K,$$

with the action of π_1 induced by the π_1 -module K . There is a splitting function $u = u_x$ such that, for $y = \tilde{y}N$ and $z \in H_3(\widehat{P}_{f,x})$, the functions A and B are given by

$$\begin{aligned} A(y, z) &= 0 \\ B(y) &= -\tilde{x}\tilde{y}\gamma q([\overline{1}] - [\overline{0}]), \end{aligned}$$

where $\gamma : \pi_2(P_{f,x}) \rightarrow \Gamma(\pi_2(P_{f,x}))$ is the universal quadratic map for the Whitehead functor Γ and $q : K \rightarrow \pi_2(P_{f,x})$, $k \mapsto 1 \otimes k$. As in 1.2, the pair A, B computes $\pi_3 X$ as a π_1 -module.

As $H_3(\widehat{X})$ is free abelian, the exact sequence (1.1) always allows a splitting function which is a homomorphism of abelian groups. This leads, for $X = P_{f,x}$, to the injective function

$$\tau : \text{Ext}_{\pi_1}(H_3(\widehat{X}), \Gamma(\pi_2 X)) \rightarrow \text{coker}(\beta),$$

with

$$\beta : \text{Hom}_{\mathbb{Z}}(H_3(\widehat{X}), \Gamma(\pi_2 X)) \rightarrow \text{Hom}_{\mathbb{Z}}(H_3(\widehat{X}), \Gamma(\pi_2 X)), t \mapsto \beta_t,$$

given by

$$\beta_t(\ell) = -t(\ell^1) + (t(\ell))^1.$$

The function τ maps the equivalence class of an extension to the element in $\text{coker}\beta$ represented by $B = B_u$, where u is a \mathbb{Z} -homomorphic splitting function for the extension. Hence the equivalence class, $\{\pi_3 X\}$, of the extension $\pi_3 X$ in (1.1) is determined by the image $\tau\{\pi_3 X\} \in \text{coker}(\beta)$. For the family of examples in 1.3 we show

Theorem 1.4. *Let $X = P_{f,x}$ be a pseudo-projective 3-space with $x = \tilde{x}([\overline{1}] - [\overline{0}])$, $\tilde{x} \in \mathbb{Z}$, $\tilde{x} \neq 0$ and $f > 1$. Then $\beta : \Gamma((\mathbb{Z}/\tilde{x}\mathbb{Z}) \otimes_{\mathbb{Z}} K) \rightarrow \Gamma((\mathbb{Z}/\tilde{x}\mathbb{Z}) \otimes_{\mathbb{Z}} K)$ maps ℓ to $-\ell + \ell^1$ and $\tau\{\pi_3 X\} \in \text{coker}(\beta)$ is represented by $\tilde{x}\gamma q([\overline{1}] - [\overline{0}]) \in \Gamma(\pi_2)$. Hence $\tau\{\pi_3 X\} = 0$ if \tilde{x} is odd, so that, in this case, $\pi_3 X$ in (1.1) is a split extension over π_1 . If both \tilde{x} and f are even, then $\tau\{\pi_3 X\}$ is a non-trivial element of order 2, and the extension $\pi_3 X$ in (1.1) does not split over π_1 . Moreover, $\tau\{\pi_3 X\}$ is represented by B in 1.3. If \tilde{x} is even and f is odd, then $\tau\{\pi_3 X\}$ is trivial and the extension $\pi_3 X$ in (1.1) does split over π_1 .*

This result is a corollary of 1.3, the computations are contained at the end of Section 8.

Given a pseudo-projective 3-space, $P_{f,x}$, and an element $z \in \pi_3(P_{f,x})$, we obtain a pseudo-projective 4-space, $X = P_{f,x,z} = S^1 \cup e^2 \cup e^3 \cup e^4$, where z is the attaching map of the 4-cell e^4 . For $n \geq 2$, the attaching map z of an $(n+1)$ -cell in a CW-complex, X , is *homologically non-trivial* if the image of z under the Hurewicz homomorphism is non-trivial in $H_n \widehat{X}^n$.

Theorem 1.5. *Let $X = S^1 \cup e^2 \cup e^3 \cup e^4$ be a pseudo-projective 4-space with $\pi_1 X = \mathbb{Z}/2\mathbb{Z}$ and homologically non-trivial attaching maps of cells in dimension 3 and 4. Then the action of $\pi_1 X$ on $\pi_3 X$ is trivial.*

Theorem 1.5 is a corollary to Theorem 9.1.

2. CROSSED MODULES

We recall the notions of pre-crossed module, Peiffer commutator, crossed module and $\text{nil}(2)$ -module, which are ingredients of algebraic models of 2- and 3-dimensional CW-complexes used in the proofs of our results, see [B] and [BHS]. In particular, Theorem 2.2 provides an exact sequence in the algebraic context of a $\text{nil}(2)$ -module equivalent to Whitehead's Certain Exact Sequence (1.1).

A *pre-crossed module* is a homomorphism of groups, $\partial : M \rightarrow N$, together with an action of N on M , such that, for $x \in M$ and $\alpha \in N$,

$$\partial(x^\alpha) = -\alpha + \partial x + \alpha.$$

Here the action is given by $(\alpha, x) \mapsto x^\alpha$ and we use additive notation for group operations even where the group fails to be abelian. The *Peiffer commutator* of $x, y \in M$ in such a pre-crossed module is given by

$$\langle x, y \rangle = -x - y + x + y^{\partial x}.$$

The subgroup of M generated by all iterated Peiffer commutators $\langle x_1, \dots, x_n \rangle$ of length n is denoted by $P_n(\partial)$ and a *nil(n)-module* is a pre-crossed module $\partial : M \rightarrow N$ with $P_{n+1}(\partial) = 0$. A *crossed module* is a $\text{nil}(1)$ -module, that is, a pre-crossed module in which all Peiffer commutators vanish. We also consider $\text{nil}(2)$ -modules, that is, pre-crossed modules for which $P_3(\partial) = 0$.

A morphism or map $(m, n) : \partial \rightarrow \partial'$ in the category of pre-crossed modules is given by a commutative diagram

$$\begin{array}{ccc} M & \xrightarrow{m} & M' \\ \partial \downarrow & & \downarrow \partial' \\ N & \xrightarrow{n} & N' \end{array}$$

in the category of groups, where m is n -equivariant, that is, $m(x^\alpha) = m(x)^{n(\alpha)}$, for $x \in M$ and $\alpha \in N$. The categories of crossed modules and $\text{nil}(2)$ -modules are full subcategories of the category of pre-crossed modules.

Note that $P_{n+1}(\partial) \subseteq \ker \partial$ for any pre-crossed module, $\partial : M \rightarrow N$. Thus we obtain the *associated nil(n)-module* $r_n(\partial) : M/P_{n+1}(\partial) \rightarrow N$, where the action on the quotient is determined by demanding that the quotient map $q : M \rightarrow M/P_{n+1}(\partial)$ be equivariant. For $n = 1$ we write $\partial^{cr} = r_1(\partial) : M^{cr} = M/P_2(\partial) \rightarrow N$ for the *crossed module associated to ∂* .

Given a set, Z , let $\langle Z \rangle$ denote the free group generated by Z . Now take a group, N , and a group homomorphism, $f : F = \langle Z \rangle \rightarrow N$. Then the *free N -group* generated by Z is the free

group, $\langle Z \times N \rangle$, generated by elements denoted by $x^\alpha = ((x, \alpha))$ with $x \in Z$ and $\alpha \in N$. These are elements in the product $Z \times N$ of sets. The action is determined by

$$(2.1) \quad ((x, \alpha))^\beta = ((x, \alpha + \beta)).$$

Define the group homomorphism $\partial_f : \langle Z \times N \rangle \rightarrow N$ by $((x, \alpha)) \mapsto -\alpha + f(x) + \alpha$, for generators $((x, \alpha)) \in \langle Z \times N \rangle$, to obtain the pre-crossed module ∂_f with associated $\text{nil}(n)$ -module $r_n(\partial_f) : \langle Z \times N \rangle / P_{n+1}(\partial_f) \rightarrow N$. Note that $r_n(\partial_f)\iota = f$, where $\iota = p\iota_F$ is the composition of the inclusion $\iota_F : F = \langle Z \rangle \rightarrow \langle Z \times N \rangle$ and the projection $p : \langle Z \times N \rangle \rightarrow M = \langle Z \times N \rangle / P_{n+1}(\partial_f)$ onto the quotient.

Remark 2.1. The $\text{nil}(n)$ -module, $r_n(\partial_f) : M = \langle Z \times N \rangle / P_{n+1}(\partial_f) \rightarrow N$, satisfies the following universal property: For every $\text{nil}(n)$ -module, $\partial' : M' \rightarrow N'$, and every pair of group homomorphisms, $m_F : F = \langle Z \rangle \rightarrow M'$, and $n : N \rightarrow N'$ with $\partial' m_F = n f$, there is a unique group homomorphism, $m : M \rightarrow M'$, such that $m\iota = m_F$, and $(n, m) : r_n(\partial_f) \rightarrow \partial'$ is a map of $\text{nil}(n)$ -modules.

$$\begin{array}{ccc} M & \overset{m}{\dashrightarrow} & M' \\ \downarrow r_n(\partial_f) & \swarrow \iota & \nearrow m_F \\ & F & \\ \downarrow & \searrow f & \downarrow \partial' \\ N & \xrightarrow{n} & N' \end{array}$$

Thus $r_n(\partial_f)$ is called the *free nil(n)-module with basis f*. A free $\text{nil}(n)$ -module is *totally free* if N is a free group.

Given a path connected space Y and a space X obtained from Y by attaching 2-cells, let Z_2 be the set of 2-cells in $X - Y$, and let $f : Z_2 \rightarrow \pi_1(Y)$ be the attaching map. J.H.C. Whitehead [W1] showed that

$$(2.2) \quad \partial : \pi_2(X, Y) \rightarrow \pi_1(Y)$$

is a free crossed module with basis f . Then $\ker \partial = \pi_2(X)$, $\text{coker } \partial = \pi_1(X)$ and ∂ is totally free if Y is a one-point union of 1-spheres. Whitehead also proved that the abelianisation of the group $\pi_2(X, Y)$ is the free R -module $\langle Z_2 \rangle_R$ generated by the set Z_2 , where $R = \mathbb{Z}[\pi_1(X)]$ is the group ring [W1].

Now take a totally free $\text{nil}(2)$ -module $\partial : M \rightarrow N$ with associated crossed module $\partial^{cr} : M^{cr} \rightarrow N$. Let

$$M \overset{q}{\twoheadrightarrow} M^{cr} \overset{h_2}{\twoheadrightarrow} C = (M^{cr})^{ab}$$

be the composition of projections. Put $K = h_2(\ker(\partial^{cr}))$. Further, let Γ be *Whitehead's quadratic functor* and $\tau : \Gamma(K) \rightarrow K \otimes K \subset C \otimes C$ the composition of the injective homomorphism induced by the quadratic map $K \rightarrow K \otimes K, k \mapsto k \otimes k$ and the inclusion. The *Peiffer commutator map*, $w : C \otimes C \rightarrow M$, is given by $w(\{x\} \otimes \{y\}) = \langle x, y \rangle$, for $x, y \in M$ with $\{x\} = h_2(q(x)), \{y\} = h_2(q(y))$. Lemma (IV 1.6) and Theorem (IV 1.8) in [B] imply

Theorem 2.2. *Let $\partial : M \rightarrow N$ be a totally free nil(2)-module. Then the sequence*

$$\Gamma(K) \xrightarrow{\tau} C \otimes C \xrightarrow{w} M \overset{q}{\twoheadrightarrow} M^{cr}$$

is exact and the image of w is central in M .

3. PSEUDO-PROJECTIVE SPACES IN DIMENSIONS 2 AND 3

Real projective n -space $\mathbb{R}P^n$ has a cell structure with precisely one cell in each dimension $\leq n$. More generally, a CW-complex,

$$X = S^1 \cup e^2 \cup \dots \cup e^n,$$

with precisely one cell in each dimension $\leq n$, is called a *pseudo-projective n -space*. For $n = 2$ we obtain *pseudo-projective planes*, see [O]. In this section we fix notation and consider pseudo-projective spaces in dimensions 2 and 3. In particular, we determine the totally free crossed module associated with a pseudo-projective plane and begin to investigate the totally free $\text{nil}(2)$ -module associated with a pseudo-projective 3-space.

The fundamental group of a pseudo-projective plane $P_f = S^1 \cup e^2$, with attaching map $f \in \pi_1(S^1) = \mathbb{Z}$, is the cyclic group $\pi_1 = \pi_1(P_f) = \mathbb{Z}/f\mathbb{Z}$. We obtain $\pi_1 = \mathbb{Z}$ for $f = 0$, $\pi_1 = \{0\}$ for $f = 1$, and the bijection of sets

$$\{0, 1, 2, \dots, f-1\} \rightarrow \pi_1 = \mathbb{Z}/f\mathbb{Z}, \quad k \mapsto \bar{k} = k + f\mathbb{Z},$$

for $1 < f$. Addition in π_1 is given by

$$\bar{k} + \bar{\ell} = \begin{cases} \overline{k + \ell} & \text{for } k + \ell < f; \\ \overline{k + \ell - f} & \text{for } k + \ell \geq f. \end{cases}$$

Denoting the integral group ring of the cyclic group π_1 by $R = \mathbb{Z}[\pi_1]$, an element $x \in R$ is a linear combination

$$x = \sum_{\alpha \in \pi_1} x_\alpha [\alpha] = \sum_{k=0}^{f-1} x_{\bar{k}} [\bar{k}],$$

with $x_\alpha, x_{\bar{k}} \in \mathbb{Z}$. Note that $1_R = [\bar{0}]$ is the neutral element with respect to multiplication in R and, for $x = \sum_{\alpha \in \pi_1} x_\alpha [\alpha], y = \sum_{\beta \in \pi_1} y_\beta [\beta]$,

$$xy = \sum_{\alpha, \beta \in \pi_1} x_\alpha y_\beta [\alpha + \beta] = \sum_{\ell=0}^{f-1} \left(\sum_{k=0}^{\ell} x_{\bar{k}} y_{\overline{\ell-k}} + \sum_{k=\ell+1}^{f-1} x_{\bar{k}} y_{\overline{f+\ell-k}} \right) [\bar{\ell}].$$

The *augmentation* $\varepsilon = \varepsilon_R : R \rightarrow \mathbb{Z}$ maps $\sum_{\alpha \in \pi_1} x_\alpha [\alpha]$ to $\sum_{\alpha \in \pi_1} x_\alpha$. The *augmentation ideal*, K , is the kernel of ε . For a right R -module, C , we write the action of $\alpha \in \pi_1$ on $x \in C$ exponentially as $x^\alpha = x[\alpha]$.

Given a pseudo-projective plane $P_f = S^1 \cup e^2$ with attaching map $f \in \pi_1(S^1) = \mathbb{Z}$, Whitehead's results on the free crossed module (2.2) imply that

$$(3.1) \quad \partial : \pi_2(P_f, S^1) \rightarrow \pi_1(S^1)$$

is a totally free crossed module with one generator, e_i , in dimensions $i = 1, 2$, and basis $\tilde{f} : Z_2 = \{e_2\} \rightarrow \pi_1(S^1)$ given by $\tilde{f}(e_2) = fe_1$. Note that ∂ has cokernel $\pi_1(P_f) = \mathbb{Z}/f\mathbb{Z} = \pi_1$ and kernel $\pi_2(P_f)$.

Lemma 3.1. *The diagram*

$$\begin{array}{ccc} \pi_2(P_f, S^1) & \xrightarrow{\partial} & \pi_1(S^1) \\ \cong \downarrow & & \parallel \\ R & \xrightarrow{f \cdot \varepsilon_R} & \mathbb{Z} \end{array}$$

is an isomorphism of crossed modules, where $\varepsilon_R : R \rightarrow \mathbb{Z}$ is the augmentation.

Proof. By Whitehead's results [W1] on the free crossed module (2.2), it is enough to show that $\pi_2(P_f, S^1)$ is abelian. As ∂ is a totally free crossed module with basis \tilde{f} , $\pi_2(P_f, S^1)$ is generated by elements $e^n = ((e_2, n))$, see (2.1). Note that we obtain e^n by the action of $n \in \mathbb{Z}$ on $\iota(e_2) = ((e_2, 0)) = e^0$ and $\partial(e^n) = -n + \partial e + n = \partial e = f$ as $\pi_1(S^1) = \mathbb{Z}$ is abelian. We obtain

$$\begin{aligned} \langle e^n, e^m \rangle - \langle e^m, e^m \rangle &= -e^n - e^m + e^n + (e^m)^{\partial(e^n)} - (-e^m - e^m + e^m + (e^m)^{\partial(e^m)}) \\ &= -e^n - e^m + e^n + (e^m)^f - (e^m)^f + e^m \\ &= (e^n, e^m), \end{aligned}$$

where $(a, b) = -a - b + a + b$ denotes the commutator of a and b . Thus commutators of generators are sums of Peiffer commutators which are trivial in a crossed module. \square

With the notation of Theorem 2.2 and $M = \pi_2(P_f, S^1)$, Lemma 3.1 shows that $M = M^{cr} = (M^{cr})^{ab} = R$ and that $\pi_2(P_f) = \ker \partial = \ker \partial^{cr} = \ker(f \cdot \varepsilon) = K$ is the augmentation ideal of R , for $f \neq 0$. Thus the homotopy type of a pseudo-projective 3-space,

$$(3.2) \quad P_{f,x} = S^1 \cup e^2 \cup e^3,$$

is determined by the pair (f, x) of attaching maps, $f \in \pi_1(S^1) = \mathbb{Z}$ of the 2-cell e^2 , and $x \in \pi_2(P_f) = K \subseteq R$ of the 3-cell e^3 . We obtain the totally free nil(2)-module

$$(3.3) \quad M = \pi_2(P_{f,x}, S^1) \xrightarrow{\partial} N = \pi_1(S^1).$$

In the next section we use Theorem 2.2 to describe the group structure of $\pi_2(P_{f,x}, S^1)$, as well as the action of N on $\pi_2(P_{f,x}, S^1)$. The formulæ we derive are required to compute the homotopy group $\pi_3(P_{f,x})$ as a π_1 -module.

4. COMPUTATIONS IN NIL(2)-MODULES

In this Section we consider totally free nil(2)-modules, $\partial : M \rightarrow N$, generated by one element, e_i , in dimensions $i = 1, 2$, with basis $\tilde{f} : \{e_2\} \rightarrow N \cong \mathbb{Z}$. Then $\pi_1 = \text{coker } \partial = \mathbb{Z}/f\mathbb{Z}$ and, with $R = \mathbb{Z}[\pi_1]$, we obtain $(M^{cr})^{ab} = C = R$. Thus Theorem 2.2 yields the short exact sequence

$$(4.1) \quad (R \otimes R)/\Gamma(K) \xrightarrow{w} M \xrightarrow{q} R$$

with the image of $(R \otimes R)/\Gamma(K)$ central in M . This allows us to compute the group structure of M , as well as the action of $N = \mathbb{Z}$ on M , by computing the cross-effects of a set-theoretic splitting s of (4.1) with respect to addition and the action of N , even though here M need not be commutative.

The element $x \otimes y \in R \otimes R$ represents an equivalence class in $R \otimes R/\Gamma(K)$, also denoted by $x \otimes y$, so that $w(x \otimes y) = \langle \hat{x}, \hat{y} \rangle$ is the Peiffer commutator for $x, y \in R$, with $x = q(\hat{x})$ and $y = q(\hat{y})$. As a group, M is generated by elements $e^n = ((e_2, n))$, in particular, $e = e^0 = ((e_2, 0))$, see (2.1). We write

$$ke^n = \begin{cases} e^n + \dots + e^n & (k \text{ summands}) & \text{for } k > 0, \\ 0 & & \text{for } k = 0 \text{ and} \\ -e^n - \dots - e^n & (-k \text{ summands}) & \text{for } k < 0, \end{cases}$$

and define the set-theoretic splitting s of (4.1) by

$$s : R \longrightarrow M, \quad \sum_{k=0}^{f-1} x_{\bar{k}}[\bar{k}] \longmapsto x_{\bar{0}}e^0 + x_{\bar{1}}e^1 + \dots + x_{\bar{f-1}}e^{f-1}.$$

Then every $m \in M$ can be expressed uniquely as a sum $m = s(x) + w(m^\otimes)$ with $x \in R$ and $m^\otimes \in (R \otimes R)/\Gamma(K)$. The following formulæ for the cross-effects of s with respect to addition and the action provide a complete description of the nil(2)-module M in terms of R and $R \otimes R/\Gamma(K)$.

Given a function, $f : G \rightarrow H$, between groups, G and H , we write

$$(4.2) \quad f(x|y) = f(x + y) - (f(x) + f(y)), \quad \text{for } x, y \in G.$$

Lemma 4.1. *Take $x = \sum_{m=0}^{f-1} x_{\bar{m}}$, $y = \sum_{n=0}^{f-1} y_{\bar{n}}$ in R . Then*

$$s(x|y) = w(\nabla(x, y)),$$

where

$$\nabla(x, y) = \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} x_{\bar{m}} y_{\bar{n}} w([\bar{n}] \otimes [\bar{m}] - [\bar{m}] \otimes [\bar{n}]).$$

Thus $\nabla(x, y)$ is linear in x and y , yielding a homomorphism $\nabla : R \otimes R \rightarrow R \otimes R$.

Proof. First note that, by definition, $\nabla(k[\overline{m}], \ell[\overline{n}]) = 0$ unless $m > n$. To deal with the latter case, recall that commutators are central in M and use induction, first on k , then on ℓ , to show that

$$(ke^m, \ell e^n) = k\ell(e^m, e^n),$$

for $k, \ell > 0$. To show equality for negative k or ℓ , replace e^m or e^n by $-e^m$ and $-e^n$, respectively. Furthermore, note that the equality

$$(4.3) \quad (e^n, e^m) = -e^n - e^m + e^n + e^m = \langle e^n, e^m \rangle - \langle e^m, e^n \rangle$$

for commutators of generators of totally free cyclic crossed modules derived in the proof of Lemma 3.1 holds in any totally free $\text{nil}(n)$ -module generated by one element in each dimension. Taking $x = \sum_{m=0}^{f-1} x_{\overline{m}}[\overline{m}]$ and $y = \sum_{n=0}^{f-1} y_{\overline{n}}[\overline{n}]$, we obtain

$$\begin{aligned} s(x+y) &= (x_{\overline{0}} + y_{\overline{0}})e + \dots + (x_{\overline{m}} + y_{\overline{m}})e^m + \dots + (x_{\overline{f-1}} + y_{\overline{f-1}})e^{f-1} \\ &= (x_{\overline{i}}e + \dots + x_{\overline{f-1}}e^{f-1}) + (y_{\overline{0}}e + \dots + y_{\overline{f-1}}e^{f-1}) + \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} x_{\overline{m}}y_{\overline{n}}(e^n, e^m) \\ &= s(x) + s(y) + \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} x_{\overline{m}}y_{\overline{n}}(\langle e^n, e^m \rangle - \langle e^m, e^n \rangle) \\ &= s(x) + s(y) + \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} x_{\overline{m}}y_{\overline{n}}w([\overline{n}] \otimes [\overline{m}] - [\overline{m}] \otimes [\overline{n}]). \end{aligned}$$

□

Corollary 4.2. *Take $x \in R$ and $r \in \mathbb{Z}$. Then*

$$s(rx) = rs(x) + \binom{r}{2}w(\nabla(x, x)), \quad \text{where} \quad \binom{r}{2} = \frac{r(r-1)}{2}.$$

As $N = \mathbb{Z}$ is cyclic, the action of N on M is determined by the action of the generator, $1 \in \mathbb{Z}$. The formula for general $k \in \mathbb{Z}$ provided in the next lemma is required for the definition of the set-theoretic splitting u_x of (1.1) and the explicit computation of A and B in Theorem 1.3.

Lemma 4.3. *Take $x = \sum_{n=0}^{f-1} x_{\overline{n}}[\overline{n}] \in R$ and $\overline{k} \in \pi_1$. Write $R = \mathbb{Z}[\overline{0}, \dots, \overline{f-1}] = R_k \times \widehat{R}_k$, where $R_k = \mathbb{Z}[\overline{0}, \dots, \overline{f-k-1}]$ and $\widehat{R}_k = \mathbb{Z}[\overline{f-k}, \dots, \overline{f-1}]$. Then*

$$(s(x))^{\overline{k}} = s(x^{\overline{k}}) + w(\overline{\nabla}_k(a, b)),$$

where $x = (a, b)$ and

$$\overline{\nabla}_k : R_k \times \widehat{R}_k \rightarrow R \otimes R, \quad (a, b) \mapsto Q_k(a, b) + L_k(b)$$

with

$$\begin{aligned} Q_k(a, b) &= \sum_{p=0}^{f-\ell-1} \sum_{q=0}^{\ell-1} x_{\overline{p}}x_{\overline{q+f-\ell}}([\overline{p+\ell}] \otimes [\overline{q}] - [\overline{q}] \otimes [\overline{q}]) \\ L_k(b) &= \sum_{q=0}^{\ell-1} x_{\overline{q+f-\ell}}[\overline{q}] \otimes [\overline{q}]. \end{aligned}$$

Thus Q_k is linear in a and b and L_k is linear in b .

Proof. For $\overline{j} \in \pi_1$ and $p \in \mathbb{Z}$,

$$\begin{aligned} e^{j+f} &= (e^j)^{\partial(e)} \\ &= e^j + (e^j, e) + \langle e, e^j \rangle \\ &= e^j - (\langle e, e^j \rangle - \langle e^j, e^j \rangle) + \langle e, e^j \rangle \\ &= e^j + \langle e^j, e^j \rangle. \end{aligned}$$

Thus, for $\bar{n}, \bar{k} \in \pi_1$, with $\bar{n} + \bar{k} = \bar{j}$,

$$\begin{aligned} (s([\bar{n}]))^k &= \begin{cases} e^j, & \text{for } 0 \leq n < f - k, \\ e^j + \langle e^j, e^j \rangle, & \text{for } f - k \leq n < f \end{cases} \\ &= \begin{cases} s([\bar{n}]^{\bar{k}}), & \text{for } 0 \leq n < f - k, \\ s([\bar{n}]^{\bar{k}}) + w([\bar{j}] \otimes [\bar{j}]), & \text{for } f - k \leq n < f. \end{cases} \end{aligned}$$

Hence, for $x = \sum_{p=0}^{f-1} x_{\bar{p}} [\bar{p}]$,

$$\begin{aligned} &(s(x))^k \\ &= x_{\bar{0}} s([\bar{0}])^k + x_{\bar{1}} s([\bar{1}])^k + \dots + x_{\bar{f-1}} s([\bar{f-1}])^k \\ &= x_{\bar{0}} s([\bar{0}]^{\bar{k}}) + x_{\bar{1}} s([\bar{1}]^{\bar{k}}) + \dots + x_{\bar{f-1}} s([\bar{f-1}]^{\bar{k}}) + \sum_{n=f-k}^{f-1} x_{\bar{n}} w([\bar{n} + k - f] \otimes [\bar{n} + k - f]) \\ &= x_{\bar{f-k}} s([\bar{f-k}]^{\bar{k}}) + \dots + x_{\bar{f-1}} s([\bar{f-1}]^{\bar{k}}) + x_{\bar{0}} s([\bar{0}]^{\bar{k}}) + \dots + x_{\bar{f-k-1}} s([\bar{f-k-1}]^{\bar{k}}) \\ &\quad + \sum_{p=0}^{f-k-1} \sum_{n=f-k}^{f-1} (x_{\bar{p}} s([\bar{p} + \bar{k}]), x_{\bar{n}} s([\bar{n} + \bar{k}])) + \sum_{q=0}^{k-1} x_{\bar{q+f-k}} w([\bar{q}] \otimes [\bar{q}]) \\ &= s(x^{\bar{k}}) + \sum_{p=0}^{f-k-1} \sum_{q=0}^{k-1} x_{\bar{p}} x_{\bar{q+f-k}} w([\bar{p} + \bar{k}] \otimes [\bar{q}] - [\bar{q}] \otimes [\bar{q}]) + \sum_{q=0}^{k-1} x_{\bar{q+f-k}} w([\bar{q}] \otimes [\bar{q}]). \end{aligned}$$

□

Remark 4.4. We use the final results of this section to define and establish the properties of the set-theoretic splitting u_x of (1.1). The next result shows how the cross-effects interact with multiplication in R .

Lemma 4.5. *Take $x, y \in R$. Then*

$$\sum_{i=0}^{f-1} y_{\bar{i}} (s(x))^i = s(xy) + w(\mu(x, y)),$$

where $\mu : R \times R \rightarrow R \otimes R$ is given by

$$\mu(x, y) = - \sum_{i < j} y_{\bar{i}} y_{\bar{j}} \nabla(x^{\bar{i}}, x^{\bar{j}}) + \sum_{i=0}^{f-1} (\bar{\nabla}_i(y_{\bar{i}} x) - \binom{y_{\bar{i}}}{2} \nabla(x, x^{\bar{i}})).$$

Proof. By Lemmata 4.1 and 4.3 and Corollary 4.2, we obtain, for $x, y \in R$,

$$\begin{aligned} \sum_{i=0}^{f-1} y_{\bar{i}} (s(x))^i &= \sum_{i=0}^{f-1} (y_{\bar{i}} s(x))^i \\ &= \sum_{i=0}^{f-1} (s(y_{\bar{i}} x) - \binom{y_{\bar{i}}}{2} w(\nabla(x, x)))^i \\ &= \sum_{i=0}^{f-1} s(y_{\bar{i}} x^{\bar{i}}) + w(\bar{\nabla}_i(y_{\bar{i}} x)) - \left(\binom{y_{\bar{i}}}{2} w(\nabla(x, x)) \right)^i \\ &= s\left(\sum_{i=0}^{f-1} y_{\bar{i}} x^{\bar{i}}\right) - \sum_{i < j} w(\nabla(y_{\bar{i}} x^{\bar{i}}, y_{\bar{j}} x^{\bar{j}})) + \sum_{i=0}^{f-1} w(\bar{\nabla}_i(y_{\bar{i}} x)) - \binom{y_{\bar{i}}}{2} w(\nabla(x, x^{\bar{i}})). \end{aligned}$$

□

Finally, the definitions and a simple calculation yield

Lemma 4.6. For $x, y, z \in R$ and with the notation in (4.2),

$$\mu(x, y|z) = - \sum_{i < j} (y_{\bar{i}} z_{\bar{j}} + z_{\bar{i}} y_{\bar{j}}) \nabla(x^{\bar{i}}, x^{\bar{j}}) + 2 \sum_{i=1}^{f-1} y_{\bar{i}} z_{\bar{i}} Q_i(x) - \sum_{i=0}^{f-1} y_{\bar{i}} z_{\bar{i}} \nabla(x, x)^{\bar{i}}.$$

Hence, for fixed $x \in R$, $\mu(x, \cdot) : R \times R \rightarrow R \otimes R$, $(y, z) \mapsto \mu(x, y|z)$ is bilinear.

5. QUADRATIC MODULES

In dimension 3, quadratic modules assume the role played by crossed modules in dimension 2. We recall the notion of quadratic modules and totally free quadratic modules, see [B], which we require for the description of the third homotopy group $\pi_3(P_{f,x})$ of a 3-dimensional pseudo-projective space $P_{f,x}$, as in (3.2).

A *quadratic module* $(\omega, \delta, \partial)$ consists of a commutative diagram of group homomorphisms

$$\begin{array}{ccccc} & & C \otimes C & & \\ & \swarrow \omega & \downarrow w & & \\ L & \xrightarrow{\delta} & M & \xrightarrow{\partial} & N, \end{array}$$

such that

- $\partial : M \rightarrow N$ is a nil(2)-module with quotient map $M \twoheadrightarrow C = (M^{cr})^{ab}$, $x \mapsto \{x\}$, and Peiffer commutator map w given by $w(\{x\} \otimes \{y\}) = \langle x, y \rangle$;
- the *boundary homomorphisms* ∂ and δ satisfy $\partial\delta = 0$, and the *quadratic map* ω is a lift of w , that is, for $x, y \in M$,

$$\delta\omega(\{x\} \otimes \{y\}) = \langle x, y \rangle;$$

- N acts on L , all homomorphisms are equivariant with respect to the action of N and, for $a \in L$ and $x \in M$,

$$(5.1) \quad a^{\partial(x)} = a + \omega(\{\delta a\} \otimes \{x\} + \{x\} \otimes \{\delta a\});$$

- finally, for $a, b \in L$,

$$(5.2) \quad (a, b) = -a - b + a + b = \omega(\{\delta a\} \otimes \{\delta b\}).$$

A map $\varphi : (\omega, \delta, \partial) \rightarrow (\omega', \delta', \partial')$ of quadratic modules is given by a commutative diagram

$$\begin{array}{ccccccc} C \otimes C & \xrightarrow{\omega} & L & \xrightarrow{\delta} & M & \xrightarrow{\partial} & N \\ \varphi_* \otimes \varphi_* \downarrow & & \downarrow l & & \downarrow m & & \downarrow n \\ C' \otimes C' & \xrightarrow{\omega'} & L' & \xrightarrow{\delta'} & M' & \xrightarrow{\partial'} & N' \end{array}$$

where l is n -equivariant, and (m, n) is a map between pre-crossed modules inducing $\varphi_* : C \rightarrow C'$.

Given a nil(2)-module $\partial : M \rightarrow N$, a free group F and a homomorphism $\tilde{f} : F \rightarrow M$ with $\partial\tilde{f} = 0$, a quadratic module $(\omega, \delta, \partial)$ is *free with basis* \tilde{f} , if there is a homomorphism $i : F \rightarrow L$ with $\delta i = \tilde{f}$, such that the following universal property is satisfied: For every quadratic module $(\omega', \delta', \partial')$ and map $(m, n) : \partial \rightarrow \partial'$ of nil(2)-modules and every homomorphism $l_F : F \rightarrow L'$ with $m\tilde{f} = \delta' l_F$, there is a unique map (l, m, n) of quadratic modules with $li = l_F$.

$$\begin{array}{ccccc} L & \xrightarrow{\delta} & M & \xrightarrow{\partial} & N \\ \downarrow l & \swarrow i & \nearrow \tilde{f} & & \downarrow n \\ & & F & & \\ \downarrow l_F & \swarrow & \nearrow m & & \\ L' & \xrightarrow{\delta'} & M' & \xrightarrow{\partial'} & N' \end{array}$$

For $F = \langle Z \rangle$, the homomorphism \tilde{f} is determined by its restriction $\tilde{f}|_Z$ which is then called a *basis* for $(\omega, \delta, \partial)$. A quadratic module $(\omega, \delta, \partial)$ is *totally free* if it is free, if ∂ is a free nil(2)-module and if N is a free group.

6. THE HOMOTOPY GROUP π_3 OF A PSEUDO-PROJECTIVE 3-SPACE AND THE ASSOCIATED SPLITTING FUNCTION u_x

In this section we return to pseudo-projective 3-spaces

$$P_{f,x} = S^1 \cup e^2 \cup e^3,$$

determined by the pair (f, x) of attaching maps, $f \in \pi_1(S^1) = \mathbb{Z}$ and $x \in \pi_2(P_f) = K \subseteq R$, as in (3.2). Using results on totally free quadratic modules in [B], we investigate the structure of the third homotopy group $\pi_3(P_{f,x})$ as a π_1 -module by defining a set-theoretic splitting u_x of J.H.C. Whitehead's Certain Exact Sequence of the universal cover, $\widehat{P}_{f,x}$,

$$(6.1) \quad \Gamma(\pi_2(P_{f,x})) \twoheadrightarrow \pi_3(P_{f,x}) \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{u_x} \end{array} \mathbb{H}_3(\widehat{P}_{f,x}).$$

Recall that $\pi_1 = \pi_1(P_f) = \mathbb{Z}/f\mathbb{Z}$ with augmentation ideal $K = \ker f\varepsilon$, and let B be the image of $d_x : R \rightarrow R, y \mapsto xy$. Then

$$(6.2) \quad \pi_2(P_{f,x}) = \mathbb{H}_2(\widehat{P}_{f,x}) = K/B = (\ker f\varepsilon)/xR.$$

The functor σ in (IV 6.8) in [B] assigns a totally free quadratic module $(\omega, \delta, \partial)$ to the pseudo-projective 3-space $P_{f,x}$ and we obtain the commutative diagram

$$\begin{array}{ccccccc} \Gamma(\pi_2(P_{f,x})) & \twoheadrightarrow & R \otimes R/\Delta_B & \xrightarrow{q} & R \otimes R/\Gamma(K) & & \\ \downarrow & & \downarrow \omega & & \downarrow w & & \\ \pi_3(P_{f,x}) & \twoheadrightarrow & L & \xrightarrow{\delta} & M & \xrightarrow{\partial} & N \\ \downarrow u_x & & \downarrow t_x & & \downarrow s & & \parallel \\ \mathbb{H}_3(\widehat{P}_{f,x}) & \twoheadrightarrow & R & \xrightarrow{d_x} & R & \xrightarrow{f \cdot \varepsilon} & \mathbb{Z} \end{array}$$

of straight arrows. Here the generators $e_3 \in L$, $e_2 \in M$ and $e_1 = 1 \in N = \mathbb{Z}$ correspond to the cells of $P_{f,x}$ and ∂ is the totally free nil(2)-module of Lemma 3.1. The right hand column is the short exact sequence (4.1) with the set theoretic splitting s defined in Section 4. The short exact sequence in the middle column is described in (IV 2.13) in [B], where the product $[\alpha, \beta]$ of $\alpha \in K$ and $\beta \in B$ is given by $[\alpha, \beta] = \alpha \otimes \beta + \beta \otimes \alpha \in R \otimes R$ and

$$\Delta_B = \Gamma(B) + [K, B].$$

By Corollary (IV 2.14) in [B], taking kernels yields Whitehead's short exact sequence (6.1) in the left hand column of the diagram, that is, $\ker q = \Gamma(\pi_2(\widehat{P}_{f,x}))$, $\ker \delta = \pi_3(P_{f,x})$ and $\ker d_x = \mathbb{H}_3(\widehat{P}_{f,x})$. As $(\omega, \delta, \partial)$ is a quadratic module associated to $P_{f,x}$, we may assume that $\delta(e_3) = s(x)$.

In Section 4 we determined the structure of M as an N -module by computing the cross-effects of the set-theoretic splitting s with respect to addition and the action. Analogously to the definition of s , we now define a set-theoretic splitting of the short exact sequence in the second column of this diagram by

$$t_x : R \longrightarrow L, \quad \sum_{k=0}^{f-1} y_{\overline{k}} [\overline{k}] \longmapsto y_{\overline{0}} e_3^0 + \dots + y_{\overline{f-1}} e_3^{f-1}.$$

The cross-effects of t_x with respect to addition and the action determine the N -module structure of L , but we want to determine the module structure of $\pi_3(P_{f,x})$. To obtain a set-theoretic splitting of the first column which will allow us to do so, we must adjust t_x , such that the image of $\mathbb{H}_3(\widehat{P}_{f,x})$ under the new splitting is contained in $\ker \delta = \pi_3(P_{f,x})$. Recall that δ is a homomorphism which

is equivariant with respect to the action of N and $\delta(e_3) = s(x)$. Thus Lemma 4.5 yields, for $y \in \mathbb{H}_3(\widehat{P}_{f,x}) = \ker d_x$, that is, for $d_x(y) = xy = 0$,

$$\begin{aligned} \delta(t_x(y)) &= \delta\left(\sum_{i=0}^{f-1} y_i e_3^{\bar{i}}\right) = \sum_{i=0}^{f-1} y_i \delta(e_3)^{\bar{i}} = \sum_{i=0}^{f-1} y_i (s(x))^{\bar{i}} \\ &= s(xy) + w(\mu(x, y)) \\ &= \delta\omega\mu(x, y). \end{aligned}$$

Hence $t_x(y) - \omega\mu(x, y) \in \ker \delta = \pi_3(P_{f,x})$, giving rise to the set theoretic splitting

$$u_x : \mathbb{H}_3(\widehat{P}_{f,x}) \longrightarrow \pi_3(P_{f,x}), \quad y \longmapsto t_x(y) - \omega\mu(x, y)$$

of the Hurewicz map $\pi_3 \rightarrow \mathbb{H}_3$. The cross-effects of u_x with respect to addition and the action determine (6.1) as a short exact sequence of π_1 -modules. In Section 7 we determine the cross-effects of t_x and investigate the properties of the functions A and B describing the cross-effects of u_x .

7. COMPUTATIONS IN FREE QUADRATIC MODULES

The first two results of this Section describe the cross-effects of t_x with respect to addition and the action, respectively. We then turn to the properties of the cross-effects of u_x .

Lemma 7.1. *Take $z, y \in R$. Then, with the notation in (4.2),*

$$t_x(z|y) = \omega(\Psi(z, y)),$$

where

$$\Psi(z, y) = \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} z_{\bar{m}} y_{\bar{n}} x[\bar{n}] \otimes x[\bar{m}].$$

Thus $\Psi(z, y)$ is linear in z and y , yielding a homomorphism $\Psi : R \otimes R \rightarrow R \otimes R$.

Proof. As in the proof of Lemma 4.1, we obtain

$$t_x(z|y) = \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} z_{\bar{m}} y_{\bar{n}} (e_3^{\bar{n}}, e_3^{\bar{m}}).$$

Note that $\{\delta(e_3^{\bar{n}})\} = \{\delta(t_x([\bar{n}]))\} = d_x([\bar{n}]) = x[\bar{n}]$. Thus (5.2) yields

$$t_x(z|y) = \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} z_{\bar{m}} y_{\bar{n}} \omega(\{\delta(e_3^{\bar{n}})\} \otimes \{\delta(e_3^{\bar{m}})\}) = \sum_{m=1}^{f-1} \sum_{n=0}^{m-1} z_{\bar{m}} y_{\bar{n}} \omega(x[\bar{n}] \otimes x[\bar{m}]).$$

□

As $N = \mathbb{Z}$ is cyclic, the action of N on L is determined by the generator $1 \in \mathbb{Z}$.

Lemma 7.2. *Take $x \in R$. Then*

$$(t_x(y))^1 = t_x(y^{\bar{1}}) + \omega(\overline{\Psi}_1(a, b)),$$

where

$$\overline{\Psi}_1 = \sum_{p=0}^{f-2} y_{\bar{p}} y_{\overline{f-1}} x[\overline{p+1}] \otimes x[\bar{0}] + y_{\overline{f-1}} (x \otimes [\bar{0}] + [\bar{0}] \otimes x).$$

Proof. With $\{\delta(e_3^{\bar{n}})\} = x[\bar{n}]$ from above and (5.1), we obtain

$$\begin{aligned} e_3^{1+f} &= (e_3^1)^f = (e_3^1)^{\partial(e)} = e^1 + \omega(\{\delta(e_3^1)\} \otimes \{e\} + \{e\} \otimes \{\delta(e_3^1)\}) \\ &= t_x([\bar{n}]^{\bar{1}}) + \omega(x[\bar{1}] \otimes [\bar{0}] + [\bar{0}] \otimes x[\bar{1}]). \end{aligned}$$

Thus, for $\bar{n} \in \pi$,

$$(t_x([\bar{n}]))^1 = \begin{cases} \omega(t_x([\bar{n}]^{\bar{1}})) & \text{for } 0 \leq n < f-1, \\ \omega(t_x([\bar{n}]^{\bar{1}}) + x[\bar{1}] \otimes [\bar{0}] + [\bar{0}] \otimes x[\bar{1}]) & \text{for } n = f-1. \end{cases}$$

With (5.2), we obtain, for $y = \sum_{n=0}^{f-1} y_{\overline{n}}[\overline{n}]$,

$$\begin{aligned}
(t_x(y))^1 &= y_{\overline{0}} e_3^1 + y_{\overline{1}} e_3^2 \dots + y_{\overline{f-2}} e_3^{f-1} + y_{\overline{f-1}} e_3^f \\
&= y_{\overline{0}} t_x([\overline{0}]^{\overline{1}}) + \dots + y_{\overline{f-2}} t_x([\overline{f-1}]^{\overline{1}}) + y_{\overline{f-1}} t_x([\overline{f-1}]^{\overline{1}}) + y_{\overline{f-1}} \omega(x \otimes [\overline{0}] + [\overline{0}] \otimes x) \\
&= t_x(y^{\overline{1}}) + \sum_{p=0}^{f-2} y_{\overline{p}} y_{\overline{f-1}} (e_3^{p+1}, e_3) + y_{\overline{f-1}} \omega(x \otimes [\overline{0}] + [\overline{0}] \otimes x) \\
&= t_x(y^{\overline{1}}) + \sum_{p=0}^{f-2} y_{\overline{p}} y_{\overline{f-1}} x[\overline{p+1}] \otimes x[\overline{0}] + y_{\overline{f-1}} (x \otimes [\overline{0}] + [\overline{0}] \otimes x)
\end{aligned}$$

□

The next two results concern the properties of the maps A and B which describe the cross-effects of u_x with respect to addition and the action, respectively.

Lemma 7.3. *For $x \in K$ the map*

$$A : H_3 \widehat{P}_{f,x} \times H_3 \widehat{P}_{f,x} \rightarrow \Gamma(\pi_2 P_{f,x}), (y, z) \mapsto u_x(y|z)$$

is bilinear.

Proof. Take $x \in K$ and $y, z \in H_3 \widehat{P}_{f,x}$. By definition

$$A(y, z) = u_x(y|z) = t_x(y|z) - \omega \mu(x, y|z) = \omega(\Psi(y, z) - \mu(x, y|z)).$$

Thus Lemmata 4.6 and 7.1 imply that A is bilinear. □

Lemma 7.4. *For $x \in K$ define*

$$B : H_3 \widehat{P}_{f,x} \rightarrow \Gamma(\pi_2 P_{f,x}), y \mapsto (u_x(y))^1 - u_x(y^1)$$

Then

$$H_3 \widehat{P}_{f,x} \times H_3 \widehat{P}_{f,x} \rightarrow \Gamma(\pi_2 P_{f,x}), (y, z) \mapsto B(y|z)$$

is bilinear.

Proof. Take $x \in K$ and $y, z \in H_3 \widehat{P}_{f,x}$. Then

$$\begin{aligned}
(A(y, z))^1 &= (u_x(y+z) - (u_x(y) + u_x(z)))^1 \\
&= (u_x(y+z))^1 - (u_x(y))^1 - (u_x(z))^1 \\
&= B(y+z) + u_x((y+z)^1) - (B(y) + u_x(y^1) + B(z) + u_x(z^1)). \\
&= B(y|z) + A(y^1, z^1)
\end{aligned}$$

Thus

$$(7.1) \quad B(y|z) = (A(y, z))^1 - A(y^1, z^1)$$

and bilinearity follows from that of A and the properties of an action. □

8. EXAMPLES OF PSEUDO-PROJECTIVE 3-SPACES

In this Section we provide explicit computations for examples of pseudo-projective 3-spaces, including proofs for Theorem 1.1, Theorem 1.3 and Theorem 1.4.

Note that, as abelian group, the augmentation ideal K of a pseudo-projective 3-space $P_{f,x}$, as in (3.2), is freely generated by $\{[\overline{1}] - [\overline{0}], \dots, [\overline{f-1}] - [\overline{0}]\}$. We consider pseudo-projective 3-spaces, $P_{f,x}$, with $x = \tilde{x}([\overline{1}] - [\overline{0}])$ and $\tilde{x} \in \mathbb{Z}$. We compute $\pi_2(P_{f,x}), H_3(\widehat{P}_{f,x})$, as well as the cross-effects of u_x for this special case. For $f = 2$, the general case coincides with the special case and provides an example where π_1 acts trivially on $\Gamma\pi_2(P_{2,\tilde{x}})$ and on $H_3(\widehat{P}_{2,\tilde{x}})$, but non-trivially on $\pi_3(P_{2,\tilde{x}})$.

Lemma 8.1. For $x = \tilde{x}([\overline{1}] - [\overline{0}])$ with $\tilde{x} \in \mathbb{Z}$,

$$H_3(\widehat{P}_{f,x}) = \{\tilde{y}N \mid \tilde{y} \in \mathbb{Z}\} \cong \mathbb{Z},$$

is generated by the norm element $N = \sum_{k=0}^{f-1} [\overline{k}]$. Hence π_1 acts trivially on $H_3(\widehat{P}_{f,x})$. Furthermore,

$$\pi_2(P_{f,x}) = (\mathbb{Z}/\tilde{x}\mathbb{Z}) \otimes_{\mathbb{Z}} K.$$

Hence $\tilde{x}^2\ell = 0$ for every $\ell \in \Gamma(\pi_2(P_{f,x}))$.

Proof. Take $x = \tilde{x}([\overline{1}] - [\overline{0}])$ with $\tilde{x} \in \mathbb{Z}$ and $y = \sum_{k=0}^{f-1} y_{\overline{k}}[\overline{k}] \in \ker d_x$. Then

$$\begin{aligned} d_x(y) = xy = 0 &\iff \tilde{x} \sum_{k=0}^{f-1} y_{\overline{k}}([\overline{k+1}] - [\overline{k}]) = 0 \\ &\iff y_{\overline{f-1}} = y_{\overline{0}} = y_{\overline{1}} = y_{\overline{2}} = \dots = y_{\overline{f-2}} = \tilde{y}, \end{aligned}$$

for some $\tilde{y} \in \mathbb{Z}$. Hence $y = \tilde{y}N$.

By (6.2), $\pi_2(P_{f,x}) = K/xR$. As abelian group, $K = \ker \varepsilon$ is freely generated by $\{[\overline{k}] - [\overline{0}]\}_{1 \leq k \leq f-1}$ and hence also by $\{[\overline{k}] - [\overline{k-1}]\}_{1 \leq k \leq f-1}$. For $y = \sum_{i=0}^{f-1} y_{\overline{i}}[\overline{i}] \in R$ we obtain

$$\begin{aligned} xy &= \tilde{x} \sum_{i=1}^{f-1} y_{\overline{i}}([\overline{i}] - [\overline{i-1}]) + \tilde{x}y_{\overline{f-1}}([\overline{0}] - [\overline{f-1}]) \\ &= \tilde{x} \sum_{i=1}^{f-1} y_{\overline{i}}([\overline{i}] - [\overline{i-1}]) - \tilde{x}y_{\overline{f-1}} \sum_{i=1}^{f-1} ([\overline{i}] - [\overline{i-1}]) \\ &= \tilde{x} \sum_{i=1}^{f-1} (y_{\overline{i}} - y_{\overline{f-1}})([\overline{i}] - [\overline{i-1}]). \end{aligned}$$

As $\tilde{x}K \subseteq xR$, we obtain $xR = \tilde{x}K$ and hence

$$\pi_2(P_{f,x}) = K/xR = K/\tilde{x}K = (\mathbb{Z}/\tilde{x}\mathbb{Z}) \otimes_{\mathbb{Z}} K.$$

If \tilde{x} is odd, then every element $\ell \in \Gamma(\pi_2(P_{f,x}))$ has order \tilde{x} . If \tilde{x} is even, an element $\ell \in \Gamma(\pi_2(P_{f,x}))$ has order $2\tilde{x}$ or \tilde{x} . In either case, $\tilde{x}^2\ell = 0$ for every $\ell \in \Gamma(\pi_2(P_{f,x}))$. \square

Lemma 8.2. Take $x = \tilde{x}([\overline{1}] - [\overline{0}])$ and $y, z \in H_3(\widehat{P}_{f,x})$. Then

$$A(y, z) = 0.$$

Proof. By definition,

$$A(y, z) = u_x(y|z) = t_x(y|z) - \omega\mu(x, y|z) = \omega(\Psi(y, z) - \mu(x, y|z)).$$

The definition of Ψ and Lemma 4.6 yield

$$\Psi(y, z) - \mu(x, y|z) = \tilde{y}\tilde{z} \left(\sum_{p=1}^{f-1} \sum_{q=0}^{p-1} x[\overline{q}] \otimes x[\overline{p}] + 2 \sum_{q=1}^{f-1} \sum_{p=0}^{q-1} \nabla(x^{\overline{p}}, x^{\overline{q}}) - 2 \sum_{p=1}^{f-1} Q_p(x) + \sum_{p=0}^{f-1} (\nabla(x, x))^{\overline{p}} \right).$$

Recall that $\tilde{x}^2\ell = 0$ for every $\ell \in \Gamma(\pi_2(P_{f,x}))$ and note that, by the properties of Q and ∇ , each summand in the above sum has a factor of \tilde{x}^2 . \square

Lemma 8.3. Let $\gamma : \pi_2(P_{f,x}) \rightarrow \Gamma(\pi_2(P_{f,x}))$ be the universal quadratic map for the Whitehead functor Γ . Take $q : K \rightarrow \pi_2(P_{f,x})$, $k \mapsto 1 \otimes k$, $x = \tilde{x}([\overline{1}] - [\overline{0}])$ and $y = \tilde{y}N$. Then

$$B(y) = -\tilde{x}\tilde{y}\gamma q([\overline{1}] - [\overline{0}]).$$

Proof. Note that $y^\beta = y$ for $\beta \in \pi_1$. As $\tilde{x}^2 \ell = 0$ for every $\ell \in \Gamma(\pi_2(P_{f,x}))$, any summand with a factor \tilde{x}^2 is equal to 0. By Lemma 7.2,

$$\begin{aligned} \overline{\Psi}_1(y) &= \sum_{p=0}^{f-2} \tilde{y}^2 (\tilde{x}([\overline{1}] - [\overline{0}])[\overline{p+1}] \otimes (\tilde{x}([\overline{1}] - [\overline{0}])) + \tilde{y}(\tilde{x}([\overline{1}] - [\overline{0}]) \otimes [\overline{0}] + [\overline{0}] \otimes \tilde{x}([\overline{1}] - [\overline{0}])) \\ &= \tilde{x}\tilde{y}([\overline{1}] - [\overline{0}]) \otimes [\overline{0}] + [\overline{0}] \otimes ([\overline{1}] - [\overline{0}]). \end{aligned}$$

Lemma 4.5 yields

$$\begin{aligned} \mu(x, y) &= - \sum_{q=0}^{f-1} \sum_{p=0}^{q-1} \tilde{x}^2 \tilde{y}^2 \nabla([\overline{p+1}] - [\overline{p}], [\overline{q+1}] - [\overline{q}]) + \sum_{p=0}^{f-1} \overline{\nabla}_p(\tilde{y}\tilde{x}([\overline{1}] - [\overline{0}])) \\ &\quad - \tilde{x}^2 \binom{\tilde{y}}{2} (\nabla([\overline{1}] - [\overline{0}], ([\overline{1}] - [\overline{0}]))^{\overline{p}} \\ &= \overline{\nabla}_{f-1}(\tilde{x}\tilde{y}([\overline{1}] - [\overline{0}])) \\ &= -\tilde{x}^2 \tilde{y}^2 ([\overline{f-1}] \otimes [\overline{0}] - [\overline{0}] \otimes [\overline{0}]) + \tilde{x}\tilde{y} [\overline{0}] \otimes [\overline{0}] \\ &= \tilde{x}\tilde{y} [\overline{0}] \otimes [\overline{0}]. \end{aligned}$$

Thus

$$B(y) = (u_x(y))^1 - u_x(y^{\overline{1}}) = \omega(\overline{\Psi}_1(y) - (\mu(x, y))^1 + \mu(x, y)) = -\tilde{x}\tilde{y}\gamma q([\overline{1}] - [\overline{0}]).$$

□

Together Lemmata 8.1, 8.2 and 8.3 provide a proof of Theorem 1.3.

For $f = 2$ the special case coincides with the general case and we obtain

Theorem 8.4. *Let $X = P_{2,x}$ be a pseudo-projective 3-space with $x = \tilde{x}([\overline{1}] - [\overline{0}])$, for $\tilde{x} \in \mathbb{Z}$ and $\tilde{x} \neq 0$. Then u_x is a homomorphism and the fundamental group $\pi_1 = \mathbb{Z}/2\mathbb{Z}$ acts trivially on $\Gamma(\pi_2 P_{2,x})$ and on $H_3 \widehat{P}_{2,x}$. The action of π_1 on $\pi_3 P_{2,x}$ is non-trivial if and only if \tilde{x} is even.*

Proof. For $f = 2$ the augmentation ideal K is generated by $k = [\overline{1}] - [\overline{0}]$. Since $k[\overline{1}] = -k$, the action of $\pi_1 = \mathbb{Z}/2\mathbb{Z}$ on K and hence on $\pi_2 P_{2,x} = K/xR = \mathbb{Z}/\tilde{x}\mathbb{Z}$ is multiplication by -1 . As the Γ -functor maps multiplication by -1 to the identity morphism, the action on π_1 on $\Gamma(\pi_2 P_{2,x})$ is trivial. The group $H_3 \widehat{P}_{2,x}$ is generated by the norm element $N = [\overline{0}] + [\overline{1}]$. As $N[\overline{1}] = N$, π_1 acts trivially on $H_3 \widehat{P}_{2,x}$. As $\pi_2 = \mathbb{Z}/\tilde{x}\mathbb{Z}$ is cyclic, $\Gamma\pi_2 = \pi_2$ if \tilde{x} is odd and $\Gamma\pi_2 = \mathbb{Z}/2\tilde{x}\mathbb{Z}$ if \tilde{x} is even, that is,

$$(8.1) \quad \Gamma\pi_2 = \mathbb{Z}/\gcd(\tilde{x}, 2)\tilde{x}\mathbb{Z}.$$

By Lemma 8.3 and (8.1), the action of π_1 on $\pi_3 X$ is non-trivial if and only if \tilde{x} is even. □

Theorem 1.1 is a corollary to Theorem 8.4.

Proof of 1.4. Note that $\mathbb{Z}/\tilde{x}\mathbb{Z} \otimes_{\mathbb{Z}} K$ is generated by $\{\alpha_k = q([\overline{k}] - [\overline{k-1}])\}_{0 < k < f}$, where $q : K \rightarrow \mathbb{Z}/\tilde{x}\mathbb{Z} \otimes_{\mathbb{Z}} K, k \mapsto 1 \otimes k$. Thus $\Gamma(\pi_2(P_{f,x})) = \Gamma(\mathbb{Z}/\tilde{x}\mathbb{Z} \otimes K) \subseteq (\mathbb{Z}/\tilde{x}\mathbb{Z} \otimes_{\mathbb{Z}} K) \otimes (\mathbb{Z}/\tilde{x}\mathbb{Z} \otimes_{\mathbb{Z}} K)$ is generated by $\{\gamma q(\alpha_k), [q(\alpha_j), q(\alpha_k)]\}_{0 < j < k, 0 < k < f}$. With $\alpha_k^1 = \alpha_{k+1}$ for $1 < k < f-1$ and $\alpha_{f-1}^1 = [\overline{0}] - [\overline{f-1}] = -\sum_{i=1}^{f-1} \alpha_i$, we obtain, for $\ell = \sum_{k=1}^{f-1} \ell_k \gamma(\alpha_k) + \sum_{k=2}^{f-1} \sum_{j=1}^{k-1} \ell_{j,k} [\alpha_j, \alpha_k] \in$

$\Gamma(\pi_2(P_{f,\tilde{x}})),$

$$\begin{aligned}
\ell^1 - \ell &= \sum_{k=1}^{f-1} \ell_k \gamma q(\alpha_k)^1 + \sum_{k=2}^{f-1} \sum_{j=1}^{k-1} \ell_{j,k} [q(\alpha_j), q(\alpha_k)]^1 - \sum_{k=1}^{f-1} \ell_k \gamma q(\alpha_k) - \sum_{k=2}^{f-1} \sum_{j=1}^{k-1} \ell_{j,k} [q(\alpha_j), q(\alpha_k)] \\
&= \sum_{k=1}^{f-2} \ell_k \gamma q(\alpha_{k+1}) + \ell_{f-1} \gamma q(-\sum_{i=1}^{f-1} \alpha_i) + \sum_{k=2}^{f-2} \sum_{j=1}^{k-1} \ell_{j,k} [q(\alpha_{j+1}), q(\alpha_{k+1})] \\
&\quad + \sum_{j=1}^{f-1} \ell_{j,f-1} [\gamma q(\alpha_{j+1}), \gamma q(-\sum_{i=1}^{f-1} \alpha_i)] - \sum_{k=1}^{f-1} \ell_k \gamma q(\alpha_k) - \sum_{k=2}^{f-1} \sum_{j=1}^{k-1} \ell_{j,k} [q(\alpha_j), q(\alpha_k)] \\
&= (\ell_{f-1} - \ell_1) \gamma q(\alpha_1) + \sum_{k=2}^{f-1} (\ell_{k-1} - \ell_k + \ell_{f-1} - 2\ell_{k-1,f-1}) \gamma q(\alpha_k) \\
&\quad + \sum_{k=2}^{f-1} (\ell_{f-1} - \ell_{1,k} - \ell_{k-1,f-1}) [q(\alpha_1), q(\alpha_k)] \\
&\quad + \sum_{k=3}^{f-1} \sum_{j=2}^{k-1} (\ell_{f-1} + \ell_{j-1,k-1} - \ell_{j,k} - \ell_{j-1,f-1} - \ell_{k-1,f-1}) [q(\alpha_j), q(\alpha_k)].
\end{aligned}$$

Thus the sequence (1.1) splits if and only if there is at least one solution of the system of equations

$$\begin{aligned}
(A) \quad & 0 = \ell_{f-1} - \ell_1 && \text{mod } 2\tilde{x} \\
(B_k) \quad & 0 = \ell_{k-1} - \ell_k + \ell_{f-1} - 2\ell_{k-1,f-1} && \text{mod } 2\tilde{x} \text{ for } 2 \leq k \leq f-1 \\
(C_k) \quad & 0 = \ell_{f-1} - \ell_{1,k} - \ell_{k-1,f-1} && \text{mod } \tilde{x} \text{ for } 2 \leq k \leq f-1 \\
(D_{j,k}) \quad & 0 = \ell_{f-1} + \ell_{j-1,k-1} - \ell_{j,k} - \ell_{j-1,f-1} - \ell_{k-1,f-1} && \text{mod } \tilde{x} \text{ for } 2 \leq j \leq k, 2 < k < f-1.
\end{aligned}$$

For odd f , a solution of the system is given by $\ell_{j,k} = 0$ for $1 \leq j \leq k-1, 1 < k < f-1, \ell_k = 0$ for k odd, and $\ell_k = \tilde{x}$ for k even. Hence (1.1) splits if f is odd. It remains to show that there are no solutions for even $f > 2$.

For $2 \leq j < \frac{1}{2}(f-2)$, subtract the equation $(D_{i,f-j+i})$ from the equation $(D_{i,f-j+i-1})$ for $2 \leq i < j$. Add $(D_{j,f-1})$ and (C_{f-j}) , then subtract (C_{f-j+1}) . Adding the resulting equations yields

$$(E_j) \quad 0 = \ell_{f-1} - \ell_{j,f-1} - \ell_{f-j-1,f-1} \quad \text{mod } \tilde{x}.$$

Multiplying the equations (C_{f-1}) and $(E_j), 2 \leq j \leq \frac{1}{2}(f-2)$ by 2 and adding them we obtain

$$0 = (f-2)\ell_{f-1} - 2 \sum_{j=1}^{f-2} \ell_{j,f-1} \quad \text{mod } 2\tilde{x}.$$

On the other hand, adding the equations (A) and $(B_k), 1 < k < f-1$, the resulting equation is

$$\tilde{x} = (f-2)\ell_{f-1} - 2 \sum_{j=1}^{f-2} \ell_{j,f-1} \quad \text{mod } 2\tilde{x}.$$

Hence there are no solutions for f even. \square

9. PSEUDO-PROJECTIVE SPACES IN DIMENSION 4

In the final section we consider 4-dimensional pseudo-projective spaces and provide a proof of Theorem 1.5. We begin by constructing a 4-dimensional pseudo-projective space associated to given algebraic data. Namely, take $f \in \mathbb{Z}$ with $f \geq 0, x, y \in R = \mathbb{Z}[Z/f\mathbb{Z}]$ with $xy = 0$ and $f\varepsilon(x) = 0$, where ε is the augmentation of the group ring, R , so that $xR \subseteq \ker \varepsilon$. Finally, take $\gamma \in \Gamma((\ker f\varepsilon)/xR)$. Given such data, (f, x, y, α) , take a 3-dimensional pseudo-projective space $P_{f,x}$ as in (3.2). Then the set-theoretic splitting u_x of the short exact sequence

$$\Gamma(\pi_2(P_{f,x})) \longrightarrow \pi_3(P_{f,x}) \longrightarrow H_3(\widehat{P}_{f,x})$$

implies that every element of $\pi_3(P_{f,x})$ may be expressed uniquely as a sum $u_x(v) + \beta$ with $v \in H_3(\widehat{P}_{f,x})$, that is, $xv = 0$, and $\beta \in \Gamma(\pi_2(P_{f,x})) = \Gamma((\ker f\varepsilon)/xR)$, see (6.2). Using $u_x(y) + \alpha \in \pi_3(P_{f,x})$ to attach a 4-cell to $P_{f,x}$ we obtain the 4-dimensional pseudo-projective space,

$$P = P_{f,x,y,\alpha} = S_1 \cup e^2 \cup e^3 \cup e^4.$$

Note that the homotopy type of $P = P_{f,x,y,\alpha}$ is determined by (f, x, y, α) and that every 4-dimensional pseudo-projective space is of this form. The cellular chain complex, $C_*(\widehat{P})$, of the universal cover, $\widehat{P} = \widehat{P}_{f,x,y,\alpha}$, is the complex of free R -modules,

$$\langle e_4 \rangle_R \xrightarrow{d_4} \langle e_3 \rangle_R \xrightarrow{d_3} \langle e_2 \rangle_R \xrightarrow{d_2} \langle e_1 \rangle_R \xrightarrow{d_1} \langle e_0 \rangle_R,$$

given by $d_1(e_1) = e_0([\overline{1}] - [\overline{0}])$, $d_2(e_2) = e_1N$, that is, multiplication by the *norm element*, $N = \sum_{i=0}^{f-1} [\overline{i}]$, $d_3(e_3) = e_2x$, and $d_4(e_4) = e_3y$. Let $\bar{b} : R \rightarrow \pi_3 P_{f,x}$ be the homomorphism of R -modules which maps the generator $[\overline{0}] \in R$ to $\bar{b}([\overline{0}]) = u_x(y) + \alpha$, so that composition with the projection onto $H_3 \widehat{P}_{f,x}$ yields the homomorphism of R -modules induced by the boundary operator d_4 . Thus we obtain the commutative diagram

$$\begin{array}{ccccc} H_4 \widehat{P} & \xrightarrow{b} & \Gamma \pi_2 P & & \\ \downarrow & & \downarrow & \searrow j & \\ R & \xrightarrow{\bar{b}} & \pi_3 P_{f,x} & \twoheadrightarrow & \pi_3 P \\ & \searrow \bar{d}_4 & \downarrow & & \downarrow h \\ & & H_3 \widehat{P}_{f,x} & \twoheadrightarrow & H_3 \widehat{P} \end{array}$$

in the category of R -modules, where the middle column is the short exact sequence (6.1) and

$$(9.1) \quad H_4 \widehat{P} \xrightarrow{b} \Gamma \pi_2 P \xrightarrow{j} \pi_3 P \xrightarrow{h} H_3 \widehat{P}$$

is Whitehead's Certain Exact Sequence of the universal cover, $\widehat{P} = \widehat{P}_{f,x,y,\alpha}$.

Now we restrict attention to the case $f = 2$. Then $\pi_1 = \pi_1 P = \mathbb{Z}/2\mathbb{Z}$ and the augmentation ideal, K is generated by $[\overline{1}] - [\overline{0}]$. Thus

$$x = \tilde{x}([\overline{1}] - [\overline{0}]) \quad \text{and} \quad y = \tilde{y}([\overline{1}] + [\overline{0}]), \quad \text{for some } \tilde{x}, \tilde{y} \in \mathbb{Z}.$$

We assume that x and y are non-trivial, that is, $\tilde{x}, \tilde{y} \neq 0$.

Theorem 9.1. *For $P = P_{2,x,y,\alpha}$, with x and y as above, $\pi_1 P = \mathbb{Z}/2\mathbb{Z}$ acts on $\pi_2 P = \mathbb{Z}/\tilde{x}\mathbb{Z}$ via multiplication by -1 , trivially on $H_3 \widehat{P} = \mathbb{Z}/\tilde{y}\mathbb{Z}$ and via multiplication by -1 on $H_4 \widehat{P} = \mathbb{Z} = \langle [\overline{1}] - [\overline{0}] \rangle$. The exact sequence (9.1) is given by*

$$(9.2) \quad H_4 \widehat{P} = \mathbb{Z} \xrightarrow{b} \Gamma \pi_2 P = \Gamma(\mathbb{Z}/\tilde{x}\mathbb{Z}) \xrightarrow{j} \pi_3 P \xrightarrow{h} H_3 \widehat{P} = \mathbb{Z}/\tilde{y}\mathbb{Z}.$$

Denoting the generator of $\Gamma \pi_2 P$ by ξ , the boundary b is determined by

$$b([\overline{1}] - [\overline{0}]) = \tilde{x}\tilde{y}\xi,$$

and the action of $\pi_1 P$ on $\pi_3 P$ is trivial. As abelian group, $\pi_3 P$ is the extension of $H_3 \widehat{P}$ by $\text{coker } b$ given by the image of $-\alpha \in \Gamma \pi_2$ under the homomorphism

$$\tau : \Gamma \pi_2 \twoheadrightarrow \text{coker } b \twoheadrightarrow \text{coker } b / \tilde{y} \text{coker } b = \text{Ext}(\mathbb{Z}/\tilde{y}\mathbb{Z}, \text{coker } b).$$

Hence the extension $\pi_3 P$ over \mathbb{Z} determines α modulo $\ker \tau$.

Theorem 1.5 is a corollary to Theorem 9.1.

Proof. As the augmentation ideal $K \cong \mathbb{Z}$ is generated by $k = [\overline{1}] - [\overline{0}]$, the action of $\pi_1 = \mathbb{Z}/2\mathbb{Z}$ on $K = \pi_2 P_2$ and hence on $\pi_2 P = K/xR = \mathbb{Z}/\tilde{x}\mathbb{Z}$ is multiplication by -1 , since $k[\overline{1}] = -k$. But the Γ -functor maps multiplication by -1 to the identity morphism, so that π_1 acts trivially on $\Gamma(\pi_2 P)$.

As $d_3(e_3) = e_2 x$, we obtain $H_3 \widehat{P}_{2,x} \cong \mathbb{Z}$, generated by the norm element $N = [\overline{1}] + [\overline{0}]$. Since $N[\overline{1}] = N$, the action of π_1 on $H_3 \widehat{P}_{2,x}$ is trivial.

As $d_4(e_4) = e_3 y$, we obtain $H_3 \widehat{P} \cong \mathbb{Z}/\tilde{y}\mathbb{Z}$ and $H_4 \widehat{P} \cong \mathbb{Z}$, generated by $k = [\overline{1}] - [\overline{0}]$. Hence the action of π_1 on $H_4 \widehat{P}$ is multiplication by -1 .

Now let $\xi = ([\overline{1}] - [\overline{0}]) \otimes ([\overline{1}] - [\overline{0}])$ be the generator of $\Gamma(K)$. Note that $v[\overline{1}] = v$ and $\beta[\overline{1}] = \beta$, for $v \in H_3 \widehat{P}_{2,x}$ and $\beta \in \Gamma(\pi_2 P)$, since π_1 acts trivially on both $H_3 \widehat{P}_{2,x}$ and $\Gamma(\pi_2 P)$. Lemma 8.3 implies

$$(u(v) + \beta)[\overline{1}] = -\tilde{x}\tilde{y}\omega(\xi) + u(v[\overline{1}]) + \omega(\beta)[\overline{1}] = -\tilde{x}\tilde{y}\omega(\xi) + u(v) + \omega(\beta).$$

We obtain

$$\begin{aligned} \bar{b}(e_4([\overline{1}] - [\overline{0}])) &= (u(y) + \omega(\alpha))([\overline{1}] - [\overline{0}]) \\ &= -\tilde{x}\tilde{y}\omega(\xi) + u(y) + \omega(\alpha) - (u(y) + \omega(\alpha)) \\ &= -\tilde{x}\tilde{y}\omega(\xi). \end{aligned}$$

By definition of \bar{b} ,

$$\pi_3 P = \pi_3 P_{2,x} / \text{im } \bar{b}.$$

Hence π_1 acts trivially on $\pi_3(P)$.

Sequence (9.1) yields the short exact sequence

$$(9.3) \quad G = \text{coker } b \longrightarrow \pi_3 P \xrightarrow{h} H_3 \widehat{P} \cong \mathbb{Z}/\tilde{y}\mathbb{Z},$$

which represents $\pi_3 P$ as an extension of $\mathbb{Z}/\tilde{y}\mathbb{Z}$ by $G = \text{coker } b$. Thus the extension $\pi_3 P$ over \mathbb{Z} determines γ modulo the kernel of the map

$$\tau : \Gamma\pi_2 \longrightarrow \text{coker } b \longrightarrow \text{coker } b / \tilde{y}\text{coker } b = \text{Ext}(\mathbb{Z}/\tilde{y}\mathbb{Z}, \text{coker } b) .$$

□

REFERENCES

- [B] H.J. Baues, *Combinatorial Homotopy and 4-dimensional CW-complexes*, de Gruyter expositions in mathematics (1991).
- [BHS] R. Brown, P. J. Higgins and R. Sivera, *Nonabelian Algebraic Topology*, EMS Publishing House (2011).
- [O] P. Olum, *Self-equivalences and pseudo-projective planes*, *Topology* **4**, (1965), 109–127.
- [W1] J.H.C. Whitehead, *Combinatorial homotopy II*, *Bull. AMS* **55** (1949), 213–245.
- [W2] J.H.C. Whitehead, *A certain exact sequence*, *Ann. of Math.* **52** (1950), 51–110.

MAX PLANCK INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, D-53111 BONN, GERMANY
E-mail address: baues@mpim-bonn.mpg.de

SECOND AUTHOR'S HOME INSTITUTION: SCHOOL OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF NEW ENGLAND,
 NSW 2351, AUSTRALIA
E-mail address: bbleile@une.edu.au