

On the enumeration of complex plane curves with two singular points

Dmitry Kerner
Ben Gurion University, Israel

ABSTRACT. We study equi-singular strata of plane curves with two singular points of prescribed types. The method of the previous work [Kerner06] is generalized to this case. In particular we consider the enumerative problem for plane curves with two singular points of *linear* singularity types.

First the problem for two ordinary multiple points of fixed multiplicities is solved. Then the enumeration for arbitrary linear types is reduced to the case of ordinary multiple points and to the understanding of "merging" of singular points. Many examples and numerical answers are given.

CONTENTS

1. Introduction	1
2. Some relevant notions and auxiliary results	4
3. The method	10
4. Examples	16
Appendix A. Some numerical results for enumeration of curves with two singular points	22
References	23

1. INTRODUCTION

1.1. **The setup and the problem.** We work with (complex) algebraic curves in \mathbb{P}^2 . Consider the parameter space of degree- d curves, i.e. the complete linear system $|\mathcal{O}_{\mathbb{P}^2}(d)|$. It is a projective space of dimension $N_d = \binom{d+2}{2} - 1$.

A classical enumerative problem is: *given the singularity types $\mathbb{S}_1 \dots \mathbb{S}_r$, "how many" curves of the linear system $|\mathcal{O}_{\mathbb{P}^2}(d)|$ possess singular points of these types?* (To make this number finite one imposes a sufficient amount of generic base points. The degree d is assumed big enough to avoid various pathologies. By the singularity type in this paper we always mean the local, embedded, topological singularity type of a reduced plane curve. For more details and related notions cf.§2.)

In this paper we consider the case of two prescribed singular points (the case of one singular point was considered in [Kerner06]). First reformulate the problem. The parameter space $|\mathcal{O}_{\mathbb{P}^2}(d)|$ is stratified according to the singularity types of curves. The generic point of $|\mathcal{O}_{\mathbb{P}^2}(d)|$ corresponds to an (irreducible, reduced) smooth curve. The set of points corresponding to the singular curves is called the *discriminant* of plane curves. It is an irreducible projective hypersurface in $|\mathcal{O}_{\mathbb{P}^2}(d)|$.

Definition 1.1. *For the given singularity types $\mathbb{S}_1 \dots \mathbb{S}_r$, the equisingular stratum $\Sigma_{\mathbb{S}_1 \dots \mathbb{S}_r} \subset |\mathcal{O}_{\mathbb{P}^2}(d)|$ is the set of points corresponding to the curves with $\mathbb{S}_1 \dots \mathbb{S}_r$ singularities.*

The generic point of the discriminant lies in the stratum of nodal curves ($\Sigma = \overline{\Sigma}_{A_1}$). Other strata correspond to higher singularities. The stratum of δ -nodal curves $\Sigma_{\delta A_1}$ (whose closure $\overline{\Sigma}_{\delta A_1}$ contains the stratum of curves of a given genus) is the classical Severi variety. Other strata are Σ_{A_k} , Σ_{D_k} , Σ_{E_k} etc. The strata are quasi-projective varieties [Greuel-Lossen96]. For a comprehensive introduction to these equi-singular families and related notions cf. [GLS-book1],[GLS-book2].

For small d various pathologies can occur, but for sufficiently high degrees (given the singularity types $\mathbb{S}_1 \dots \mathbb{S}_r$) the strata are non-empty, irreducible, smooth, of expected (co-)dimension. One sufficient condition for this is [Dimca-book,

Date: January 13, 2011.

2000 Mathematics Subject Classification. Primary 14N10; 14N35; Secondary 14H10; 14H50;

Key words and phrases. Enumeration of singular curves, Severi-type varieties, equi-singular families of curves, Thom polynomials.

This paper is an updated version of the preprint: Max Planck Institut für Mathematik Bonn, Germany, MPIM2007-76.

The author was supported by the Max Planck Institut für Mathematik (Bonn, Germany) and the Skirball postdoctoral fellowship of the Center of Advanced Studies in Mathematics at the Mathematics Department of Ben-Gurion University (Be'er-Sheva, Israel).

§I.3]: $d \geq \sum o.d.(\mathbb{S}_i) + r - 1$, here $o.d.$ are the orders of determinacy. There are also other sufficient conditions, formulated in terms of the Milnor or Tjurina numbers of the types or the γ invariant (cf. [GLS06, §3] and [GLS-book2, §IV.2]), e.g. $\sum \gamma(\mathbb{S}_i) \leq (d+3)^2$. In this paper we always assume d big enough.

By definition each stratum is embedded into $|\mathcal{O}_{\mathbb{P}^2}(d)|$, thus its natural compactification is just the topological closure. The closures of the strata are singular (often in co-dimension 1). The closed stratum defines the homology class $[\overline{\Sigma}_{\mathbb{S}_1 \dots \mathbb{S}_r}] \in H_*(|\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z}) \approx \mathbb{Z}$ in the homology of the corresponding dimension. The degree of this class is the degree of the stratum $\deg(\overline{\Sigma}_{\mathbb{S}_1 \dots \mathbb{S}_r})$, obtained by the intersection with the generic plane in $|\mathcal{O}_{\mathbb{P}^2}(d)|$ of the complementary dimension. This corresponds to imposing generic base points. So this degree is "the number" of curves possessing the prescribed singularities.

In [Kerner06] we proposed a method to compute the degrees of strata $\overline{\Sigma}_{\mathbb{S}}$ for curves with just one singular point. In [Kerner08] the method was generalized to some singular hypersurfaces in \mathbb{P}^n . The current paper is an application of the method to the enumeration of curves with two singular points, i.e. to compute the classes $[\overline{\Sigma}_{\mathbb{S}_1 \mathbb{S}_2}]$.

Acknowledgements. This work is a tail of my PhD, done under the supervision of E.Shustin, to whom I wish to express my deepest gratitude. The conversations and advices of P.Aluffi, G.-M.Greuel, I.Tyomkin were highly important. Careful reading of two anonymous referees has improved the text significantly.

Most of the work was done during my stay in Max Planck Institut für Mathematik, Bonn in 2006-2007. Many thanks for excellent working conditions.

1.2. The known results. Since the question is completely classical, lots of *particular* results are known. We mention only a few (for a much better discussion cf. [Kleiman76, Kleiman85], [Kleiman-Piene01, Remark 3.7] and [Kazarian01]).

- The most classical case is $\deg \overline{\Sigma}_{\delta A_1}$. The history of solved cases starts from $\delta = 1$ in [Steiner1848], $\delta = 2$ in [Cayley1866] and $\delta = 3$ in [Roberts1875]. The few-nodal cases were recalculated many times by various methods, e.g. for $\delta \leq 3$ in [Harris-Pandharipande95], for $\delta \leq 5$ in [Vainsencher81, Vainsencher95], for $\delta \leq 8$ in [Kleiman-Piene98, Kleiman-Piene01, Kleiman-Piene04]. Various general algorithms have been given (e.g. [Ran89, Ran02], [Caporaso-Harris98-1, Caporaso-Harris98-2]). It was conjectured [Göttsche98] that $\deg \overline{\Sigma}_{\delta A_1}$ is a polynomial in d of degree 2δ , for $\delta < 2d - 1$. This was proved in [Liu2000], where a general method of calculation of $\deg \overline{\Sigma}_{\delta A_1}$ was also proposed. Another proof has appeared recently: [Fomin-Mikhalkin09].

Finally, some many-nodal cases were treated, as they amount to enumeration of low genus plane curves (with Gromov-Witten invariants, quantum cohomology, [Kontsevich-Manin94] and all that.) The later approach is effective for low genus curves, i.e. when the number of nodes (or higher singularities) is almost maximal ($\delta \lesssim \frac{(d-1)(d-2)}{2}$). It seems to be non-effective for high genus computations (e.g. in the case of just a few singular points).

- The $\deg \overline{\Sigma}_{A_2}$ was predicted in [Enriques32] and (re)proved in [Lascoux77, pg.151], [Vainsencher81, 8.6.3, pg.416], [di Francesco-Itzykson95, pg.86-88] and [Aluffi98, pg.10]. In the later paper $\deg \overline{\Sigma}_{A_3}$ was also obtained. In [Kleiman-Piene98, theorem 1.2] the degrees of $\overline{\Sigma}_{D_4, A_1}$, $\overline{\Sigma}_{D_4, 2A_1}$, $\overline{\Sigma}_{D_4, 3A_1}$, $\overline{\Sigma}_{D_6, A_1}$ and $\overline{\Sigma}_{E_7}$ were computed. The $\deg(\overline{\Sigma}_{A_2 A_1})$ was computed in [Vainsencher81, 8.6.4, pg.416]

- The impressive breakthrough has been recently achieved in [Kazarian01]-[Kazarian03-2]. The proposed topological method allows (in principle) to compute the degree of *any* stratum (with lots of explicit results in [Kazarian03-hab]). In particular he presents the answers $\deg \overline{\Sigma}_{\mathbb{S}_1 \dots \mathbb{S}_r}$ for the total codimension: $\sum \text{codim}(\mathbb{S}_i) \leq 7$. For higher singularities the method is not quite efficient, as it solves the problem *simultaneously* for all the singularity types of a given codimension. So, first one should classify the singularities (by now the classification seems to exist up to codimension 16 only). Even if this is done, one faces the problem of enumerating huge amount of cases (the number of types grows exponentially with the codimension). And of course, each computation can give a result for a specific choice of singularity types, it is not clear whether the method allows to obtain results for some *series* of singularities.

- In [Kerner06] the problem was solved for curves with one singular point of an *arbitrary* given singularity type. The proposed method gives immediate answer (explicit formulas) for some specific series of types (the so-called *linear*). For all other (series of) types it gives an explicit algorithm.

Despite numerous isolated results as above it seems that currently there is no effective universal method to compute $\deg \overline{\Sigma}_{\mathbb{S}_1 \dots \mathbb{S}_k}$. Not speaking about a general formula giving this degree for various choices. And most methods use the information about adjacency of types: which singularity is obtained when the singular points of the types $\mathbb{S}_1 \dots \mathbb{S}_k$ are merged (collide) generically?

1.3. Our results. We consider the equisingular strata of curves with two singular points: $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$. We restrict mostly to the case of linear singularity types (cf. definition 2.2.2). The simplest examples of linear singularity types are $x_1^p + x_2^q$ for $p \leq q \leq 2p$. (In particular they include ordinary multiple point, $A_{k \leq 3}$, $D_{k \leq 6}$, $E_{k \leq 8}$, J_{10} , $Z_{k \leq 13}$ etc.)

We construct the partial resolutions $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y} \rightarrow \overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ as on the diagram. Here Aux is an auxiliary space tracing parameters of the singularities, e.g. the singular points, the lines of the tangent cones, the line $l = \overline{xy}$. The cohomology class $[\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}] \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)|)$ is completely determined by the class $[\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}] \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux})$, so we compute the later.

$$\begin{array}{ccc} \overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y} & \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux} \\ \downarrow & & \downarrow \\ \overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y} & \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| \end{array}$$

In §4.1 the enumerative problem for two ordinary multiple points is solved, i.e. the cohomology class of $\overline{\Sigma}_{x_1^{p+1}+x_2^{p+1}, y_1^{q+1}+y_2^{q+1}}$ is computed. The enumeration is done in two ways: stepwise intersection with hypersurfaces and degeneration to reducible curves.

In §3.2.2 we use the method of degenerations from [Kerner06] to express the class $[\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}]$ (for $\mathbb{S}_x, \mathbb{S}_y$ arbitrary linear singularities) via $[\overline{\Sigma}_{x_1^{p+1}+x_2^{p+1}, y_1^{q+1}+y_2^{q+1}}]$ and some necessary classes $[\overline{\Sigma}_{\mathbb{S}}]$ (the classes of some strata of curves with one singular point of type \mathbb{S}).

The strata $\overline{\Sigma}_{\mathbb{S}}$ appear inevitably, they are irreducible components of the restriction to the diagonal: $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}$. Therefore the enumerative problem is reduced to the description of such "collision of singular points". This later problem is very complicated in general. But it is manageable when the types $\mathbb{S}_x, \mathbb{S}_y$ are *linear*. In [Kerner07-2] we gave an explicit algorithm to classify the "results of collisions", i.e. the irreducible components of $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}$. In fact the algorithm even gives the defining ideals of the corresponding strata, this is briefly recalled in §2.3.2. So, each class $[\overline{\Sigma}_{\mathbb{S}}]$ is computable by methods of [Kerner06].

So, for linear singularities, the method allows to compute the class $[\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}]$. Many numerical answers are given in the Appendix. As mentioned above, the results are valid for $d \gg 0$. In §3.3 we give a cheap sufficient bound (in terms of the orders of determinacy of $\mathbb{S}_x\mathbb{S}_y$), though the bound is far from being necessary.

For small d (but big enough so that the strata $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}, \overline{\Sigma}_{\mathbb{S}}$ are irreducible, of expected dimension) the problem is still well defined, but our formulas are inapplicable. For each such d the enumeration should be done separately, cf. [Kerner06, §5].

Our approach is most naive and classical. In some sense it is a brute-force calculation. Correspondingly it is of-ten long and cumbersome. The advantages of the method are:

- The method gives a recursive algorithm, consisting of routine parts.
- The computation does not assume any preliminary classification of singularities.
- The method seems to be more effective than other approaches (to the best of our knowledge). In particular, in Appendix we present the results for some *series* of types (as compared to *single, isolated* results previously known). Thus we can treat the question: how do the degrees of the strata depend on the invariants of the singularity types for some series of singularities?

In the case of one singular point many examples suggested in [Kerner06] that (at least for linear types) the degrees depend *algebraically* on the parameters of series (e.g. for ordinary multiple point there is a polynomial dependence on the multiplicity).

One might conjecture that in the multi-singular case the dependence will be also algebraic. Unfortunately, the simplest case already provides a counterexample: for the two ordinary multiple points of multiplicities $p+1, q+1$ the degree of the corresponding stratum depends on $\max(p, q)$ and $\min(p, q)$, cf. remark 4.4.

- As the final result we obtain the multi-degree of the (partial) resolution of the stratum $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$. (The actual degree is just a particular coefficient in a big polynomial.) This multi-degree contains many important numerical invariants, e.g. enumeration of $(\mathbb{S}_x, \mathbb{S}_y)$ with one or two singular points restricted to some curves, or with some conditions on the tangents to the branches. More generally: when the parameters of the singularity (the points, the tangents) are restricted to a subvariety of the original parameter space. So, this solves a whole class of related enumerative problems.

1.4. Contents of the paper. In §2 we fix the notations and recall some necessary notions. In particular, *linear singularities* are introduced in §2.2.2 and *collisions* of singular points are discussed in §2.3.2. In §3 we describe the method. First we recall in §3.1 the case of curves with one singular point. In §3.2 we discuss the case of two singular

points. §3.2.1 presents the direct approach: obtaining the cohomology class as the chain of intersections. §3.2.2 presents the indirect approach: degeneration to ordinary multiple points of higher multiplicities.

The method is applied to several particular cases in §4, illustrating the computation. Explicit formulas are given in Appendix, for the details of numerical computation see the attached Mathematica file [Kerner07-3].

2. SOME RELEVANT NOTIONS AND AUXILIARY RESULTS

2.1. Notations and some classical facts.

2.1.1. *Coordinates and variables.* We work with various projective spaces and their subvarieties. Adopt the following notation.

A curve is denoted by $C \subset \mathbb{P}^2$ or by the defining polynomial f , the parameter space of such curves of degree d is $|\mathcal{O}_{\mathbb{P}^2}(d)|$. Let $x \in \mathbb{P}_x^2$ be a point in the projective plane, the homogeneous coordinates are (x_0, x_1, x_2) . The generator of the cohomology ring of this \mathbb{P}_x^2 is denoted by the upper-case letter X , so that $H^*(\mathbb{P}_x^2, \mathbb{Z}) = \mathbb{Z}[X]/(X^3)$. By the same letter we denote the class of a line in the homology of \mathbb{P}_x^2 . Since it is always clear, where we speak about coordinates and where about (co)homology classes, no confusion arises. For example, consider the hypersurface

$$(1) \quad V = \{(x, y, f) \mid f(x, y) = 0\} \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|$$

Here f is a bi-homogeneous polynomial of bi-degree d_x, d_y in homogeneous coordinates $(x_0, x_1, x_2) \in \mathbb{P}_x^2, (y_0, y_1, y_2) \in \mathbb{P}_y^2$. The coefficients of f are the homogeneous coordinates on the parameter space $|\mathcal{O}_{\mathbb{P}^2}(d)|$. The cohomology class of this hypersurface is

$$(2) \quad [V] = d_x X + d_y Y + F \in H^2(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z})$$

A (projective) line through the point $x \in \mathbb{P}_x^2$ is defined by a 1-form l (so that $l \in \check{\mathbb{P}}_l^2, l(x) = 0$). Correspondingly the generator of $H^*(\check{\mathbb{P}}_l^2)$ is denoted by L .

For a variety $V \xrightarrow{i} |\mathcal{O}_{\mathbb{P}^2}(d)| \times \mathbb{P}_x^2 \times \mathbb{P}_y^2$ the pullbacks $i^*(X), i^*(Y), i^*(F)$ are constantly used. To simplify the formulas we denote the pulled back classes by the same letters (X, Y, F) . This brings no confusion as the ambient space of intersection is always specified.

We often work with symmetric p -forms $\Omega^{(p)} \in \text{Sym}^p(\check{V}_3)$, here \check{V}_3 is a 3-dimensional vector space of linear forms. Fix coordinates in \check{V}_3 , then the form is a symmetric tensor with p indices $(\Omega_{i_1, \dots, i_p}^{(p)})$. We write $\Omega^{(p)}(\underbrace{x \dots x}_k)$ as a shorthand for the tensor, multiplied k times by the point $x \in V_3$:

$$(3) \quad \Omega^{(p)}(\underbrace{x \dots x}_k) := \sum_{0 \leq i_1, \dots, i_k \leq 2} \Omega_{i_1, \dots, i_p}^{(p)} x_{i_1} \dots x_{i_k}$$

So, for example, the expression $\Omega^{(p)}(x)$ is a $(p-1)$ -form. Unless stated otherwise, we assume the symmetric form $\Omega^{(p)}$ to be generic (in particular non-degenerate, i.e. the corresponding curve $\{\Omega^{(p)}(\underbrace{x \dots x}_p) = 0\} \subset \mathbb{P}_x^2$ is smooth).

Symmetric forms occur typically as tensors of derivatives of order p in homogeneous coordinates: $f^{(p)}$. Sometimes, to emphasize the point at which the derivatives are calculated we assign it. So, e.g. $f|_x^{(p)}(\underbrace{y, \dots, y}_k)$ means a symmetric $(p-k)$ form: the tensor of derivatives of order p , calculated at the point x , and multiplied k times with y .

Sometimes we address a particular component of the tensor, e.g. $(f|_x^{(p)})_{i_1 \dots i_p} := \frac{\partial}{\partial x_{i_1}} \dots \frac{\partial}{\partial x_{i_p}} f|_x \equiv \partial_{i_1} \dots \partial_{i_p} f|_x$.

The Euler identity $f|_x^{(1)}(x) \equiv \sum x_i \partial_i f|_x = df$ and its consequences (e.g. $f|_x^{(p)}(\underbrace{x \dots x}_k) \sim f|_x^{(p-k)}$) are tacitly assumed.

2.1.2. *Blowup along the diagonal.* The diagonal $\Delta = \{x = y\} \subset \mathbb{P}_x^n \times \mathbb{P}_y^n$ appears constantly. Its class is [Fulton-book, example 8.4.2]

$$(4) \quad [\Delta] = \sum_{i=0}^n X^{n-i} Y^i \in H^{2n}(\mathbb{P}_x^n \times \mathbb{P}_y^n, \mathbb{Z})$$

For example, the proportionality condition of two symmetric forms $f^{(p)}|_x \sim g^{(p)}|_x$, for $x \in \mathbb{P}^2$, is just the coincidence of the corresponding points in a big projective space, thus its class is given by the above type formula (with $n = \binom{p+2}{2} - 1$).

The blowup of $\mathbb{P}_x^2 \times \mathbb{P}_y^2$ over the diagonal $\{x = y\}$ is described nicely as the incidence variety of triples: a line and a pair of its points.

$$(5) \quad Bl_\Delta(\mathbb{P}_x^2 \times \mathbb{P}_y^2) = \{(x, y, l) \mid x \in l \ni y\} \xrightarrow{i} \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2, \quad E_\Delta = \{x = y \in l\} \subset Bl_\Delta(\mathbb{P}_x^2 \times \mathbb{P}_y^2)$$

Lemma 2.1. • $Bl_\Delta(\mathbb{P}_x^2 \times \mathbb{P}_y^2) \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2$ is a complete intersection, with cohomology class: $[Bl_\Delta(\mathbb{P}_x^2 \times \mathbb{P}_y^2)] = (L + X)(L + Y) \in H^4(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2)$.

• The class of the exceptional divisor is: $[E_\Delta] = (L + X)(X^2 + XY + Y^2) \in H^6(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2)$ and $[E_\Delta] = X + Y - L \in H^2(Bl_\Delta(\mathbb{P}_x^2 \times \mathbb{P}_y^2))$.

The two formulas for E_Δ are related by the pushforward i_* . The identity $i_*(X + Y - L) = (L + X)(X^2 + XY + Y^2) \in H^6(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2)$ is directly verified.

Proof. Note that E_Δ is a transversal intersection of the two conditions ($x = y$ and $l(x) = 0$). Therefore: $[E_\Delta] = (L + X)(X^2 + XY + Y^2) \in H^6(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2)$.

To obtain the class of the exceptional divisor in the ring $H^*(Bl_\Delta(\mathbb{P}_x^2 \times \mathbb{P}_y^2))$ note that the hypersurface $\begin{pmatrix} x_0 & x_1 \\ y_0 & y_1 \end{pmatrix} = 0$ contains the exceptional divisor E_Δ and also the residual divisor $l_2 = 0$. ■

2.2. Some notions from singularities.

Definition 2.2. [GLS-book1, pg.202] Let $(C_x, x) \subset (\mathbb{C}_x^2, x)$ and $(C_y, y) \subset (\mathbb{C}_y^2, y)$ be two germs of isolated curve singularities. They are (locally, embedded, topologically) equivalent if there exist a local homeomorphism $(\mathbb{C}_x^2, x) \mapsto (\mathbb{C}_y^2, y)$ mapping (C_x, x) to (C_y, y) .

The corresponding equivalence class is called the (local embedded topological) singularity type. The variety of points (in the parameter space $|\mathcal{O}_{\mathbb{P}^2}(d)|$), corresponding to the curves with prescribed singularity types $\mathbb{S}_x \dots \mathbb{S}_z$ is called the equisingular stratum $\Sigma_{\mathbb{S}_x \dots \mathbb{S}_z}$.

The singularity type can be specified by a polynomial representative of the type. Several simplest types have the following representatives (all the notations are from [AGLV, §I.2], we omit the moduli of analytic classification):

$$(6) \quad \begin{aligned} A_k &: x_2^2 + x_1^{k+1}, \quad D_k : x_2^2 x_1 + x_1^{k-1}, \quad E_{6k} : x_2^3 + x_1^{3k+1}, \quad E_{6k+1} : x_2^3 + x_2 x_1^{2k+1}, \quad E_{6k+2} : x_2^3 + x_1^{3k+2} \\ J_{k \geq 1, i \geq 0} &: x_2^3 + x_2^2 x_1^k + x_1^{3k+i}, \quad Z_{6k-1} : x_2^3 x_1 + x_1^{3k-1}, \quad Z_{6k} : x_2^3 x_1 + x_2 x_1^{2k}, \quad Z_{6k+1} : x_2^3 x_1 + x_1^{3k} \\ X_{k \geq 1, i \geq 0} &: x_2^4 + x_2^3 x_1^k + x_2^2 x_1^{2k} + x_1^{4k+i}, \quad W_{12k} : x_2^4 + x_1^{4k+1}, \quad W_{12k+1} : x_2^4 + x_2 x_1^{3k+1} \end{aligned}$$

For a curve defined by $f = f_p + f_{p+1} + \dots \in \mathbb{C}\{x_1, x_2\}$ the projectivized tangent cone is the (non-reduced) scheme $\{f_p = 0\} \subset \mathbb{P}^1$. We denote it as $\mathbb{P}T_{(C,0)} = l_1^{p_1} \dots l_k^{p_k}$, where $\{l_\alpha\}$ are the distinct lines of the tangent cone and $\{p_\alpha\}$ are their multiplicities, i.e. the local multiplicities of the scheme $\{f_p = 0\}$. Thus in particular $\sum_\alpha p_\alpha = \text{mult}(C)$.

Associated to the tangent cone is the tangential decomposition: $(C, 0) = \cup(C_\alpha, 0)$. Here the tangent cone of each $(C_\alpha, 0)$ is just one line and $T_{(C_\alpha,0)} \neq T_{(C_\beta,0)}$, but the germs $(C_\alpha, 0)$ can be further reducible.

2.2.1. Newton-non-degenerate singularities. Given the representative $f = \sum a_{\mathbf{I}} \mathbf{x}^{\mathbf{I}} \in \mathbb{C}\{\mathbf{x}\}$ of the singularity type, one can draw the *Newton diagram* Γ_f of the singularity. Namely, one marks the points \mathbf{I} corresponding to non-vanishing monomials in f , and takes the convex hull of the sets $\mathbf{I} + \mathbb{R}_+^2$. The envelope of the convex hull (the chain of segments) is the Newton diagram.

Definition 2.3.

- The function is called *Newton-non-degenerate* with respect to its diagram if the truncation f_σ to every face of the diagram is non-degenerate (i.e. the truncated function has no singular points in the torus $(\mathbb{C}^*)^2$).
- The curve-germ is called *generalized Newton-non-degenerate* if it can be brought to a Newton-non-degenerate form by a locally analytic transformation.
- The singular type is called *Newton-non-degenerate* if it has a Newton-non-degenerate representative.

The Newton-non-degenerate type is completely specified by the Newton diagram of (any of) its Newton-non-degenerate representative [Oka79].

In the tangent cone of the singularity $T_C = \{l_1^{p_1} \dots l_k^{p_k}\}$, consider the lines appearing with the multiplicity 1. They correspond to smooth branches, not tangent to any other branch of the singularity.

Definition 2.4. The branches as above are called *free branches*. The tangents to the non-free branches are called *non-free tangents*.

Newton-non-degeneracy implies strong restrictions on the tangent cone: there are at most two non-free tangents.

Property 2.5. Let $T_C = \{l_1^{p_1}, \dots, l_k^{p_k}\}$ be the tangent cone of the germ $C = \cup C_j$. If the germ is generalized Newton-non-degenerate then $p_\alpha > 1$ for at most two tangents l_α .

Indeed, suppose $(C, 0)$ is Newton-non-degenerate in some coordinates, let $f = f_p + f_{p+1} + \dots$ be its locally defining function. The non-degeneracy means that f_p has no singular points in \mathbb{C}^* , hence the statement.

2.2.2. *Linear singularities.* By definition the equisingular stratum $\overline{\Sigma}_{\mathbb{S}}$ is a subvariety of $|\mathcal{O}_{\mathbb{P}^2}(d)|$. Let the tangent cone be $T_{(C,0)} = l_1^{p_1} \dots l_k^{p_k}$ where $l_1 \dots l_k$ are (distinct) lines passing through x . Consider the subvariety

$$(7) \quad \overline{\Sigma}_{\mathbb{S}} \supset \overline{\Sigma}_{\mathbb{S}|x, \{l_i\}} := \overline{\{\text{curves of degree } d \text{ with } \mathbb{S} \text{ at } x \in \mathbb{P}^2 \text{ and } T_{(C,x)} = l_1^{p_1} \dots l_k^{p_k}\}}$$

Definition 2.6. The singularity (the germ, the stratum, the type) is called linear if $\overline{\Sigma}_{\mathbb{S}|x, \{l_i\}}$ is a linear subspace of $|\mathcal{O}_{\mathbb{P}^2}(d)|$ for some (and hence for any) choice of $x, \{l_i\}$.

For example, an ordinary multiple point is linear. For a linear singularity type \mathbb{S} an appropriate modification of the equisingular stratum $\overline{\Sigma}_{\mathbb{S}} \rightarrow \overline{\Sigma}_{\mathbb{S}}$ (defined later) fibres over the auxiliary space of lines through the points $\text{Aux} = \{(x, l_1 \dots l_k) \mid \forall i : x \in l_i\}$.

Linear types are abundant by the following observation. Let $(C, 0) = \cup_\alpha (C_\alpha, 0)$ be the *tangential* decomposition as above. Let \mathbb{S}_α be the singularity type of $(C_\alpha, 0)$, note that the types $\mathbb{S}_1 \dots \mathbb{S}_k$ are completely determined by \mathbb{S} . For each non-smooth $(C_\alpha, 0)$ choose the coordinates: one axis is tangent to the branch, the other axis is generic. Assume the Newton diagram $\Gamma_{(C_\alpha, 0)}$ is commode, i.e. intersects all the coordinate axes, and $(C_\alpha, 0)$ is non-degenerate in the chosen coordinates. For each face σ of the Newton diagram let a_σ be the angle between the face and the coordinate axis \hat{x}_1 .

Proposition 2.7. [Kerner06, section 3.1] Under the assumptions as above the type \mathbb{S} is linear iff each \mathbb{S}_α is linear. And \mathbb{S}_α is linear iff every face σ of the Newton diagram $\Gamma_{(C_\alpha, 0)}$ has a bounded slope: $\frac{1}{2} \leq |tg(a_\sigma)| \leq 2$.

Example 2.8. The simplest class of examples of linear singularities is defined by the series: $f = x^p + y^q$, $p \leq q \leq 2p$. In general, for a given series only for a few types of singularities the strata can be linear. In the low modality cases the linear types are:

- Simple singularities (no moduli): $A_{1 \leq k \leq 3}$, $D_{4 \leq k \leq 6}$, $E_{6 \leq k \leq 8}$
- Unimodal singularities: $X_9 (= X_{1,0})$, $J_{10} (= J_{2,0})$, $Z_{11 \leq k \leq 13}$, $W_{12 \leq k \leq 13}$
- Bimodal: $Z_{1,0}$, $W_{1,0}$, $W_{1,1}$, W_{17} , W_{18}

Most singularity types are nonlinear. For example, if a curve has an A_4 point, then the choice of one axis as the generic tangent brings it to the Newton diagram of A_3 , but with the defining function: $(\alpha x_2 + \beta x_1^2)^2 + \gamma x_1^5 + \dots$. The stratum of curves whose local defining equation begins with expansion of this type (i.e. α, β are not fixed) is not a linear subspace in $|\mathcal{O}_{\mathbb{P}^2}(d)|$.

2.2.3. *Finite determinacy.* The finite determinacy theorem of Tougeron (cf. [AGLV, §I.1.5] or [GLS-book1, §I.2.2]) states that the topological type of the curve-germ is fixed by a finite jet of the defining series. Namely, for every type \mathbb{S} , there exists k such that for all bigger $n \geq k$: $(\{\text{jet}_n(f) = 0\} \text{ has type } \mathbb{S})$ implies $(\{f = 0\} \text{ has type } \mathbb{S})$. The minimal such k is called: *the order of determinacy* (for contact equivalence). E.g. $o.d.(A_k) = k + 1$, $o.d.(D_k) = k - 1$. The classical theorem [GLS-book1, thm I.2.23] reads: if $m^{k+1} \subset m^2 \text{Jac}(f)$ then $o.d.(f) \leq k$.

2.3. Equisingular strata and related questions.

2.3.1. *Resolution of the singularities of closed strata.*

The following situation occurs frequently. Let Aux be a smooth, irreducible projective variety $\widetilde{\Sigma} \subset \overline{\Sigma} \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|$ and the projection $\widetilde{\Sigma} \xrightarrow{\pi} \text{Aux}$ a locally trivial fibration over the base $\pi(\widetilde{\Sigma}) \subset \text{Aux}$, such that $\overline{\pi(\widetilde{\Sigma})} = \text{Aux}$ and the fibres are \mathbb{P}^n linearly embedded into $|\mathcal{O}_{\mathbb{P}^2}(d)|$. In particular $\widetilde{\Sigma}$ is irreducible and smooth.

We want to compactify $\widetilde{\Sigma}$ in a smooth way, preserving the bundle structure. Start from the topological closure $\overline{\widetilde{\Sigma}} \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|$. Note that the projection $\overline{\widetilde{\Sigma}} \rightarrow \text{Aux}$ is well defined (being the restriction of π) and is surjective.

Proposition 2.9. There exists a birational morphism of smooth varieties $\widetilde{\text{Aux}} \xrightarrow{\phi} \text{Aux}$ with the properties:

- It is an isomorphism over $\pi(\widetilde{\Sigma})$ and a finite collection of blowups with smooth centers in $\text{Aux} \setminus \pi(\widetilde{\Sigma})$
- The corresponding lifting $\widetilde{\Sigma} \subset \widetilde{\text{Aux}} \times |\mathcal{O}_{\mathbb{P}^2}(d)|$ is a projective fibration over $\widetilde{\text{Aux}}$, with fibres \mathbb{P}^n linearly embedded into $|\mathcal{O}_{\mathbb{P}^2}(d)|$.

In particular, $\widetilde{\Sigma}'$ is smooth.

Proof. As the fibres of the projection $\widetilde{\Sigma} \rightarrow \pi(\widetilde{\Sigma}) \subset \text{Aux}$ are linear subspaces of $|\mathcal{O}_{\mathbb{P}^2}(d)|$ we have a natural morphism: $\pi(\widetilde{\Sigma}) \rightarrow \text{Gr}(\mathbb{P}^n, |\mathcal{O}_{\mathbb{P}^2}(d)|)$. Hence there is a rational map: $\text{Aux} \dashrightarrow \text{Gr}(\mathbb{P}^n, |\mathcal{O}_{\mathbb{P}^2}(d)|)$, whose indeterminacy locus lies in $\text{Aux} \setminus \pi(\widetilde{\Sigma})$.

Resolve this map by a chain of blowups $\widetilde{\text{Aux}} \xrightarrow{\phi} \text{Aux}$ (note that Aux is itself smooth), then get the diagram on the right.

Now, the construction ensures that the projection $\widetilde{\Sigma}' \rightarrow \widetilde{\text{Aux}}$ is a locally trivial fibration, whose fibres are linear subspaces of $|\mathcal{O}_{\mathbb{P}^2}(d)|$.

$$\begin{array}{ccc} \widetilde{\text{Aux}} \times |\mathcal{O}_{\mathbb{P}^2}(d)| \supset \widetilde{\Sigma}' & \rightarrow & \widetilde{\Sigma} \supset \widetilde{\Sigma} \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)| \\ \downarrow & & \searrow \downarrow \\ \widetilde{\text{Aux}} & \xrightarrow{\phi} & \text{Aux} \supset \pi(\widetilde{\Sigma}) \\ \downarrow & & \swarrow \downarrow \\ \text{Gr}(\mathbb{P}^n, |\mathcal{O}_{\mathbb{P}^2}(d)|) & & \end{array}$$

This follows from the pull-back of the universal family on the Grassmanian. In more detail, let $pt \in \widetilde{\text{Aux}} \setminus \pi(\widetilde{\Sigma})$. Let $(D, pt) \subset \widetilde{\text{Aux}}$ be the germ of a smooth curve such that $D \setminus \{pt\} \subset \pi(\widetilde{\Sigma})$. The fibration over $D \setminus \{pt\}$ is locally trivial and extends (e.g. by the topological closure in the Grassmanian) to a locally trivial fibration over D .

Hence the fibre of $\widetilde{\Sigma}' \rightarrow \widetilde{\text{Aux}}$ over pt contains a linear subspace $L_{pt,D} \approx \mathbb{P}^n \subset |\mathcal{O}_{\mathbb{P}^2}(d)|$. But, this linear subspace does not depend on the curve D , it is determined just by the point of the Grassmanian, the image of $\{pt\}$. Hence, by taking all the possible curves at $pt \in \widetilde{\text{Aux}}$ we get: all the fibres are linear subspace of $|\mathcal{O}_{\mathbb{P}^2}(d)|$ of constant dimension. Hence the statement. ■

Note that the theorem is completely general. In practice it is quite hard to realize the resolved base space $\widetilde{\text{Aux}}$ as some nice variety with a simple presentation of the (co)homology ring, explicit generators, etc.

2.3.2. Flat limits and the decomposition of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ into irreducible components. The following question occurs frequently. Given the singularity types $\mathbb{S}_x \mathbb{S}_y$ find the defining equations of the equisingular stratum $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}(x, y, \dots)$ near the diagonal $x = y$. And then describe the decomposition of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ into irreducible components and find their defining ideals and multiplicities. We call this process: collision.

In general this collision/adjacency problem is very complicated. It is manageable in the case of linear singularity types, because the equations of $\widetilde{\Sigma}_{\mathbb{S}_x}(x, \dots)$, $\widetilde{\Sigma}_{\mathbb{S}_y}(y, \dots)$ are known (and hence the equations of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ away from the diagonal). To obtain the needed set of equations one expands $y = x + \epsilon v$ where $v \in l = \overline{xy}$, and considers the ideal $\langle I_{\mathbb{S}_x}(x, \dots), I_{\mathbb{S}_y}(x + \epsilon v, \dots) \rangle$. Now one should take the flat limit as $\epsilon \rightarrow 0$, a well known operation in commutative algebra (cf. for example [Eisenbud-book, chapter XV]).

From the computational point of view one does the following. Start from the ideal $\langle I_{\mathbb{S}_x}(x, \dots), I_{\mathbb{S}_y}(x + \epsilon v, \dots) \rangle$, where in $I_{\mathbb{S}_y}(x + \epsilon v, \dots)$ all the equations are series in ϵ . If for a member of this ideal the expansion in ϵ has no "constant" term, of 0'th order in ϵ , divide it by the maximal possible power of ϵ and add this normalized version to the ideal. After several such iterations the process stabilizes, i.e. no non-trivial syzygies remain. So, we have obtained the defining ideal of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ near the diagonal. Then the ideal of the restriction $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ is obtained by putting $\epsilon = 0$ in all the equations, i.e. constructing $I \otimes \mathcal{O}_{x=y} = I/(\epsilon)$.

This process is described in details in [Kerner07-2, §3.1] where many examples are considered.

2.3.3. Collision with an ordinary multiple point is the basic and most important case. Let \mathbb{S}_y denote the singularity type of an ordinary multiple point. Assume $\text{mult}(\mathbb{S}_x) = p + 1 \geq \text{mult}(\mathbb{S}_y) = q + 1$ and the collision is generic, i.e. the curve $l = \overline{xy}$ is not tangent to any of the non-free branches of \mathbb{S}_x , see definition 2.4.

We should translate the conditions at the point y to conditions at x . Outside the diagonal $x = y$ the stratum is defined by the set of conditions corresponding to $\widetilde{\Sigma}_{\mathbb{S}_x}$, and by the condition $f|_y^{(q)} = 0$. The later is the (symmetric) form of derivatives of order q , calculated at the point y (in homogeneous coordinates). In the neighborhood of x expand $y = x + \epsilon v$, here ϵ is small and v is the direction along the line $l = \overline{xy}$.

To take the flat limit, expand $f|_y^{(q)}$ around x , we get $0 = f|_y^{(q)} = f|_x^{(q)} + \dots + \frac{\epsilon^{p-q}}{(p-q)!} f|_x^{(p)}(\underbrace{v \dots v}_{p-q}) + \dots$. First several terms in the expansion vanish, up to the multiplicity of \mathbb{S}_x . Normalize by the common factor of ϵ to get the power series:

$$(8) \quad \frac{1}{(p-q+1)!} f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1-q}) + \frac{\epsilon}{(p-q+2)!} f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2-q}) + \frac{\epsilon^2}{(p-q+3)!} f|_x^{(p+3)}(\underbrace{v \dots v}_{p+3-q}) + \dots$$

There are $\binom{q+2}{2}$ series here. To take the flat limit, we should find all the syzygies between these series and the equations for $\widetilde{\Sigma}_{\mathbb{S}_x}$. First we find the "internal" syzygies among the series themselves.

Lemma 2.10. • *The standard basis, obtained by considering all the syzygies of the equation (8), is (with numerical coefficients omitted):*

$$(9) \quad \begin{array}{cccccccc} f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1-q}) + \epsilon f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2-q}) + \epsilon^2 f|_x^{(p+3)}(\underbrace{v \dots v}_{p+3-q}) + \epsilon^3 f|_x^{(p+4)}(\underbrace{v \dots v}_{p+4-q}) + \dots \\ 0 & + & f|_x^{(p+2)}(\underbrace{v \dots v}_{p+3-q}) + \epsilon f|_x^{(p+3)}(\underbrace{v \dots v}_{p+4-q}) + \epsilon^2 f|_x^{(p+4)}(\underbrace{v \dots v}_{p+5-q}) + \dots \\ 0 & + & 0 & + & f|_x^{(p+3)}(\underbrace{v \dots v}_{p+5-q}) + \epsilon f|_x^{(p+4)}(\underbrace{v \dots v}_{p+6-q}) + \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & + & 0 & + & 0 & + & \dots & + f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1}) + \dots \end{array}$$

• *If the singularity \mathbb{S}_x is an ordinary multiple point $(x_1^{p+1} + x_2^{p+1})$, with $p \geq q$ then the series as above define the stratum $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}(x, y, l) \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|$ near the diagonal $\{x = y\}$. Here $\text{Aux} = \{(x, y, l) | x \in l \ni y\} \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \mathbb{P}_l^2$. In particular, the restriction $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}(x, y, l)|_{x=y}$ is an irreducible variety, whose generic point corresponds to the singularity type of $(x_1^{p-q} + x_2^{p-q})(x_1^{q+1} + x_2^{2q+2})$ with l -the tangent line to $(x_1^{q+1} + x_2^{2q+2})$.*

Note that all the series are linear in function/its derivatives in accordance with proposition 2.9. In several simplest cases we have: $A_1 + A_1 \rightarrow A_3$, $D_4 + A_1 \rightarrow D_6$, $X_9 + A_1 \rightarrow X_{1,2}$, $D_4 + D_4 \rightarrow J_{10}$, $X_9 + D_4 \rightarrow Z_{13}$.

Proof. • The syzygies are obtained as a consequence of the Euler identity for homogeneous polynomial $\sum x_i \partial_i f = \deg(f) f$. By successive contraction of the series of equation (8) and omitting $f^{(j)}|_x$ for $j \leq p$ with x , we get the series

$$(10) \quad \begin{array}{ccc} \frac{1}{(p-q+1)!} f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1-q}) + \frac{\epsilon}{(p-q+2)!} f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2-q}) + \frac{\epsilon^2}{(p-q+3)!} f|_x^{(p+3)}(\underbrace{v \dots v}_{p+3-q}) + \dots \\ \frac{(d-p-2)}{(p-q+2)!} f|_x^{(p+1)}(\underbrace{v \dots v}_{p+2-q}) + \frac{\epsilon(d-p-3)}{(p-q+3)!} f|_x^{(p+2)}(\underbrace{v \dots v}_{p+3-q}) + \frac{\epsilon^2(d-p-4)}{(p-q+4)!} f|_x^{(p+3)}(\underbrace{v \dots v}_{p+4-q}) + \dots \\ \dots & \dots & \dots \\ \frac{\prod_{i=2}^{q+1} (d-p-i)}{(p+1)!} f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) + \frac{\epsilon \prod_{i=2}^{q+1} (d-p-1-i)}{(p+2)!} f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2}) + \frac{\epsilon^2 \prod_{i=2}^{q+1} (d-p-2-i)}{(p+3)!} f|_x^{(p+3)}(\underbrace{v \dots v}_{p+3}) + \dots \end{array}$$

Here the first row is the initial series, the second is obtained by contraction with x once, the $p+2$ 'th row is obtained by contracting $(p+1)$ times with x .

Apply now the Gaussian elimination, to bring this system to the upper triangular form.

★ Eliminate from the first column all the entries of the rows $2 \dots (p+2)$. For this contract the first row necessary number of times with v (fix the numerical coefficient) and subtract.

★ Eliminate from the second column all the entries of the rows $3 \dots (p+2)$.

★ ...

Normalize the rows (i.e. divide by the necessary power of ϵ).

In this way we get the "upper triangular" system of series in eq. (9) (we omit the numerical coefficients). It remains to check that there are no more "internal" syzygies. This is proved in the second part.

• Suppose \mathbb{S}_x is an ordinary multiple point. First we should check that the upper triangular system, obtained above, has no more syzygies. Alternatively, that it generates a prime ideal. This can be done directly (by methods of commutative algebra). However in our case a simpler way is to check the dimension of the restriction $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ and to compare it to the dimension obtained from (9).

Note that $\dim(\text{Aux}) = 4$ and $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ is irreducible. Hence the fibre over the generic point (x, y, l) has dimension $\dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}) - 4$. For the diagonal: $\dim\{(x, y, l) | x = y \in l\} = 3$. Therefore

$$(11) \quad \dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}) \geq (\dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}) - 4) + 3 = \dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}) - 1$$


On the other hand, by the irreducibility of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ one has: $\dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}) \leq \dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}) - 1$. Thus we get the dimension of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ and of its fibre over the generic point $\{x = y \in l\}$.

Now check the scheme defined by equation (9). Take the limit $\epsilon \rightarrow 0$ (i.e. omit the higher order terms in each row):

$$(12) \quad f|_x^{(p)} = 0, \quad f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1-q}) = 0, \quad f|_x^{(p+2)}(\underbrace{v \dots v}_{p+3-q}) = 0, \quad f|_x^{(p+3)}(\underbrace{v \dots v}_{p+5-q}) = 0 \quad \dots, \quad f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1}) = 0$$

Observe that the system is linear in f . For a fixed x, v , e.g. $x = (0, 0, 1), v = (0, 1, 0)$, each equation is vanishing of some directional derivatives, and they are linearly independent. Hence the fibres over (x, v) are linear spaces of constant dimension and the variety is irreducible.

As each condition means the absence of some monomials in the expansion of f , the Newton diagram is easily constructed. Now the co-dimension of the fibre over $\{x = y \in l\}$ can be computed e.g. as the number of $\mathbb{Z}_{\geq 0}^2$ points strictly below the Newton diagram.



For the dimension one gets precisely $\dim(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}) - 4$, i.e. the expected dimension. This proves that the system in (9) defines an irreducible variety: the stratum $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ near the diagonal.

Finally, from the last set of equations or from the Newton diagram we get: the restriction $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ is irreducible. It is the equisingular stratum $\widetilde{\Sigma}_{\mathbb{S}}$ for the singularity type obtained from the Newton diagram. Since all the slopes of the diagram lie in the segment $[\frac{1}{2}, 2]$ the singularity type is linear.

For some fixed x, l the generic point of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ corresponds to the generic function with the given Newton diagram, i.e. (with some numerical coefficients) $f = x_1^{p+1} + \dots + x_1^{q+1}x_2^{p-q} + \dots + x_2^{p+q+2} + \text{higher order terms}$. By genericity f is non-degenerate with respect to its diagram, hence the singularity is the union of $(p-q)$ non-tangent smooth branches and $(q+1)$ smooth branches with pairwise tangency 2, i.e. has the singularity type of $(x_1^{p-q} + x_2^{p-q})(x_1^{q+1} + x_2^{2q+2})$. ■

In the more general case, when \mathbb{S}_x is not an ordinary multiple point, the generators of $I(\widetilde{\Sigma}_{\mathbb{S}_x})$ should be added and one checks again for the possible syzygies.

2.3.4. Tangency of the degenerating hypersurface and the diagonal. Each step of the method is the intersection $\widetilde{\Sigma}_1 \cap V = \widetilde{\Sigma}_2 \cup R \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$. Here the ideal $I(\widetilde{\Sigma}_1)$ is assumed to be known, V is a hypersurface (with the known equation), R is usually non-reduced, the multiplicities of its components are the degrees of tangency of V and $\widetilde{\Sigma}_1$ near the diagonal.

The multiplicities of R are computed in a standard way. The primary decomposition of the ideal $\langle I(\widetilde{\Sigma}_1), I(V) \rangle$ consists of $I(\widetilde{\Sigma}_2)$ and some ideals of schemes over the diagonal $x = y$. These ideals give the multiplicities.

As an example, consider the degeneration of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ by increasing the local degree of intersection of the curve and the line $l = \bar{x}$ at $x \in \mathbb{P}_x^2$. For the generic curve in the original stratum this degree is just the multiplicity of \mathbb{S}_x : $(p+1)$. Hence the degenerating hypersurface is $V = \{(f|_x^{(p+1)})(\underbrace{v, \dots, v}_{p+1}) = 0\}$. We assume that v does not belong to the lines

of the tangent cone at x , i.e. $l \neq l_x$. Let $\widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y}$ be the degenerated stratum.

Lemma 2.11. *Suppose $\mathbb{S}_x, \mathbb{S}_y$ are ordinary multiple points of multiplicities $p+1 \geq q+1$. The multiplicity of the residual piece is $q+1$, i.e. $[\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}] \times [V] = [\widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y}] + (q+1)[R_{\text{reduced}}]$*

Proof. The locally defining (prime) ideal of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ near the diagonal $x = y$ is obtained in §2.3.3:

$$(13) \quad f|_x^{(p)} = 0, \quad f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1-q}) + \dots = 0, \quad f|_x^{(p+2)}(\underbrace{v \dots v}_{p+3-q}) + \dots = 0, \quad f|_x^{(p+3)}(\underbrace{v \dots v}_{p+5-q}) + \dots = 0 \quad \dots, \quad f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+1+q}) + \dots = 0$$

Add the hypersurface equation $\{f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) = 0\}$ to this system of generators, decompose the so obtained ideal into primary components and find the resulting multiplicity. So, contract the first series q times with v , the second series $(q-1)$ times with v etc.:

$$(14) \quad \begin{aligned} & f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) + \epsilon^{q+1} f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) \dots = 0, \quad f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2}) + \epsilon^{p+q+2} f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) \dots = 0, \\ & f|_x^{(p+3)}(\underbrace{v \dots v}_{p+3}) + \epsilon^{p+q+2} f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) \dots = 0 \quad \dots, \quad f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1}) + \dots = 0 \end{aligned}$$

Together with $f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) = 0$ the first equation becomes: $\epsilon^{q+1} f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) \dots = 0$.

So the ideal has two components: $\langle \epsilon^{q+1}, f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}), f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2}), \dots, f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1}) \rangle$ and

$$(15) \quad \begin{aligned} & \langle f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}), f|_x^{(p+2)}(\underbrace{v \dots v}_{p+2}) + \epsilon^{p+q+2} f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) \dots, \\ & f|_x^{(p+3)}(\underbrace{v \dots v}_{p+3}) + \epsilon^{p+q+2} f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) \dots \dots, f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1}) + \dots, f|_x^{(p+q+2)}(\underbrace{v \dots v}_{p+q+2}) + \dots \rangle \end{aligned}$$

(In addition to these, the ideal contains also the equations of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ near the diagonal.)

The first component is supported on the diagonal and is of multiplicity $q + 1$, corresponding to the residual piece. The second corresponds to $\widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y}$. ■

3. THE METHOD

First we recall the case of curves with one singular point and state the necessary results. Then we consider the case of two singular points and discuss additional complications.

Recall some conventions.

- We work with many liftings, i.e. embeddings of Σ_{**} into some multi-projective spaces. To avoid messy notations (like $\widetilde{\Sigma}$) we denote by $\widetilde{\Sigma}$ any lifting. This causes no confusion, as we always specify the embedding (e.g. $\widetilde{\Sigma} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$) or the associated parameters, e.g. $\widetilde{\Sigma}(x, y, l, \dots)$.
- We always work with integral cohomology $H^*(\mathbb{P}^n) = H^*(\mathbb{P}^n, \mathbb{Z}) = \mathbb{Z}[t]/t^{n+1}$ or with integral homology.
- As was said above, once the cohomology class $[\widetilde{\Sigma}_{\mathbb{S}}] \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}, \mathbb{Z})$ is computed, the needed degree is obtained by Gysin homomorphism, corresponding to the projection $\text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)| \rightarrow |\mathcal{O}_{\mathbb{P}^2}(d)|$ (i.e. extraction of a particular coefficient). If the projection $\widetilde{\Sigma}_{\mathbb{S}} \rightarrow \Sigma_{\mathbb{S}}$ is a covering, one should also divide by the order of the symmetry group of branches. Therefore, in the following we are interested in the cohomology classes of the *lifted* strata $[\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}]$.
- Below we consider mostly linear singularities, for the definition and properties cf. §2.2.2.

3.1. Equisingular strata of curves with one singular point. Here we summarize briefly some results of [Kerner06]. The equisingular strata are resolved by lifting to a bigger ambient space. The lifted strata can be defined by some simple explicit conditions.

3.1.1. The case of an ordinary multiple point. The ordinary multiple point ($\mathbb{S} = x_1^{p+1} + x_2^{p+1}$) is the simplest case (cf. [Kerner06, §2.3]). The first lifting (tracing the singular point) is already a *smooth, globally complete* intersection:

$$(16) \quad \widetilde{\Sigma}_{\mathbb{S}}(x) = \{(x, f) \mid f|_x^{(p)} = 0\} \subset \mathbb{P}_x^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|$$

Here $f|_x^{(p)}$ is the tensor of all the partial derivatives of order p in homogeneous coordinates, calculated at the point x . This precisely encodes vanishing of the function and of all the derivatives up to order p in local coordinates, i.e. the conditions of an ordinary multiple point. The transversality of the defining conditions can be easily seen. For example, fix some point x , then the conditions are linear, so the transversality is equivalent to linear independence.

Thus in this case the (co)homology class $[\widetilde{\Sigma}_{\mathbb{S}}(x)]$ is just the product of the (co)homology classes of the defining hypersurfaces:

$$(17) \quad [\widetilde{\Sigma}_{\mathbb{S}}(x)] = \left(F + (d-p)X \right)^{\binom{p+2}{2}} \in H^*(\mathbb{P}_x^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z})$$

The class of $[\widetilde{\Sigma}_{\mathbb{S}}]$ is obtained by Gysin projection, which in this case means: to extract the coefficient of X^2 . Hence $[\widetilde{\Sigma}_{\mathbb{S}}] = \binom{p+2}{2} (d-p)^2 F^{\binom{p+2}{2}-2} \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z})$.

3.1.2. The general linear singularity. In general, let \mathbb{S} be a *linear* singularity type with the tangent cone $T_{\mathbb{S}} = l_1^{p_1} \dots l_k^{p_k}$. Here $p = \sum p_i$ is the multiplicity. Consider the corresponding incidence variety, i.e. lift $\widetilde{\Sigma}_{\mathbb{S}}$ to a bigger ambient space:

$$(18) \quad \widetilde{\Sigma}_{\mathbb{S}}(x, l_1 \dots l_k) := \overline{\{(x, l_1 \dots l_k, f) \mid C = f^{-1}(0) \subset \mathbb{P}_x^2 \text{ has the singularity of type } \mathbb{S} \text{ at } x \text{ with } T_{(C,x)} = l_1^{p_1} \dots l_k^{p_k}\}}$$

The embeddings and projections of the varieties are given on the diagram. Here

$$(19) \quad \begin{array}{ccccc} \widetilde{\Sigma}_{\mathbb{S}} \subset \widetilde{\Sigma}_{\mathbb{S}} \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux} & & & \\ \downarrow & \downarrow & & & \downarrow \\ \Sigma_{\mathbb{S}} \subset \widetilde{\Sigma}_{\mathbb{S}} \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| & & & \end{array}$$

is the auxiliary incidence variety of lines through the points.

Assume that the parts of the tangential decomposition of \mathbb{S} (cf. §2.2) have distinct singularity types and are ordered.

Then for a given $(C, 0)$ the point $(x, l_1 \dots l_k, C) \in \widetilde{\Sigma}_{\mathbb{S}}$ is unique. Then the projection $\widetilde{\Sigma}_{\mathbb{S}} \rightarrow \Sigma_{\mathbb{S}}$ is an isomorphism and hence $\widetilde{\Sigma}_{\mathbb{S}} \rightarrow \overline{\Sigma}_{\mathbb{S}}$ is birational. If the singularity types for some of the parts coincide (e.g. for an ordinary multiple point) then $\widetilde{\Sigma}_{\mathbb{S}} \rightarrow \Sigma_{\mathbb{S}}$ is an un-ramified covering.

As the type \mathbb{S} is linear, the fibres of the projection $\widetilde{\Sigma}_{\mathbb{S}}(x, l_1 \dots l_k) \rightarrow \text{Aux}$ are linear subspaces of $|\mathcal{O}_{\mathbb{P}^2}(d)|$. The cohomology class $[\widetilde{\Sigma}_{\mathbb{S}}] \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}, \mathbb{Z})$ can be computed as follows.

Let $\{f = 0\} = (C, x) \subset (\mathbb{C}^2, 0)$ be the (generic) curve-germ of the type \mathbb{S} . Several initial jets of f are divisible by the one-form l_i :

$$(20) \quad \text{jet}_{p-1}(f)|_x = 0, \text{jet}_p(f)|_x = l_i^{p_i} \times (\dots), \text{jet}_{p+1}(f)|_x = l_i^{p_i - a_i} \times (\dots), \dots, \text{jet}_{p+n_i}(f)|_x = l_i \times (\dots),$$

To obtain this, choose the tangent line l_i as one of the coordinate axes, the other axis is chosen generically.

These conditions can be written globally in Aux as:

$$(21) \quad f^{(p-1)}|_x = 0, f^{(p)}|_x \sim \text{SYM}(l_i^{p_i}, A_{p-p_i}), \dots, f^{(p+n_i)}|_x \sim \text{SYM}(l_i^{q_i}, A_{p+n_i-1})$$

Here A_j are some auxiliary symmetric tensors (of rank j) and SYM means the symmetrization. Do the same for all the tangent lines to get the collection of conditions:

$$(22) \quad f^{(p-1)}|_x = 0, f^{(p)}|_x \sim \text{SYM}(l_1^{p_1}, \dots, l_k^{p_k}), \dots, f^{(p+n)}|_x \sim \text{SYM}(l_1^{n_1}, \dots, l_k^{n_k}, A_{p+n-n_1-\dots-n_k})$$

Now use the Euler identity $f^{(N)}|_x(x) \sim f^{(N-1)}|_x$ and its consequences to present these conditions in the form:

$$(23) \quad f^{(p+n)}|_x \sim \text{SYM}(l_1^{n_1}, \dots, l_k^{n_k}, A_{p+n-n_1-\dots-n_k}), A_{p+n-n_1-\dots-n_k}(x) \sim \text{SYM}(l_1^{m_1}, \dots, l_k^{m_k}, A_{p+n-\sum n_i-\sum m_i}), \dots, A_q(x) = 0$$

Now observe that distinct proportionality conditions are obviously mutually transversal. For example f appears in the first proportionality only, $A_{p+n-n_1-\dots-n_k}$ only in the first and in the second etc.

Theorem 3.1. [Kerner06, §3] *Let $\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\}, \{A_j\})$ be the lifted stratum, defined by the equation (23). Then $\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\}, \{A_j\}) \rightarrow \text{Aux}$ is the (smooth) projective bundle over the auxiliary incidence variety Aux of points, lines $\{l_i\}$ and symmetric tensors $\{A_{**}\}$. The fibres are linear subspaces of $|\mathcal{O}_{\mathbb{P}^2}(d)|$. The cohomology class of the lifted stratum is the product of the cohomology classes*

$$(24) \quad [\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\}, \{A_j\})] = \left[f^{(p+n)}|_x \sim \text{SYM}(l_1^{n_1}, \dots, l_k^{n_k}, A_{p+n-n_1-\dots-n_k}) \right] \times \left[A_{p+n-n_1-\dots-n_k}(x) \sim \text{SYM}(l_1^{m_1}, \dots, l_k^{m_k}, A_{p+n-\sum n_i-\sum m_i}) \right] \times \dots \times [A_q(x) = 0]$$

The projection $\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\}, \{A_j\}) \rightarrow \widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\}) \rightarrow \Sigma_{\mathbb{S}}$ is isomorphism (or unramified cover if some tangential components are equisingular).

Note that each of the conditions is proportionality of two tensors, i.e. coincidence of two points in some big projective space. Correspondingly its cohomology class is readily written, from §2.1.2.

Example 3.2. Let \mathbb{S} be the type of k pairwise non-tangent branches, each of the cuspidal type $x_1^{p_\alpha} + x_2^{p_\alpha+1}$. For example, a representative of this type can be chosen as: $f = \prod l_\alpha^{p_\alpha} + f_{p+1}$ where f_{p+1} is generic, and $\{l_\alpha\}$ are some distinct linear forms. Then $T_{\mathbb{S}} = l_1^{p_1} \dots l_k^{p_k}$ and any other germ $g = g_p + g_{p+1} + \dots$ with the same tangent cone and generic g_{p+1} is of this type. Hence the lifted stratum is

$$(25) \quad \begin{aligned} \widetilde{\Sigma}_{\mathbb{S}}(x, l_1 \dots l_k) &= \left\{ (x, l_1 \dots l_k, f) \mid f^{(p)} \sim \text{SYM}(l_1^{p_1} \dots l_k^{p_k}), \forall \alpha \ x \in l_\alpha \right\} \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)| \\ \text{Aux} &= \{(x, l_1 \dots l_k) \mid \forall \alpha : x \in l_\alpha\} \subset \mathbb{P}_x^2 \times \check{\mathbb{P}}_{l_1}^2 \times \dots \times \check{\mathbb{P}}_{l_k}^2 \end{aligned}$$


Thus the cohomology class of the lifted stratum is:

$$(26) \quad [\widetilde{\Sigma}_{\mathbb{S}}(x, l_1 \dots l_k)] = \prod_{\alpha=0}^k (X + L_\alpha) \sum_{j=0}^{\binom{p+2}{2}-1} (F + (d-p)X)^{\binom{p+2}{2}-1-j} \left(\sum p_\alpha L_\alpha \right)^j$$

The class of the original stratum $[\Sigma_{\mathbb{S}}]$ is obtained by extracting the coefficient of $X^2 L_1^2 \dots L_k^2$. The numerical formula is given in Appendix. If some of p_α coincide one should also divide by the corresponding symmetry factor.

3.1.3. Some degenerating divisors in the stratum $\overline{\Sigma}_{\mathbb{S}}(x, l)$.

We need the class of a specific divisor in $\overline{\Sigma}_{\mathbb{S}}$. Suppose the generic curve of the type \mathbb{S} can be brought, in local coordinates, to some fixed Newton diagram Γ . Let $(p, q) \in \Gamma$ be a particular vertex, i.e. a point of $\mathbb{Z}_{\geq 0}^2 \cap \Gamma$, not in the interior of an edge of Γ . Erase this vertex, let $\Gamma' = \text{Conv}(\Gamma \setminus (p, q))$ be the obtained diagram. Consider the substratum $\Sigma_2 \subset \Sigma_1 = \Sigma_{\mathbb{S}}$ parameterizing of all the curves that can be brought to the diagram Γ' . Assume $\Sigma_2 \subsetneq \Sigma_1$ so that this is a genuine degeneration. As d is high $\overline{\Sigma}_2$ is a hypersurface in $\overline{\Sigma}_1$. Its cohomology class can be computed as follows.



Lemma 3.3. [Kerner06, section A.1.2] *The classes satisfy:*

$$(27) \quad [\overline{\Sigma}_2(x, l)] \left(F + (d - p - 2q)X + (q - p)L \right) = [\overline{\Sigma}_1(x, l)] \in H^*(\mathbb{P}_x^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|)$$

In particular, the class of $\overline{\Sigma}_2(x, l)$ in $\overline{\Sigma}_1(x, l)$ is the pull-back: $i^*(F) + (d - p - 2q)i^*(X) + (q - p)i^*(L)$.

Proof. Let l be the tangent line corresponding to the \hat{x}_1 axis. We need to consider an additional point $v \in l$ with $v \neq x$. Correspondingly lift the strata to a bigger auxiliary space $\mathbb{P}_x^2 \times \mathbb{P}_v^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|$ as follows:

$$(28) \quad \overline{\Sigma}_j(x, v, l) := \{(x, v, l, f) \mid (x, l, f) \in \overline{\Sigma}_j(x, l), v \in l\}$$

For the cohomology classes one has: $[\overline{\Sigma}_j(x, v, l)] = [\overline{\Sigma}_j(x, l)] \times [v \in l]$.

Observe that the line l is spanned by x, v , provided $x \neq v$. Hence the degenerating condition, erasing the vertex (p, q) , is: the directional derivative vanishes $f^{(p+q)}|_x(\underbrace{v \dots v}_q \underbrace{0 \dots 0}_p) = 0$. Here we take a particular component of the tensor. Alternatively we could consider $f^{(p+q)}|_x(\underbrace{v \dots v}_q \underbrace{\tilde{v} \dots \tilde{v}}_p) = 0$ for some fixed generic vector \tilde{v} .

This condition is transversal to the stratum $\overline{\Sigma}_j(x, v, l)$ provided: $v \neq x$ and the point $(1, 0, 0)$ does not lie on the line l , i.e. $l_0^p \neq 0$.

Thus for the cohomology classes:

$$(29) \quad [\overline{\Sigma}_2(x, l)] \times \left([v \in l] \left([f^{(p+q)}|_x(\underbrace{v \dots v}_q \underbrace{0 \dots 0}_p) = 0] - [l_0^p = 0] \right) - \text{multiplicity} \times [x = v] \right) = [\overline{\Sigma}_1(x, l)] \times [v \in l]$$

Here the diagonal $[x = v]$ is subtracted with some multiplicity, corresponding to the tangency of the degeneration. The multiplicity can be computed directly, but it is easier to obtain as follows. Note that the point v moves freely along the line l , i.e. all the participating varieties are fibrations with the fibre \mathbb{P}_v^1 . Hence in the cohomology classes all the terms quadratic in V must cancel. This fixes the multiplicity uniquely: $\text{mult} = q$.

Finally, substitute to the formula the expressions for the classes

$$(30) \quad [v \in l] = V + L, [x = v] = X^2 + XV + V^2, [f^{(p+q)}|_x(\underbrace{v \dots v}_q \underbrace{0 \dots 0}_p) = 0] = F + (d - p - q)X + qV$$

From the obtained equation extract the part proportional to V , this corresponds to the projection $\mathbb{P}_x^2 \times \mathbb{P}_v^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)| \rightarrow \mathbb{P}_x^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|$. Then one gets the equation of the lemma. ■

3.1.4. Killing tangent cone. A particular kind of degeneration that happens to be especially useful is: to increase the multiplicity of \mathbb{S} by one. Given a singularity type \mathbb{S} of multiplicity p , let $\overline{\Sigma}_1 \subset \overline{\Sigma}_{\mathbb{S}}$ be the subvariety of curves with multiplicity at least $p + 1$. In general $\overline{\Sigma}_1$ is reducible and does not correspond to a specific singularity type. For example $\overline{\Sigma}_{E_6} \cup \overline{\Sigma}_{D_6} \subset \overline{\Sigma}_{A_5}$. If \mathbb{S} is a linear type, $\overline{\Sigma}_1$ is much more restricted.

Definition-Proposition 3.4. *If \mathbb{S} is linear then the subvariety as above, $\overline{\Sigma}_1 \subset \overline{\Sigma}_{\mathbb{S}}$, is an irreducible equisingular stratum of a unique linear type \mathbb{S}' . The degeneration $\mathbb{S} \rightarrow \mathbb{S}'$ is called: **killing the tangent cone**.*

Proof. Consider the defining ideal of the lifted stratum $\overline{\Sigma}_{\mathbb{S}}$ as in equation (23). By definition the ideal of $\overline{\Sigma}_1$ is obtained by adding to the equations of $\overline{\Sigma}_{\mathbb{S}}$ the equations $f^{(p)}|_x = 0$.

This bigger set of equations is still linear in the coefficients of f and the ideal is obviously prime. Hence the generic point of $\overline{\Sigma}_1$ is well defined its singularity type too. From the defining set of equations we get that \mathbb{S}' is a linear type. ■

The cohomology class of the degeneration is given by:

Lemma 3.5. Let $T_{\mathbb{S}} = l_1^{p_1} \dots l_k^{p_k}$ be the tangent cone. For the lifting $\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\})$, the subvariety $\widetilde{\Sigma}_{\mathbb{S}'} \subset \widetilde{\Sigma}_{\mathbb{S}}$ is a divisor and its cohomology class is:

$$(31) \quad [\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\})] \left(F + (d-p)X - \sum p_i L_i \right) = [\widetilde{\Sigma}_{\mathbb{S}'}(x, \{l_i\})] \in H^*(\text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|)$$

In the paper we denote this divisor by $\text{kill}T_{\mathbb{S}}$.

Proof. Intersect the lifted stratum with the hypersurface $\{(f|_x^{(p)})_{\underbrace{0 \dots 0}_p} = 0\}$. This means that the particular entry of the tensor of derivatives vanishes. (Alternatively, one could take any collection of fixed vectors $\{v_i\}$ $\{(f|_x^{(p)})(v_1 \dots v_p) = 0\}$.)


Outside the locus $(l_1^{p_1} \dots l_r^{p_r})_{0 \dots 0} = 0$ this kills the tangent cone. Therefore we get the equation for the cohomology classes:

$$(32) \quad [\widetilde{\Sigma}_{\mathbb{S}}(x, \{l_i\})] \left([(f|_x^{(p)})_{0 \dots 0} = 0] - [(l_1^{p_1} \dots l_k^{p_k})_{0 \dots 0} = 0] \right) = [\widetilde{\Sigma}_{\mathbb{S}'}(x, \{l_i\})]$$

which proves the statement. ■

3.1.5. Rectifying the branch. This degeneration is useful in the degeneration of the curve to a reducible one (chipping off a line).

Consider a singular germ $(C, 0)$ with l one of the lines of the tangent cone. The goal is to increase the degree of the intersection: $\deg(C \cap l, 0) = k$ to $(k+1)$. To do this, choose l as one of the coordinate axes, cf. the Newton diagram.



So, the degeneration is: the monomial x_2^k should be absent. The corresponding cohomology class is given in §3.1.3, in our case it is: $F + (d-2k)X + kL$.

3.1.6. Another way to compute the class $[\widetilde{\Sigma}_{\mathbb{S}}]$. Using the class of the degenerating divisors we can compute the class $[\widetilde{\Sigma}_{\mathbb{S}}]$ in another way: by a chain of intersections with hypersurfaces.

Let $p = \text{mult}(\mathbb{S})$ and $T_{\mathbb{S}} = l_1^{p_1} \dots l_k^{p_k}$. Let $x \neq v_{\alpha} \in l_{\alpha}$ be some points on the lines. Then the defining conditions (23) can be formulated as vanishing of particular entries of the derivative-tensors: $\{f^{(n)}(v_{i_1} \dots v_{i_n}) = 0\}$, each means the absence of a particular monomial corresponding to some point on the Newton diagram. Then the class $[\widetilde{\Sigma}_{\mathbb{S}}]$ is obtained as the product of the classes of these divisors.

Example 3.6. Consider the cusp $\mathbb{S} = x_1^p + x_2^{p+1}$. Locally a curve-germ with such a singularity satisfies (we assume the \hat{x}_2 axis to be the tangent line):

$$(33) \quad \text{jet}_{p-1} f = 0, \partial_2^p f = 0, \partial_2^{p-1} \partial_1 f = 0, \dots, \partial_2 \partial_1^{p-1} f = 0$$

So the stratum is obtained from the stratum of curves with an ordinary multiple point (defined by $f^{(p-1)}|_x = 0$) by intersection with p degenerating hypersurface. The cohomology class of each such hypersurface is given in proposition 3.3. In total we have:

$$(34) \quad [\widetilde{\Sigma}_{\mathbb{S}}(x, l)] = (F + (d-p+1)X)^{\binom{p+1}{2}} \prod_{i=0}^{p-1} (F + (d+i-2p)X + (p-2i)L)$$

3.2. Equisingular strata of curves with two singular points.

Assume $\text{mult}(\mathbb{S}_x) \geq \text{mult}(\mathbb{S}_y)$. As in the case of one singular point, start

from the lifting $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}_{\mathbb{S}_x} \times \text{Aux}_{\mathbb{S}_y}$. To simplify the computation it is often useful to lift further ($\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$), taking into account various parameters relating the two singular points, for example the line $l = \overline{xy}$. If the types $\mathbb{S}_x, \mathbb{S}_y$ are distinct and the types of tangential components of \mathbb{S}_x are distinct (and the same for \mathbb{S}_y) then the projection $|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux} \rightarrow |\mathcal{O}_{\mathbb{P}^2}(d)|$ restricts to the birational morphism $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \rightarrow \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$.

$$\begin{array}{ccc} \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \subset \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} & \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux} \\ \downarrow & & \downarrow \pi \\ \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \subset \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} & \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}_{\mathbb{S}_x} \times \text{Aux}_{\mathbb{S}_y} \\ \downarrow & & \downarrow \\ \Sigma_{\mathbb{S}_x \mathbb{S}_y} \subset \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} & \subset & |\mathcal{O}_{\mathbb{P}^2}(d)| \end{array}$$

Example 3.7. In the simplest case of two ordinary multiple points there is only one (natural) additional parameter: the line $l = \overline{xy}$. Note that for the generic configuration of the singularities (i.e. $x \neq y$) the line is already fixed,

while for $x = y$ the line varies in a pencil. Therefore this additional lifting is the embedded blow-up over the diagonal $\{x = y\} \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2$. Explicitly, let $\mathbb{S}_x = x_1^{p+1} + x_2^{p+1}$, $\mathbb{S}_y = y_1^{q+1} + y_2^{q+1}$ with $p > q$, then the lifting is defined by:

$$(35) \quad \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}}}(x, y, l) = \overline{\left\{ \begin{array}{l} (x, y, l, f) \\ x \neq y \end{array} \mid f|_x^{(p)} = 0 = f|_y^{(q)}, l = \overline{xy} \right\}} \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|,$$

$$\text{Aux} := \{(x, y, l) \mid x \in l \ni y\} \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2$$

Outside the diagonal $\{x = y\}$, the vanishing of the derivatives is precisely the condition of ordinary multiple points. The fibres over the diagonal correspond to the singularity obtained by "collision" of the two ordinary multiple points, the defining equations and the singularity type are obtained by the flat limit of the equations $f|_x^{(p)} = 0 = f|_y^{(q)}$, cf. §2.3.3.

3.2.1. *The direct approach: Stepwise intersection with hypersurfaces.* For linear type \mathbb{S} the defining ideal and the cohomology class of $\overline{\overline{\Sigma_{\mathbb{S}}}}$ were given in §3.1. The stratum of curves with two singularities is the closure: $\overline{\overline{\Sigma_{\mathbb{S}_x}} \cap \overline{\overline{\Sigma_{\mathbb{S}_y}}}}$. The naive intersection results in a non-pure dimensional scheme:

$$(36) \quad \overline{\overline{\Sigma_{\mathbb{S}_x}} \cap \overline{\overline{\Sigma_{\mathbb{S}_y}}}} = \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}} \cup R_{x=y}}$$

Here the "residual piece" (the contribution from the diagonal) $R_{x=y}$ is typically non-reduced and of dimension higher than that of the needed stratum.

A way to repair this situation is to split the intersection into a step-by-step procedure of intersection with hypersurfaces (cf. [Stückrad-Vogel82], [van Gastel 89]). Start from $\overline{\overline{\Sigma_{\mathbb{S}_x}}}$ whose class is known.

Let $\overline{\overline{\Sigma_{\mathbb{S}_x}}} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}_x$ be the lifting as described above. Corresponding to the lifting $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$ of the diagram above and the projection π , define $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} := \pi^{-1}(\overline{\overline{\Sigma_{\mathbb{S}_x}}} \times \text{Aux}_y) \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$. For \mathbb{S}_x linear, the stratum $\overline{\overline{\Sigma_{\mathbb{S}_x}}}$ is smooth and by the properties of π : $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}}$ is smooth too.

The cohomology class $[\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}}] \in H^*(\text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|)$ is just the pull-back of $[\overline{\overline{\Sigma_{\mathbb{S}_x}}}]$. Consider the stratum $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}}}$ as a subvariety: $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}} \subset \overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$. Correspondingly, it is enough to calculate its class: $[\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}}] \in H^*(\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} \times \mathbb{Z})$. The resulting class in $H^*(|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}, \mathbb{Z})$ is then obtained by the pushforward. So, $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}}$ is the starting point and we reach the stratum $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}}$ by successive intersections as follows.

As was explained in §3.1.6 the stratum $\overline{\overline{\Sigma_{\mathbb{S}_y}}}$ can be obtained as a chain of degenerations: $\overline{\overline{\Sigma_{\mathbb{S}_y}}} = \overline{\overline{\Sigma}_k} \subset \overline{\overline{\Sigma}_{k-1}} \subset \dots \subset \overline{\overline{\Sigma}_0} = |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}_y$. Here at each step $\overline{\overline{\Sigma}_j}$ is a divisor in $\overline{\overline{\Sigma}_{j-1}}$, whose class is known. Take the total preimages of this chain in $|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$, denote them by the same letters. Intersect $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}}$ with these subvarieties one-by-one.

$$(37) \quad \overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} \cap \overline{\overline{\Sigma}_1} = \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_1}} \cup R_1, \dots, \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_j}} \cap \overline{\overline{\Sigma}_{j+1}} = \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_{j+1}}} \cup R_{j+1},$$

$$\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_y}} \subsetneq \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_{k-1}}} \subsetneq \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_{k-2}}} \subsetneq \dots \subsetneq \overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_1}} \subsetneq \overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}}$$

Here $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_{j+1}}}$ is the needed piece (defined as the closure: $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_j} \cap \overline{\overline{\Sigma}_{j+1}}}$). Its dimension drops precisely by one with each intersection. R_j is the residual piece produced at the j 'th step (it contains all the non-enumerative contributions).

Proposition 3.8. • *Each intersection $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_j} \cap \overline{\overline{\Sigma}_{j+1}}$ results in a pure dimensional scheme $\overline{\overline{\Sigma_{\mathbb{S}_x \mathbb{S}_{j+1}}} \cup R_{j+1}$.*

• *For linear types $\mathbb{S}_x \mathbb{S}_y$, the scheme $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} \cap_{x \neq y}^{j+1} \overline{\overline{\Sigma}_i}$ is irreducible and reduced.*

The residual piece R_{j+1} is typically reducible and non-reduced.

Proof. • Note that the initial variety is irreducible, thus each components of the resulting union $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} \cap_{x \neq y}^j \overline{\overline{\Sigma}_i} \cap \overline{\overline{\Sigma}_{j+1}}$ is of (strictly) smaller dimension. Conversely, the dimension of each component cannot drop by more than one (since the intersection is with a hypersurface and the ambient space is irreducible).

• Note that the defining equations of both $\overline{\overline{\Sigma_{\mathbb{S}_x}}^{(y)}} \cap_{x \neq y}^j \overline{\overline{\Sigma}_i}$ and $\overline{\overline{\Sigma}_{j+1}}$ are linear in f (i.e. in the coordinates of $|\mathcal{O}_{\mathbb{P}^2}(d)|$).

■

Thus the contribution of the residual piece can be subtracted: $[\widetilde{\Sigma}_{\mathbb{S}_x \cap_{i=1}^{j+1} \widetilde{\Sigma}_i}] = [\widetilde{\Sigma}_{\mathbb{S}_x \cap_{i=1}^j \widetilde{\Sigma}_i}] [\widetilde{\Sigma}_{j+1}] - [R_{j+1}]$ (with multiplicities for R_{j+1}). By repeating this procedure we calculate the needed class $[\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}]$.

Therefore the enumerative problem is reduced to geometry of the residual pieces and their multiplicities. This method is applied to the case of two ordinary multiple points in §4.1. Explicit numerical results are given in the theorem 4.2, corollary 4.3 and in the Appendix A.

3.2.2. Degenerations to higher singularities. Let $\mathbb{S}_x \mathbb{S}_x'$ be two adjacent singularity types, i.e. $\widetilde{\Sigma}_{\mathbb{S}_x'} \subsetneq \widetilde{\Sigma}_{\mathbb{S}_x} \subset |\mathcal{O}_{\mathbb{P}^2}(d)|$. Let $Degen \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}_x$ be a (reduced) "degenerating cycle" such that $Degen \cap \widetilde{\Sigma}_{\mathbb{S}_x} = \widetilde{\Sigma}_{\mathbb{S}_x'}$ (at least set theoretically). These degenerating cycles and their cohomology classes are described in §3.1.3. Then for the cohomology classes we have (with some multiplicities): $[\widetilde{\Sigma}_{\mathbb{S}_x}][Degen] \sim [\widetilde{\Sigma}_{\mathbb{S}_x'}] \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux})$. The multiplicity here is the local degree of intersection of $Degen$ with $\widetilde{\Sigma}_{\mathbb{S}_x}$.

This equation allows to calculate $[\widetilde{\Sigma}_{\mathbb{S}_x'}]$ in terms of $[\widetilde{\Sigma}_{\mathbb{S}_x}]$. The key observation is that such a degeneration is always "invertible": the above equation has unique solution for $[\widetilde{\Sigma}_{\mathbb{S}_x'}]$ in terms of $[\widetilde{\Sigma}_{\mathbb{S}_x}]$ and $[Degen]$. This happens because the class $[Degen]$ contains a monomial: a power of the generator of $H^*(|\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z})$. And d is arbitrary high, so this monomial is not a zero divisor when multiplying bounded classes, i.e. lying in $\oplus_{i=0}^k H^i(|\mathcal{O}_{\mathbb{P}^2}(d)|)$ for $k \ll \dim |\mathcal{O}_{\mathbb{P}^2}(d)|$ (cf. [Kerner06, §2.2]).

Using this, we can degenerate to some higher singularity types for which the classes are known. In §4.1.2 we degenerate a curve with two ordinary multiple points to a reducible curve, with the line through the points as a component. Once the cohomology class of $\widetilde{\Sigma}_{x_1^{p+1}+x_2^{p+1}, x_1^{q+1}+x_2^{q+1}}$ is obtained, we can continue to other singularities, degenerating them to a pair of ordinary multiple points.

A suitable degeneration is by increasing the multiplicity of \mathbb{S}_x or \mathbb{S}_y . For the original stratum $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \subset |\mathcal{O}_{\mathbb{P}^2}(d)|$ such degeneration is usually of high codimension. To turn this into a degeneration in codimension 1, i.e. intersection by a hypersurface, consider the lifting that traces all the lines of the tangent cone: $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}(x, y, l, \{l_{x,i}\}, \{l_{y,j}\})$. Then the degeneration is done by intersection with $\text{kill}T_{\mathbb{S}}$ divisor from §3.1.4.

Hence the algorithm consists of several steps: each time increasing the multiplicity of \mathbb{S}_x or \mathbb{S}_y . At each step the variety is intersected by the degenerating hypersurface, the result is reducible:

$$(38) \quad \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \cap \text{kill}T_{\mathbb{S}_x} = \widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y} \cup R^{x=y} \quad \text{or} \quad \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \cap \text{kill}T_{\mathbb{S}_y} = \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y'} \cup R^{x=y},$$

Corollary 3.9. Let $\mathbb{S}_x \mathbb{S}_y$ be linear types. The degeneration approach reduces the enumerative problem to understanding the irreducible components of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$.

Proof. First note that each step gives an equation (in the cohomology ring of the ambient space $\text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|$) from which the class of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}$ is restored in terms of the classes of $\widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y}$, $\text{kill}T_{\mathbb{S}_x}$ (or $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y'}$, $\text{kill}T_{\mathbb{S}_y}$), $R^{x=y}$.

Indeed, the resulting variety $\widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y} \cup R^{x=y}$ is pure dimensional. The class of $\text{kill}T_{\mathbb{S}_x}$ is given in §3.1.4. The residual pieces $R^{x=y}$ are (as sets) unions of the equisingular strata $\widetilde{\Sigma}_{\mathbb{S}}$, where \mathbb{S} 's are the results of collision 2.3.2. In other words: $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y} = \cup \widetilde{\Sigma}_{\mathbb{S}}$. The way to compute the multiplicities of the decomposition $[\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}] = \sum m_i [\widetilde{\Sigma}_{\mathbb{S}_i}]$ is explained in §2.3.4.

Finally, after a finite number of steps the algorithm stops. This statement is immediate since e.g. one can reach an ordinary multiple point in a finite number of steps. In fact, by killing the tangent cone, cf. §3.1.4 one can do this in less than $o.d.(\mathbb{S}) - \text{mult}(\mathbb{S})$ steps. ■

Examples of the degenerations are in §4.2. Below we discuss the relevant components of $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y}$ contributing to residual pieces.

3.2.3. Residual pieces for the degeneration $\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \rightarrow \widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y}$. Assume $\mathbb{S}_x, \mathbb{S}_y$ are linear types and $\text{mult}(\mathbb{S}_x) \geq \text{mult}(\mathbb{S}_y)$. Let $\text{kill}T_{\mathbb{S}_x}$ denote the "killing tangent cone" divisor (as in lemma 3.5), so that $\widetilde{\Sigma}_{\mathbb{S}_x} \cap \text{kill}T_{\mathbb{S}_x} = \widetilde{\Sigma}_{\mathbb{S}_x'}$. For the case of two singular points:

$$(39) \quad \widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y} \cap \text{kill}T_{\mathbb{S}_x} = \widetilde{\Sigma}_{\mathbb{S}_x' \mathbb{S}_y} \cup \left(\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}|_{x=y} \cap \text{kill}T_{\mathbb{S}_x} \right)$$

Let $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y} = \cup_j \widetilde{\Sigma}_j$ be the decomposition into irreducible components. As $\dim(\widetilde{\Sigma}_j) \leq \dim(\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}) - 1$ the component $\widetilde{\Sigma}_j$ contributes to the equation in cohomologies iff $\widetilde{\Sigma}_j \subset \text{kill}T_{\mathbb{S}_x}$.

From the local defining equation of $\text{kill}T_{\mathbb{S}_x}$ (§3.1.4) we get therefore:

Proposition 3.10.

$$(40) \quad [\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}][\text{kill}T_{\mathbb{S}_x}] = [\widetilde{\Sigma}_{\mathbb{S}_x'\mathbb{S}_y}] + \sum_{j \in J} \text{tang}_j[\widetilde{\Sigma}_j]$$

where:

★ J indexes the irreducible components of $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}$ corresponding to collisions $\mathbb{S}_x + \mathbb{S}_y \rightarrow \mathbb{S}$ with $\text{mult}(\mathbb{S}) > \text{mult}(\mathbb{S}_x)$.

★ $\widetilde{\Sigma}_j$ is the corresponding stratum lifted to the given ambient space.

★ tang_j is the degree of tangency of $\text{kill}T_{\mathbb{S}_x}$ and $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ along $\widetilde{\Sigma}_j$.

3.2.4. *Residual pieces for the degeneration* $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y} \rightarrow \widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}'$. Now the degeneration is done by intersection with the divisor $\text{kill}T_{\mathbb{S}_y}$. As $\text{mult}(\mathbb{S}_y) \leq \text{mult}(\mathbb{S}_x)$ all the irreducible components of $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}$ will contribute to the cohomology class:

$$(41) \quad [\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}][\text{kill}T_{\mathbb{S}_y}] = [\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}'] + \sum_{\substack{\text{components of} \\ \widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}}} \text{tang}_j[\widetilde{\Sigma}_j]$$

where tang_j is the degree of tangency of $\text{kill}T_{\mathbb{S}_y}$ and $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ along $\widetilde{\Sigma}_j$. Let $q = \min(\text{tang}_j)$. In many cases $\text{tang}_j > q$ only for a very few components. Then we can write:

$$(42) \quad [\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}]([\text{kill}T_{\mathbb{S}_y}] - q[E_{x=y}]) = [\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}'] + \sum_{\substack{\text{components of} \\ \widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}}} (\text{tang}_j - q)[\widetilde{\Sigma}_j]$$

where $E_{x=y} = \widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}|_{x=y}$ is the "exceptional divisor". This form simplifies the computations.

3.3. On the validity range of results. From the explanation of the method it is clear that an (obvious) necessary condition for the method to be applicable is: the stratum $\Sigma_{\mathbb{S}_x\mathbb{S}_y}$ is smooth, irreducible of expected dimension.

A cheap sufficient condition is: $d \geq o.d.(\mathbb{S}_x) + o.d.(\mathbb{S}_y)$ (here $o.d.(\mathbb{S})$ is the degree of determinacy, cf. the end of §2.2). The bound can be obtained as follows. Start from the stratum $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ and degenerate to the types: ordinary multiple points of multiplicities $o.d.(\mathbb{S}_x), o.d.(\mathbb{S}_y)$. The formulas certainly hold for $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ if the corresponding formulas hold for the degenerated stratum. And in this later case an easy sufficient condition is $d \geq o.d.(\mathbb{S}_x) + o.d.(\mathbb{S}_y)$ (cf. theorem 4.2).

Of course, this sufficient bound is far from being necessary.

4. EXAMPLES

4.1. Two ordinary multiple points. Let $\mathbb{S}_x = x_1^{p+1} + x_2^{p+1}$, $\mathbb{S}_y = y_1^{q+1} + y_2^{q+1}$, $p \geq q$. As in the example 3.1.1 the natural candidate for the lifting of $\Sigma_{\mathbb{S}_x\mathbb{S}_y}$ is the variety of triples (x, y, f) with f having \mathbb{S}_y at y and \mathbb{S}_x at x . To simplify the embedding we blowup the auxiliary space $\mathbb{P}_x^2 \times \mathbb{P}_y^2$ along the diagonal $\Delta = \{x = y\}$. Geometrically we add the line $l = \overline{xy}$ (defined by a one-form, a point in the dual plane $\check{\mathbb{P}}_l^2$). The exceptional divisor is $E = \{(x, y, l) | x = y, l(x) = 0\}$. Thus the lifted stratum is defined as the strict transform:

$$(43) \quad \begin{aligned} \widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l) &:= \overline{\{(x, y, l, f), x \neq y | f|_x^{(p)} = 0 = f|_y^{(q)}, l(x) = 0 = l(y)\}} \subset \text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)| \\ \text{Aux} &= \{(x, y, l) | l(x) = 0 = l(y)\} \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2 \end{aligned}$$

4.1.1. *Stepwise intersection.*

Theorem 4.1. *The projection $\widetilde{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l) \rightarrow \text{Aux} \subset \mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2$ is a projective fibration over a smooth base, the projectivization of a vector bundle. In particular the lifted stratum is a smooth locally complete intersection.*

Proof. As the degree is high, the defining conditions are transversal outside the diagonal $x = y$. (To check a specific fibre, fix the points x, y then the conditions are just linearly independent equations in $|\mathcal{O}_{\mathbb{P}^2}(d)|$.) Therefore the topological closure $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l)$ is an irreducible reduced algebraic variety.

For $x \neq y$ the fibre of $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l) \rightarrow \text{Aux}$ over (x, y, l) is a linear subspace of $|\mathcal{O}_{\mathbb{P}^2}(d)|$. Consider the fibres over the diagonal. The points of $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l)|_{x=y}$ correspond to collision of the two ordinary multiple points. As is shown in §2.3.3, for $\mathbb{S}_x\mathbb{S}_y$ ordinary multiple points, the restriction is irreducible: $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l)|_{x=y} = \overline{\Sigma}_{\mathbb{S}}(x, l, f)$. Here \mathbb{S} is the singularity type of $(x_1^{p-q} + x_2^{p-q})(x_1^{q+1} + x_2^{2q+2})$ and l is the line tangent to $(x_1^{q+1} + x_2^{2q+2})$. Note that the fibre over $(x = y, l)$ is a linear subspace of $|\mathcal{O}_{\mathbb{P}^2}(d)|$. Its co-dimension (computed e.g. as the number of $\mathbb{Z}_{\geq 0}^2$ points strictly below the Newton diagram) is $\binom{p+2}{2} + \binom{q+2}{2}$, i.e. precisely the codimension of the general fibre over (x, y, l) .

Hence, as in the proof of proposition 2.9, there is the natural morphism: $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y} \rightarrow \text{Aux} \rightarrow \text{Gr}(|\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^2}(d)|)$, giving to $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ the structure of projective fibration as on the diagram.

$$(44) \quad \begin{array}{ccc} \overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l) & \xrightarrow{i} & \tau \quad \subset \text{Gr}(\mathbb{P}^n, |\mathcal{O}_{\mathbb{P}^2}(d)|) \times |\mathcal{O}_{\mathbb{P}^2}(d)| \\ \downarrow & & \downarrow \\ \text{Aux} & \xrightarrow{i} & \text{Gr}(\mathbb{P}^n, |\mathcal{O}_{\mathbb{P}^2}(d)|) \end{array}$$

Here τ is the tautological fibration, the fibre over $[\mathbb{P}^n] \in \text{Gr}$ is $\mathbb{P}^n \subset |\mathcal{O}_{\mathbb{P}^2}(d)|$. ■

Theorem 4.2. For $p \geq q$ the cohomology class of the lifted stratum $[\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l)] \in H^*(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z})$ is given by:

$$(45) \quad [\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l)] = (L + X)(L + Y) \left(F + (d - p)X \right)^{\binom{p+2}{2}} \prod_{i=0}^q \prod_{j=0}^{q-i} \left(F + (d - i - j)Y + iX - jL - (p + 1 + i - j)E \right)$$

here $E = X + Y - L$ is the class of the exceptional divisor. (For the notations of the cohomology generators cf. §2.1.1). The formula is applicable for $d \geq p + q + 2$.

To get the solution of the enumerative problem (i.e. the degree of $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$) we should apply the Gysin homomorphism corresponding to the projection $\overline{\Sigma}(x, y, l) \rightarrow \overline{\Sigma}$. In this case it means just to extract the coefficient of $X^2Y^2L^2$. In the case $\mathbb{S}_x = \mathbb{S}_y$ (i.e. $p = q$) the resulting answer should be also divided by 2, as the singular points are not ordered.

Corollary 4.3. In a few simplest cases the degree of $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}$ is:

- $q = 1$.
 $\deg(\overline{\Sigma}_{x_1^{p+1}+x_2^{p+1}, A_1}) = 9 \binom{p+3}{4} (d-p)^3 (d+p-2) - \frac{3}{4} \binom{p+2}{3} (10p^2 + 39p + 7)(d-p)^2 + 3 \binom{p+2}{3} (d-p)(6 + 5p)$
- $q = 2$.
 $\deg(\overline{\Sigma}_{x_1^{p+1}+x_2^{p+1}, D_4}) = \frac{45}{8} \binom{p+3}{4} (d-p)^3 (d+p-4) + 2(d-p)(8 + 3p + p^2)(35p^2 + 20p - 12) - \frac{5}{8} (p+1)(d-p)^2 (14p^4 + 105p^3 + 147p^2 + 114p - 80) - 6(85p^2 + 45p - 28)$
- $q = 3$.
 $\deg(\overline{\Sigma}_{x_1^{p+1}+x_2^{p+1}, X_9}) = \frac{135}{8} \binom{p+3}{4} (d-p)^3 (d+p-6) + 2(d-p)(16 + 3p + p^2)(270p^2 - 20p - 117) - \frac{5}{8} (d-p)^2 (54p^5 + 527p^4 + 948p^3 + 1853p^2 - 894p - 1152) - 14(830p^2 - 105p - 348)$

For $p = q = 1$ this gives the classical result [Aluffi98, pg. 2]. For $p = 2, q = 1$ this coincides the results of [Kleiman-Piene98] and [Kazarian03-hab]. For other cases the results seem to be new.

Remark 4.4. It is immediate from (43) that the stratum is symmetric with respect to the permutation $(x, p) \leftrightarrow (y, q)$. So, its cohomology class is symmetric in $(X, p) \leftrightarrow (Y, q)$. The answer in 4.2 is obtained (in a very non-symmetric way) under the assumption $p \geq q$. One might hope that this answer is still symmetric. However this happens for (p, p) and $(p + 1, p)$ cases only. So, for the general answer one should substitute $\max(p, q)$ and $\min(p, q)$, i.e. the degree is written as $\deg \overline{\Sigma}_{\max(\mathbb{S}_x, \mathbb{S}_y), \min(\mathbb{S}_x, \mathbb{S}_y)}$ with the obvious order on the types of ordinary multiple points.

This contradicts the naive expectation of algebraicity (as mentioned in §1.3): in general the cohomology classes of equisingular strata depend non-algebraically on the classical singularity invariants.

proof of the theorem:

As in §3.2.1 represent $\overline{\Sigma}_{\mathbb{S}_x\mathbb{S}_y}(x, y, l)$ as a stepwise intersection of the stratum $\overline{\Sigma}_{\mathbb{S}_x}^{(y)}(x, y, l)$ with hypersurfaces comprising

the conditions $f|_y^{(q)} = 0$. Start from the case of one singular point

$$(46) \quad \overline{\Sigma}_{\mathbb{S}_x}^{(y)}(x, y, l) := \left\{ (x, y, l, f) \mid f|_x^{(p)} = 0, l(x) = 0 = l(y) \right\} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$$

At j 'th step we have a variety $M_j \xrightarrow{i} |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$ and a hypersurface $V_{j+1} \subset |\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux}$. Think about the intersection $M_j \cap V_{j+1}$ as the pullback $i^*(V_{j+1})$. Then the task is to take the *strict transform* of V_{j+1} , i.e. to subtract from the pullback the part of the exceptional divisor over $\{x = y\}$.

The straightforward approach is just to consider the components of the tensor $f|_y^{(q)}$ (i.e. all the partials) and intersect $\overline{\Sigma}_{\mathbb{S}_x}^{(y)}(x, y, l)$ with the corresponding hypersurfaces: $\{\partial_0^{n_0} \partial_1^{n_1} \partial_2^{n_2} f|_y = 0\}_{n_0 + \dots + n_2 = q}$. This will bring various complicated residual pieces. Instead, we represent these conditions as follows (cf. §2.1.1 for notations):

$$(47) \quad \left\{ f|_y^{(i)}(\underbrace{x \dots x}_i) = 0 \right\}_{i=0}^q, \left\{ f|_y^{(i+1)}(\underbrace{x \dots x \tilde{v}}_i) = 0 \right\}_{i=0}^{q-1}, \dots, \left\{ f|_y^{(i+j)}(\underbrace{x \dots x \tilde{v} \dots \tilde{v}}_i \underbrace{\tilde{v} \dots \tilde{v}}_j) = 0 \right\}_{i=0}^{q-j}, f|_y^{(q)}(\underbrace{\tilde{v} \dots \tilde{v}}_q) = 0$$

Here \tilde{v} is a fixed point, so that the points x, y, \tilde{v} generically do not lie on one line. By direct check it is verified that for generic parameters (i.e. $y \neq x$, $\tilde{v} \notin \overline{xy} = l$) these conditions are equivalent to $f|_y^{(q)} = 0$. For non-generic situation each such equation will give a reducible hypersurface in M_j , correspondingly a residual term should be subtracted.

• $M_0 := \overline{\Sigma}_{\mathbb{S}_x}(x, y, l) \cap_{x \neq y} \{f|_y = 0\}$. The pullback of the hypersurface $\{f|_y = 0\}$ to $\overline{\Sigma}_{\mathbb{S}_x}(x, y, l)$ consists of the strict transform (the closure of the part over $x \neq y$) and the exceptional divisor E (over $x = y$). To calculate the multiplicity, expand $y = x + \epsilon v$, correspondingly:

$$(48) \quad 0 = f|_y = \underbrace{f|_x + \dots + \epsilon^p f|_x^{(p)}(v \dots v)}_{\text{vanish}} + \epsilon^{p+1} f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) + \dots$$

i.e. the exceptional divisor enters with the multiplicity $(p+1)$. So, the strict transform is $(f|_y = 0) - (p+1)E$ and the total cohomology class: $[\overline{\Sigma}_{\mathbb{S}_x}(x, y, l)] \left([f|_y = 0] - (p+1)[E] \right) \in H^*(|\mathcal{O}_{\mathbb{P}^2}(d)| \times \text{Aux})$.

The points of M_0 satisfy:

- ★ for $y \neq x$: $f|_x^{(p)} = 0$ and $f|_y = 0$
- ★ for $y = x$: $f|_x^{(p)} = 0 = f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1})$

• In the same way do all the intersections with $\left\{ f|_y^{(i)}(\underbrace{x \dots x}_i) = 0 \right\}_{i=1}^q$ (i.e. $M_i := \overline{M_{i-1} \cap_{x \neq y} \left\{ f|_y^{(i)}(\underbrace{x \dots x}_i) = 0 \right\}}$). At each step subtract the exceptional divisor with the necessary multiplicity. The resulting cohomology class is:

$$(49) \quad [\overline{\Sigma}_{\mathbb{S}_x}(x, y, l)] \prod_{i=0}^q \left([f|_y^{(i)}(\underbrace{x \dots x}_i) = 0] - [(p+1+i)E] \right)$$

The points of M_q satisfy:

- ★ for $y \neq x$: $f|_x^{(p)} = 0$ and $f|_y = 0 = f|_y^{(1)}(x) = \dots = f|_y^{(q)}(\underbrace{x \dots x}_q)$
- ★ for $y = x$: $f|_x^{(p)} = 0 = f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) = \dots = f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1})$

• Intersect with $\left\{ f|_y^{(i+1)}(\underbrace{x \dots x \tilde{v}}_i) = 0 \right\}_{i=0}^{q-1}$. Here \tilde{v} is a fixed point in \mathbb{P}^2 . The intersection of each such hypersurface with the previously obtained variety (say M_j) is reducible: it contains a component over $\tilde{v} \in \overline{xy} = l$. (Note that this component is a divisor.) So, one should take the "strict transform": $M_{j+1} = \left(\left\{ f|_y^{(i+1)}(\underbrace{x \dots x \tilde{v}}_i) = 0 \right\} - \{ \tilde{v} \in \overline{xy} \} \right) \cap M_j$.

Now, every point of M_{j+1} satisfies: $f|_y^{(i+1)}(\underbrace{x \dots x}_i) = 0$.

Now, check the situation over the diagonal $x = y$, subtract its contribution as in the previous cases:

$$(50) \quad [\overline{\Sigma}_{\mathbb{S}_x}(x, y, l)] \prod_{i=0}^q \left([f|_y^{(i)}(\underbrace{x \dots x}_i) = 0] - [(p+1+i)E] \right) \prod_{i=0}^{q-1} \left([f|_y^{(i+1)}(\underbrace{x \dots x \tilde{v}}_i) = 0] - [\tilde{v} \in l] - [(p+i)E] \right)$$

Here $\tilde{v} \in \overline{xy} = l$ is a divisor (all the lines in the plane through the fixed point \tilde{v}). Its cohomology class: $[\tilde{v} \in l] = L$. It should be subtracted with multiplicity 1. To check this expand as previously (eq. 48): $\tilde{v} = v + \epsilon \tilde{v}_1$ (with $v \in l$, $\tilde{v}_1 \notin l$), then

$$(51) \quad f|_y^{(i+1)}(\underbrace{x \dots x}_i \tilde{v}) = f|_y^{(i+1)}(\underbrace{x \dots x}_i v) + \epsilon f|_y^{(i+1)}(\underbrace{x \dots x}_i \tilde{v}_1)$$

$\underbrace{\hspace{10em}}_{\text{vanishes}}$

The points of the so obtained variety satisfy:

★ for $y \neq x$: $f|_x^{(p)} = 0$ and $\{f|_y^{(i)}(x \dots x) = 0\}_{i=1}^q$

★ for $y = x$: $f|_x^{(p)} = 0 = f|_x^{(p+1)}(\underbrace{v \dots v}_p) = \dots = f|_x^{(p+q)}(\underbrace{v \dots v}_{p+q-1}) = f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1})$

• Do the rest of intersections, at each step subtracting (with appropriate multiplicities) the exceptional divisor and the class $[\tilde{v} \in l]$. Finally we get:

$$(52) \quad [\widetilde{\Sigma}_{\mathbb{S}_x \mathbb{S}_y}(x, y, l)] = [\widetilde{\Sigma}_{\mathbb{S}_x}(x, y, l)] \prod_{i=0}^q \prod_{j=0}^{q-i} \left([f|_y^{(i+j)}(\underbrace{x \dots x}_i \underbrace{\tilde{v} \dots \tilde{v}}_j) = 0] - j[\tilde{v} \in l] - [(p+1+i-j)E] \right)$$

The points of this variety satisfy:

★ for $y \neq x$: $f|_x^{(p)} = 0 = f|_y^{(q)}$


★ for $y = x$: $f|_x^{(p)} = 0 = f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1-q}) = f|_x^{(p+2)}(\underbrace{v \dots v}_{p+3-q}) = \dots = f|_x^{(p+q+1)}(\underbrace{v \dots v}_{p+q+1})$

Substitute now the cohomology classes for the conditions (cf. §2). (Note that as \tilde{v} is a fixed point, the condition $\tilde{v} \in l$ is just one linear condition on l , its class is L . The class of the exceptional divisor is $[E] = X + Y - L$, cf. §2.1.1.) This proves the formula.

Finally, regarding the sufficient bound for validity of the formula. Note that all the conditions that have appeared in the proof concern only the $(p+q+1)$ 'th jet of the function f . So, the bound $d \geq p+q+2$ is sufficient. ■

4.1.2. Degenerations: chipping off a line.

Use the following elementary observation. Let $C_{x_{p+1}, y_{q+1}}^{(d)}$ be a curve of degree d with the points x_{p+1}, y_{q+1} of multiplicities $p+1, q+1$. Let $l = \overline{x_{p+1}, y_{q+1}}$. Suppose the intersection multiplicity satisfies: $|l \cap C_{x_{p+1}, y_{q+1}}^{(d)}| > d$. Then the curve is reducible: $C_{x_{p+1}, y_{q+1}}^{(d)} = l \cup C_{x_p, y_q}^{(d-1)}$. We degenerate to get to this situation. The degeneration is done by forcing one of the branches at x to be tangent to l with high degree of tangency, i.e. "killing" the corresponding monomials (cf. the Newton diagram).



This amounts to intersection of the stratum with the corresponding "degenerating hypersurfaces". Their cohomology

classes are given in §3.1.3. Let $\widetilde{\Sigma}_{p+1, q+1}^{(d)}$ denote the lifted stratum of degree d curves with two ordinary multiple points, of multiplicities i, j .

Theorem 4.5. *The intersection of the stratum $\widetilde{\Sigma}_{p+1, q+1}^{(d)}$ with $(d-q-p-1)$ degenerating hypersurfaces results in the stratum of reducible curves $\Xi_{p, q}^{(d-1)} := \left\{ (x, y, l, C_d) \mid C_d = l \cup C_{d-1}, (x, y, l, C_{d-1}) \in \widetilde{\Sigma}_{p, q}^{(d-1)}(x, y, l) \right\}$. In particular, the cohomology classes of the two strata are related by the equation:*

$$(53) \quad [\widetilde{\Sigma}_{p+1, q+1}^{(d)}(x, y, l)] \prod_{k=p+1}^{d-q-1} \left(F + (d-k)X + kY - (k+q+1)E \right) = [\Xi_{p, q}^{(d-1)}] \in H^*(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|), \text{ here } E = X + Y - L$$

Proof. Start from the stratum $\widetilde{\Sigma}_{p+1, q+1}^{(d)}$, the corresponding curves satisfy: $f^{(p)}|_x = 0 = f^{(q)}|_y$. Consider the degenerating hypersurfaces:

$$(54) \quad D_{p+1} := \{f|_x^{(p+1)}(\underbrace{y \dots y}_{p+1}) = 0\}, D_{p+2} := \{f|_x^{(p+2)}(\underbrace{y \dots y}_{p+2}) = 0\}, \dots, D_{d-q-1} := \{f|_x^{(d-q-1)}(\underbrace{y \dots y}_{d-q-1}) = 0\}$$

These conditions, imposed onto the curve $C_{x_{p+1}, y_{q+1}}^{(d)}$, force the intersection multiplicity: $\text{mult}_x(C_{x_{p+1}, y_{q+1}}^{(d)}, \overline{xy}) \geq (p+1) + (d-p-q-1) = d-q$. (Pass to the local coordinate system to verify that these are precisely the derivatives in the given direction.)

Hence, outside the diagonal $\{x=y\}$ one has:

$$(55) \quad \overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l) \cap_{x \neq y} \bigcap_{j=p+1}^{d-q-1} D_j = \Xi_{p, q}^{(d-1)} \cap \{x \neq y\}$$

The cohomology classes are (see §3.1.3): $[D_j] = F + (d-j)X + jL \in H^2(\mathbb{P}_x^2 \times \mathbb{P}_y^2 \times \check{\mathbb{P}}_l^2 \times |\mathcal{O}_{\mathbb{P}^2}(d)|)$.

Consider the intersection near the diagonal. Recall from §2.3.2, that the points over the diagonal $\overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l)|_{x=y}$ correspond to the type with the representative: $(x^{p-q} + y^{p-q})(x^{q+1} + y^{2q+2}) = x^{p+1} + x^{q+1}y^{p-q} + y^{p+q+2}$. The defining equations for such a fibre are:

$$(56) \quad f|_y^{(p)} = 0, \underbrace{f|_y^{(p+1)}}_{p-q+1}(v \dots v) = 0, \dots, \underbrace{f|_y^{(p+q+1)}}_{p+q+1}(v \dots v) = 0$$

here v is a point on the line l , $x \neq v$. Thus the intersection $D_{p+1} \cap \overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l)$ contains $\overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l)|_{x=y}$ as an irreducible component. To find the multiplicity of this component, expand $x = y + \epsilon v$:

$$(57) \quad f|_x^{(p+1)}(\underbrace{y \dots y}_{p+1}) = \dots + \underbrace{\epsilon^q f|_y^{(p+q+1)}(v \dots v)}_{\text{vanish on } \overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l)|_{x=y}} + \epsilon^{q+1} f|_y^{(p+q+2)}(v \dots v) + \dots$$

Therefore for the cohomology classes have:

$$(58) \quad \overline{[\Sigma_{p+1, q+1}^{(d)}(x, y, l)]} [D_{p+1}] = \overline{[\Sigma_{p+1, q+1}^{(d)}(x, y, l) \cap_{x \neq y} D_{p+1}]} + (p+q+2) \overline{[\Sigma_{p+1, q+1}^{(d)}(x, y, l)|_{x=y}]} \\ \text{i.e. } \overline{[\Sigma_{p+1, q+1}^{(d)}(x, y, l)]} ([D_{p+1}] - (p+q+2)E) = \overline{[\Sigma_{p+1, q+1}^{(d)}(x, y, l) \cap_{x \neq y} D_{p+1}]}$$

which is the first term in the product of the statement. Continue by induction.

After the k 'th step the points of the stratum $\overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l) \cap_{x \neq y} \bigcap_{i=1}^k D_{p+i}$ satisfy:

$$(59) \quad f|_x^{(p)} = 0 = f|_y^{(q)}, \underbrace{f|_x^{(p+1)}}_{p+1}(y \dots y) = 0, \dots, \underbrace{f|_x^{(p+k)}}_{p+k}(y \dots y) = 0$$

To check the intersection of D_{p+k+1} on the diagonal we should obtain the defining equations of $\overline{\Sigma}_{p+1, q+1}^{(d)}(x, y, l) \cap_{x \neq y} \bigcap_{i=1}^k D_{p+i}$ near $x = y$. This is done by the flat limit. The procedure similar to that of 2.3.2 gives the degree of the intersection D_{p+k+1} with the diagonal: $p+q+2+k$. So, the k 'th step brings to the product the factor: $[D_{p+k+1}] - (p+q+2+k)E$. ■

By applying this process $q+1$ times one arrives at a curve with an ordinary multiple point of multiplicity $(p+1)$ and a line of multiplicity $q+1$ as a component. For such a stratum the answer is known classically. As every intersection is invertible this solves the enumerative problem.

4.2. The class of $\overline{\Sigma}_{\mathbb{S}_x, A_1}$ by degenerations to ordinary multiple point. We consider here the simplest case: $\mathbb{S}_y = A_1$, then the degeneration process of $\overline{\Sigma}_{\mathbb{S}_x, A_1}$ is quite explicit.

Let $T_{\mathbb{S}_x} = l_1^{p_1} \dots l_k^{p_k}$ be the tangent cone of \mathbb{S}_x and $\overline{\Sigma}_{\mathbb{S}_x}(x, l_1 \dots l_k)$ the corresponding lifting, as in §3.1. So, we start from the lifted stratum:

$$(60) \quad \overline{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, l_1 \dots l_k) := \left\{ \overline{(x, y, l, l_1 \dots l_k, C)} \mid \begin{array}{l} C \text{ has } \mathbb{S}_x \text{ at } x \text{ and } A_1 \text{ at } y \\ x \neq y, l = \overline{xy} \\ T_{(C, x)} = l_1^{p_1} \dots l_k^{p_k} \end{array} \right\}$$

Intersect with the divisor $\text{kill}T_{\mathbb{S}_x}$, cf. §3.1.4, such that $\overline{\Sigma}_{\mathbb{S}_x}(x, l_1 \dots l_k) \cap \text{kill}T_{\mathbb{S}_x} = \overline{\Sigma}_{\mathbb{S}_x'}(x, l_1 \dots l_k)$. Here \mathbb{S}_x' is the degenerated singularity type, recall that if \mathbb{S}_x is linear singularity then \mathbb{S}_x' is well defined.

Note that $T_{\mathbb{S}_x} \neq T_{\mathbb{S}_x'}$, hence some of l_i of $\overline{\Sigma}_{\mathbb{S}_x'}(x, l_1 \dots l_k)$ can be not related to the parameters of the singularity \mathbb{S}_x' . In the extremal case, when \mathbb{S}_x' is an ordinary multiple point, one has: $\overline{\Sigma}_{\mathbb{S}_x'}(x, l_1 \dots l_k) = \overline{\Sigma}_{\mathbb{S}_x'}(x) \cap \bigcap_i \{x \in l_i\}$.

The intersection $\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, l_1 \dots l_k) \cap \text{kill}T_{\mathbb{S}_x}$ brings in addition the residual pieces over the diagonal:

Theorem 4.6. *Killing the tangent cone results in the cohomological equation in $H^*(\text{Aux} \times |\mathcal{O}_{\mathbb{P}^2}(d)|, \mathbb{Z})$*

$$(61) \quad \left[\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, \{l_i\}) \right] [\text{kill}T_{\mathbb{S}_x}] = \left[\widetilde{\Sigma}_{\mathbb{S}_x', A_1}(x, y, l, \{l_i\}) \right] + 2[x = y] \left[\widetilde{\Sigma}_{\mathbb{S}_1}(x, l, \{l_i\}) \right] + [x = y] \sum_{\substack{i \text{ such that} \\ p_i = 1}} [l = l_i] \left[\widetilde{\Sigma}_{\mathbb{S}_2}(x, l, \{l_i\}) \right]$$

where the strata for $\mathbb{S}_1, \mathbb{S}_2$ are defined by

$$(62) \quad \begin{aligned} \widetilde{\Sigma}_{\mathbb{S}_1}(x, l, \{l_i\}) &= \widetilde{\Sigma}_{\mathbb{S}_x'}(x, \{l_i\}) \cap \left\{ x \in l, f|_x^{(p)} = 0 = f|_x^{(p+1)}(\underbrace{v \dots v}_p) \forall v \in l \right\} \\ \widetilde{\Sigma}_{\mathbb{S}_2}(x, l, \{l_i\}) &= \widetilde{\Sigma}_{\mathbb{S}_x'}(x, \{l_i\}) \cap \left\{ x \in l, f|_x^{(p)} = 0 = f|_x^{(p+1)}(\underbrace{v \dots v}_{p+1}) \forall v \in l \right\} \end{aligned}$$

The needed class $\left[\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, \{l_i\}) \right]$ is determined from this equation uniquely.

This theorem expresses the class $\left[\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, \{l_i\}) \right]$ in terms of the degenerated stratum $\left[\widetilde{\Sigma}_{\mathbb{S}_x', A_1}(x, y, l, \{l_i\}) \right]$ and the known classes $\left[\widetilde{\Sigma}_{\mathbb{S}_1}(x, l, \{l_i\}) \right]$ and $\left[\widetilde{\Sigma}_{\mathbb{S}_2}(x, l, \{l_i\}) \right]$, the later two correspond to curves with one singular point (of a linear type). Hence, after a few degenerating steps (not more than the order of determinacy of \mathbb{S}_x minus the multiplicity of \mathbb{S}_x), one arrives at the stratum of ordinary multiple points, the class of which was computed in the preceding section.

In particular, if $\text{o.d.}(\mathbb{S}_x) - \text{mult}(\mathbb{S}_x) = 1$, i.e. \mathbb{S}_x has a representative of the form $f_p + f_{p+1}$, with f_{p+1} generic, then the problem is solved in one step. If \mathbb{S}_x is already an ordinary multiple point then the theorem can be used to compute the class $\left[\widetilde{\Sigma}_{\mathbb{S}_x', A_1}(x, y, l) \right]$ in terms of $\left[\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l) \right]$, i.e. it gives an alternative (and fast) way to solve the enumerative problem for ordinary multiple points.

Various numerical answers are given in Appendix.

Proof. We need only to understand the residual pieces over the diagonal $\{x = y\}$ and their multiplicities. Consider the locally defining equations of $\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, \{l_i\})$ off $x = y$. As in §2.3.2 take the limit *ytox*, i.e. expand $y = x + \epsilon v$. Then the preliminary set of equations consists of the equations of \mathbb{S}_x and local expansion of $\mathbb{S}_y = A_1$:

$$(63) \quad \begin{aligned} f^{(p)}|_x &\sim \text{SYM}(l_1^{p_1}, \dots, l_k^{p_k}), \dots \text{ further equations of } \mathbb{S}_x, \dots \\ f^{(p)}|_x(\underbrace{v \dots v}_{p-1}) + \epsilon f^{(p+1)}|_x(\underbrace{v \dots v}_p) + \dots &= 0 = f^{(p+1)}|_x(\underbrace{v \dots v}_{p+1}) + \epsilon f^{(p+2)}|_x(\underbrace{v \dots v}_{p+2}) + \dots \end{aligned}$$

Here the proportionality of tensors is the only condition of \mathbb{S}_x involving $f^{(p)}|_x$.

This equation gives rise to the syzygies. We should take the flat limit as $\epsilon \rightarrow 0$. The following cases are possible:

- $\exists i : p_i \geq 2$ and $l_i = l$. Then in the equation above the term $f^{(p)}|_x(\underbrace{v \dots v}_{p-1})$ necessarily vanishes (as $v \in l_i$). So, the

equations place no conditions on $f^{(p)}|_x$ except for $f^{(p)}|_x \sim \text{SYM}(l_1^{p_1}, \dots, l_k^{p_k})$. Thus, generically $f^{(p)}|_x \neq 0$ and this locus of Aux does not contribute to the cohomology class (cf. §3.2.3).

- $\exists i : p_i = 1$ and $l_i = l$, but $\{l_i\}$ are pairwise distinct. Then in the equations above the number of independent (linear) conditions on $f^{(p)}|_x$ equals precisely the number of independent entries. So, the equations of this piece are $f^{(p)}|_x = 0 = f^{(p+1)}|_x(\underbrace{v \dots v}_{p+1})$ and some equations of \mathbb{S}_x on higher derivatives.

- $\forall i : l_i \neq l$ and $\{l_i\}$ are distinct. Then the number of independent (linear) conditions on $f^{(p)}|_x$ is bigger by one than the number of its independent entries. So, the equations of the contributing residual piece are: $f^{(p)}|_x = 0 = f^{(p+1)}|_x(\underbrace{v \dots v}_p)$.

Note that in both cases the residual pieces are reduced, as all the equations are linear in f or its derivatives.

Now compute the tangency degrees of the divisor $\text{kill}T_{\mathbb{S}_x}$ and the residual pieces. Namely, add to the locally defining ideal of $\widetilde{\Sigma}_{\mathbb{S}_x, A_1}(x, y, l, \{l_i\})$ near the contributing residual piece the local equation of $\text{kill}T_{\mathbb{S}_x}$ and check the primary decomposition of the bigger ideal. In the two relevant cases we have:

- $\exists i : p_i = 1$ and $l_i = l$, but $\{l_i\}$ are pairwise distinct. Then the enlarged ideal is generated by

(64)

$$f^{(p)}|_x = 0, \epsilon f^{(p+1)}|_x(\underbrace{v \dots v}_p) + \epsilon^2 f^{(p+2)}|_x(\underbrace{v \dots v}_{p+1}) \dots, f^{(p+1)}|_x(\underbrace{v \dots v}_{p+1}) + \epsilon f^{(p+2)}|_x(\underbrace{v \dots v}_{p+2}) + \dots, \dots \text{ conditions on higher derivatives}$$

And its primary decomposition consists of the two parts: $\left\{ f^{(p)}|_x, \epsilon, f^{(p+1)}|_x(\underbrace{v \dots v}_{p+1}), \dots \right\}$ and $\left\{ f^{(p)}|_x, f^{(p+1)}|_x(\underbrace{v \dots v}_p) + \epsilon^2 f^{(p+2)}|_x(\underbrace{v \dots v}_{p+1}) \dots, \dots \right\}$. So the tangency in this case is 1.

- $\forall i : l_i \neq l$ and $\{l_i\}$ are distinct. Then the enlarged ideal is generated by

$$(65) \quad f^{(p)}|_x = 0, \epsilon^2 f^{(p+2)}|_x(\underbrace{v \dots v}_{p+1}) + \dots = 0 = f^{(p+1)}|_x(\underbrace{v \dots v}_p) + \dots, \dots \text{ conditions on higher derivatives} \dots$$

So the tangency in this case is 2. ■

APPENDIX A. SOME NUMERICAL RESULTS FOR ENUMERATION OF CURVES WITH TWO SINGULAR POINTS

We give below explicit expressions $\deg \overline{\Sigma}_{\mathbb{S}_x, \mathbb{S}_y}$ for various types \mathbb{S}_x and \mathbb{S}_y an ordinary multiple point. This is only a tiny amount of possible enumerative results, for more formulas cf. the attached Mathematica file in the Arxiv [Kerner07-3] or on author's homepage.

As is well known (cf. [Kazarian01, Kazarian03-1]) for a collection of singularities $\mathbb{S}_1 \dots \mathbb{S}_r$ the degree of the stratum $\overline{\Sigma}_{\mathbb{S}_1 \dots \mathbb{S}_r}$ is expressed as:

$$(66) \quad \deg(\overline{\Sigma}_{\mathbb{S}_1 \dots \mathbb{S}_r}) = \frac{1}{|Aut|} \sum_{J_1 \sqcup \dots \sqcup J_k} S_{\{\mathbb{S}_i\}_{i \in J_1}} \dots S_{\{\mathbb{S}_i\}_{i \in J_k}}$$

Here the sum is over all the possible decompositions $\{\mathbb{S}_1, \dots, \mathbb{S}_r\} = \sqcup J_i$ and $S_{\{\mathbb{S}_i\}_{i \in J_i}}$ is the specialization of Thom polynomial, $S_{\mathbb{S}_x} = \deg(\overline{\Sigma}_{\mathbb{S}_x})$. One divides by $|Aut|$ the cardinality of the automorphisms (depending on the coincidence of the types). In particular, for two singular points: $\deg(\overline{\Sigma}_{\mathbb{S}_x, \mathbb{S}_y}) = \deg(\overline{\Sigma}_{\mathbb{S}_x}) \deg(\overline{\Sigma}_{\mathbb{S}_y}) + S_{\mathbb{S}_x, \mathbb{S}_y}$.

The degrees $\deg(\overline{\Sigma}_{\mathbb{S}_i})$ are known e.g. from [Kerner06, §A.2], for particular cases cf. example 3.2. For completeness we present them here.

- For the ordinary multiple point: $\deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}}) = \binom{p+2}{2} (d-p)^2$.
- For the singularity type \mathbb{S}_x with the representative $f_p + x_1^{p+1} + x_2^{p+1}$, and the tangent cone $T_{\mathbb{S}_x} = l_1^{p_1} \dots l_k^{p_k}$, with $\sum p_i = p$:

$$(67) \quad \deg(\overline{\Sigma}_{\mathbb{S}_x}) = \frac{1}{|G|} \left(\prod p_i \right) \left(k! \binom{p+2}{2} - 1 - k \right) (d-p)^2 + (k-1)! \binom{k+1}{2} \binom{p+2}{1} - 2 - k \binom{p+2}{1} (d-p) \sum p_i + (k-2)! \binom{k}{2} \binom{k+2}{2} \sum_{1 \leq i < j \leq k} p_i p_j \right)$$

Here G is the symmetry group, permuting lines with coinciding p_j 's.

- For the singularity type \mathbb{S}_x with the representative $(x_1^{p-1} + x_2^p)(x_1 + x_2^2)$,

$$(68) \quad \deg(\overline{\Sigma}_{\mathbb{S}_x}) = \frac{p(p+4)(p-1)}{8} (2p^3 + 7p^2 - 5p - 2)(d-p)^2 + \left(\binom{p+2}{2} - 3 \right) p^2 (d-p)(d-2(p+1)),$$

Below are the degrees of strata for curves with two singular points. For two ordinary multiple points the final answers are also given in §4.1. Recall, for $\overline{\Sigma}_{\mathbb{S}_x, \mathbb{S}_y}$ we always assume: $\text{mult}(\mathbb{S}_x) \geq \text{mult}(\mathbb{S}_y)$. If $\mathbb{S}_x = \mathbb{S}_y$ the final answer should be divided by 2.

(69)

$$\begin{aligned} \deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}, A_1}) &= \deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}}) \deg(\overline{\Sigma}_{A_1}) - \frac{3}{4} \binom{p+2}{3} (d-p)^2 (3p+4)(p^2+3p+4) + 3 \binom{p+2}{3} (d-p)(5p+6) \\ &\quad - \frac{5}{8} (d-p)^2 (p+1)(3p-1)(p^2+3p+8)(p^2+3p+10) \\ \deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}, D_4}) &= \deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}}) \deg(\overline{\Sigma}_{D_4}) + \left(\begin{aligned} &- \frac{5}{8} (d-p)^2 (p+1)(3p-1)(p^2+3p+8)(p^2+3p+10) \\ &+ 2(d-p)(p^2+3p+8)(35p^2+20p-12) - 6(85p^2+45p-28) \end{aligned} \right) \\ \deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}, X_9}) &= \deg(\overline{\Sigma}_{x_1^{p+1} + x_2^{p+1}}) \deg(\overline{\Sigma}_{X_9}) + \left(\begin{aligned} &- \frac{5}{8} (d-p)^2 (3p+2)(3p-2)(p^2+3p+16)(p^2+3p+18) \\ &+ 2(d-p)(p^2+3p+16)(270p^2-20p-117) - 14(830p^2-105p-348) \end{aligned} \right) \end{aligned}$$

$$(70) \quad \deg(\overline{\Sigma}_{x_1^p + x_2^p, A_1}) = \deg(\overline{\Sigma}_{x_1^p + x_2^p}) \deg(\overline{\Sigma}_{A_1}) - \frac{3}{8} p^4 (3+p)(d-p)^2 (p^2+3p-2) - \frac{3}{2} (p-1) p^3 (d-p)(p^2+3p-2) + 3p^4$$

(71)

$$\deg(\overline{\Sigma}_{y(x_1^p+x_2^{p+1}),A_1}) = \deg(\overline{\Sigma}_{y(x_1^p+x_2^{p+1})}) \deg(\overline{\Sigma}_{A_1}) + \left(\frac{-p^2(d-p)^2(5+p)}{(p+1)(p+5)p^3} (2+5p+p^2)(2+11p+6p^2) - \frac{p^2(p-1)}{8} (6p^4+41p^3+55p^2+64p+92) \right)$$

(72)

$$\deg(\overline{\Sigma}_{(x_2+y^2)(x_1^p+x_2^{p+1}),A_1}) = \deg(\overline{\Sigma}_{(x_2+y^2)(x_1^p+x_2^{p+1})}) \deg(\overline{\Sigma}_{A_1}) + \left(\frac{-p(d-p)^2(1+p)(p+6)}{+p(d-p)(p^2+7p+4)(9p^4+79p^3+220p^2+216p+63)} (p^2+7p+4)(9p^2+34p+24) - \frac{p(9p^6+124p^5+587p^4+1316p^3+1480p^2+654p+60)}{p^2+7p+4} \right)$$

(73)

$$\deg(\overline{\Sigma}_{(x_1^{p-1}+x_2^p)(x_1-x_2^2),A_1}) = \deg(\overline{\Sigma}_{(x_1^{p-1}+x_2^p)(x_1-x_2^2)}) \deg(\overline{\Sigma}_{A_1}) + \left(\frac{-9(p+3)p(d-p)^2(4+p+2p^2)+3p(p^4+3p^3+3p^2+4p-4)}{-\frac{3}{2}p^2(3+p)(d-p)(p^3-3p^2-p-8)} \right)$$

(74)

$$p > 2, \deg(\overline{\Sigma}_{(x_1^{p-2}+x_2^{p-2})(x_1^2-x_2^4),A_1}) = \deg(\overline{\Sigma}_{(x_1^{p-2}+x_2^{p-2})(x_1^2-x_2^4)}) \deg(\overline{\Sigma}_{A_1}) + \left(\frac{-\frac{3(d-p)^2}{8}(p^2+3p+6)(p^2+3p+8)(3p^3+6p^2+12p+5)}{+\frac{3(d-p)}{2}(p^2+3p+6)(3p^4+23p^3+48p^2+81p+29)} - \frac{3(57+182p+118p^2+51p^3+10p^4)}{3(57+182p+118p^2+51p^3+10p^4)} \right)$$

For small values $p = 2, 3$ these formulas reproduce the results of [Kazarian03-hab].

REFERENCES

- [AGLV] V.I.Arnol'd, V.V.Goryunov, O.V.Lyashko, V.A.Vasil'ev, *Singularity theory. I*. Springer-Verlag, Berlin, 1998.
- [Aluffi98] P.Aluffi *Characteristic classes of discriminants and enumerative geometry*, Comm. Algebra 26 (1998), no. 10, 3165–3193.
- [Caporaso-Harris98-1] L. Caporaso, J. Harris: *Enumerating rational curves: the rational fibration method*. *Compositio Math.* **113** (1998), no. 2, 209–236.
- [Caporaso-Harris98-2] L. Caporaso, J. Harris: *Counting plane curves of any genus*. *Invent. Math.* **131** (1998), no. 2, 345–392.
- [Cayley1866] A. Cayley, *On the theory of involution*, Trans. Cambridge Phil. Soc., XI. Part I (1866), 21–38. = “Coll. Math. Papers of A. Cayley,” **V**, [348], 1892, pp. 295–312.
- [Diaz-Harris86] S.Diaz, J.Harris, *Geometry of Severi varieties. II. Independence of divisor classes and examples*. Algebraic geometry (Sundance, UT, 1986), 23–50, Lecture Notes in Math., 1311, Springer, Berlin, 1988.
- [Diaz-Harris88] S.Diaz, J.Harris, *Geometry of the Severi variety*. Trans. Amer. Math. Soc. 309 (1988), no. 1, 1–34
- [Dimca-book] A.Dimca *Singularities and topology of hypersurfaces*. Springer-Verlag, Universitext. Springer-Verlag, New York, 1992.
- [Enriques32] F.Enriques, *Intorno ad alcune serie invarianti ...* Atti. Acad. Naz. Lincei 16(1932)
- [Fomin-Mikhalkin09] S.Fomin, G.Mikhalkin *Labeled floor diagrams for plane curves*, arXiv:0906.3828
- [di Francesco-Itzykson95] P. di Francesco, C. Itzykson: *Quantum intersection rings*, in: *The Moduli Space of Curves* (R. Dijkgraaf, C. Faber, and G. van der Geer, eds.), Progress in Mathematics, vol. 129, Birkhäuser, Boston, 1995, pp. 81–148
- [Fulton-book] W.Fulton: “Intersection theory.” Second edition. A Series of Modern Surveys in Mathematics, Springer-Verlag, Berlin, 1998.
- [van Gastel 89] L. van Gastel, *Characteristic numbers of plane curves: an excess intersection theoretical approach*. Enumerative algebraic geometry (Copenhagen, 1989), 259–265, Contemp. Math., 123.
- [van Gastel 91] L. van Gastel, *Excess intersections and a correspondence principle*. *Invent. Math.* 103 (1991), no. 1, 197–222.
- [Greuel-Lossen96] G.M.Greuel, C.Lossen, *Equianalytic and equisingular families of curves on surfaces*. *Manuscripta Math.* 91 (1996), no. 3, 323–342.
- [GLS01] G.-M. Greuel, C. Lossen, E. Shustin: *The variety of plane curves with ordinary singularities is not irreducible*. *Internat. Math. Res. Notices* **11** (2001), 543–550.
- [GLS06] G.-M. Greuel, C. Lossen, E. Shustin: *Equisingular Families of Projective Curves*, arXiv: math.AG/0612310
- [GLS-book1] G.-M. Greuel, C. Lossen, E. Shustin: *Introduction to Singularities and Deformations*. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [GLS-book2] G.-M. Greuel, C. Lossen, E. Shustin: *Singular algebraic curves*, to appear in Series: Springer Monographs in Mathematics 2008.
- [Göttsche98] L. Göttsche, *A conjectural generating function for numbers of curves on surfaces*, *Comm. Math. Phys* **196** 1998 523–33.
- [Harris-Pandharipande95] J. Harris and R. Pandharipande, *Severi degrees in cogenus 3*, preprint, arXiv:alg-geom/9504003.
- [Eisenbud-book] D.Eisenbud, *Commutative algebra. With a view toward algebraic geometry*. Graduate Texts in Mathematics, 150. Springer-Verlag, New York, 1995
- [Kazarian01] M. Kazarian. *Classifying spaces of singularities and Thom polynomials*. In New developments in Singularity Theory (Cambridge 2000), NATO Sci. Ser. II Math. Phys. Chem, 21, Kluwer Acad. Publ., Dordrecht, 2001, pp. 117–134.
- [Kazarian03-1] M. Kazarian. *Thom polynomials for Lagrange, Legendre, and critical point function singularities*. *Proc. Lond. Math. Soc.* (3) **86** (2003), 707–734.
- [Kazarian03-2] M. Kazarian. *Multisingularities, cobordisms, and enumerative geometry*. *uss. Math. Surveys* **58(4)** (2003), 665–724.
- [Kazarian03-hab] M. Kazarian. *Characteristic Classes in Singularity theory*, Doctoral Dissertation (habilitation thesis), Steklov Math. Inst., 2003. The file with many numerical results is on: “<http://www.mi.ras.ru/~kazarian/>”.
- [Kerner06] D.Kerner *Enumeration of singular algebraic curves*, *Israel Journal of Math.* 155 (2006), pp1-56. See updated and corrected version: arXiv:math.AG/0407358
- [Kerner08] D.Kerner *Enumeration of uni-singular algebraic hypersurfaces*, *Proc. Lond. Math. Soc.* (3) 96 (2008), no. 3, 623–668, cf. also arXiv:math/0505375
- [Kerner07-1] D.Kerner *On the geometry of some strata of uni-singular curves*, arXiv:math/0702735, preprint: Max Planck Institut für Mathematik Bonn, Germany, MPIM2007-68.
- [Kerner07-2] D.Kerner *On the collisions of singular points of complex algebraic plane curves.*, *Singularities II*, 89–110, Contemp. Math., 475, Amer. Math. Soc., Providence, RI, 2008, cf. also arXiv:0708.1228.

- [Kerner07-3] D.Kerner *On the enumeration of complex plane curves with two singular points* arXiv:0708.1229, cf. the attached Mathematica file. Available also on <http://www.math.bgu.ac.il/~kernerdm/index.html>
- [Kleiman76] S.L. Kleiman, *The enumerative theory of singularities*. Real and complex singularities (*Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Oslo, 1976*), pp. 297–396. Sijthoff and Noordhoff, Alphen aan den Rijn, 1977.
- [Kleiman85] S.L. Kleiman, *Intersection theory and enumerative geometry: a decade in review*. Proc. Sympos. Pure Math., 46, Part 2, Algebraic geometry, Bowdoin, 1985, 321–370, Amer. Math. Soc., Providence, RI, 1987.
- [Kleiman-Piene98] S.Kleiman, R.Piene, *Enumerating singular curves on surfaces*. *Algebraic geometry: Hirzebruch 70 (Warsaw, 1998)*, 209–238, *Contemp. Math.*, 241, Amer. Math. Soc., Providence, RI, 1999.)
- [Kleiman-Piene01] S.Kleiman, R.Piene, *Node polynomials for families: results and examples* preprint, arXiv:math/0111299
- [Kleiman-Piene04] S.Kleiman, R.Piene, Node polynomials for families: methods and applications. *Math. Nachr.* 271, 69–90 (2004).
- [Kontsevich-Manin94] M.Kontsevich, Yu.Manin, Gromov-Witten Classes, Quantum Cohomology, and Enumerative Geometry, *Comm. Math. Phys.* 164 (1994), no. 3, 525–562
- [Liu2000] Ai-Ko Liu, Family blowup formula, admissible graphs and the enumeration of singular curves. I. *J. Differential Geom.* 56 (2000), no. 3, 381–579
- [Lascoux77] A.Lascoux, *Sistemi lineari di divisori sulle curve e sulle superficie*. *Ann. Mat. Pura Appl.* (4) 114 (1977), 141–153.
- [Oka79] M.Oka, *On the bifurcation of the multiplicity and topology of the Newton boundary*. *J. Math. Soc. Japan* 31 (1979), no. 3, 435–450
- [Ran89] Z. Ran. *Enumerative geometry of singular plane curves*. *Invent. Math.* 97 (1989), no. 3, 447–465.
- [Ran02] Z. Ran. *Enumerative geometry of divisorial families of rational curves*, Preprint arXiv:math.AG/0205090.
- [Roberts1875] S. Roberts, *On a simplified method of obtaining the order of algebraical conditions*, Proc. Lond. Math. Soc. 1875 101–13.
- [Steiner1848] J. Steiner, *Allgemeine Eigenschaften der algebraischen Curven*, Berlin. Ber. 1848, 310–15 = *Journal Reine Anfew. Math.* 47 1853 1–6 = “Ges. Werke,” herausg. von K. Weierstrass, Berlin 1882, 2, 495–500.
- [Stückrad-Vogel82] J.Stückrad, W.Vogel, *An algebraic approach to the intersection theory*. The curves seminar at Queens, Vol. II (Kingston, Ont., 1981/1982), Exp. No. A, 32 pp., Queen’s Papers in Pure and Appl. Math., 61, Queen’s Univ., Kingston, ON, 1982.
- [Vainsencher81] I. Vainsencher. *Counting divisors with prescribed singularities*. *Trans. Amer. Math. Soc.* 267 (1981), no. 2, 399–422.
- [Vainsencher95] I.Vainsencher, *Enumeration of n-fold tangent hyperplanes to a surface*. *J. Algebraic Geom.* 4 (1995), no. 3, 503–526.
- [Vainsencher03] I. Vainsencher, *Hypersurfaces with up to six double points*. *Comm. Algebra* 31 (2003), no. 8, 4107–4129.

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, P.O.B. 653, BE’ER SHEVA 84105, ISRAEL.
E-mail address: kernerdm@math.bgu.ac.il