

## BIPARTITE $S_2$ GRAPHS ARE COHEN-MACAULAY

HASSAN HAGHIGHI, SIAMAK YASSEMI, AND RAHIM ZAARE-NAHANDI

ABSTRACT. In this paper we show that if the Stanley-Reisner ring of the simplicial complex of independent sets of a bipartite graph  $G$  satisfies Serre's condition  $S_2$ , then  $G$  is Cohen-Macaulay. As a consequence, the characterization of Cohen-Macaulay bipartite graphs due to Herzog and Hibi carries over this family of bipartite graphs. We check that the equivalence of Cohen-Macaulay property and the condition  $S_2$  is also true for chordal graphs and we classify cyclic graphs with respect to the condition  $S_2$ .

### INTRODUCTION

Let  $k$  be a field. To any finite simple graph  $G$  with vertex set  $V = [n] = \{1, \dots, n\}$  and edge set  $E(G)$  one associates an ideal  $I(G) \subset k[x_1, \dots, x_n]$  generated by all monomials  $x_i x_j$  such that  $\{i, j\} \in E(G)$ . The ideal  $I(G)$  and the quotient ring  $k[x_1, \dots, x_n]/I(G)$  are called the edge ideal of  $G$  and the edge ring of  $G$ , respectively. The simplicial complex of  $G$  is defined by

$$\Delta_G = \{A \subseteq V \mid A \text{ is an independent set in } G\},$$

where  $A$  is an independent set in  $G$  if none of its elements are adjacent. Note that  $\Delta_G$  is precisely the simplicial complex with the Stanley-Reisner ideal  $I(G)$ .

A graph  $G$  is said to be Cohen-Macaulay (resp. Buchsbaum) over  $k$ , if the edge ring of  $G$   $k[x_1, \dots, x_n]/I(G)$  is Cohen-Macaulay (resp. Buchsbaum), and is called Cohen-Macaulay (resp. Buchsbaum) if it is Cohen-Macaulay (resp. Buchsbaum) over any field. A graph is said to be chordal if each cycle of length  $> 3$  has a chord.

Let  $\Delta$  be a simplicial complex. This complex is called disconnected if the vertex set  $V$  of  $\Delta$  is the disjoint union of two nonempty sets  $V_1$  and  $V_2$  such that no face of  $\Delta$  has vertices in both  $V_1$  and  $V_2$ , otherwise it is called connected. A simplicial complex  $\Delta$  is called Cohen-Macaulay (resp. Buchsbaum) over an infinite field  $k$  if its Stanley-Reisner ring  $k[\Delta]$  is Cohen-Macaulay (resp. Buchsbaum).

It is known that if  $\Delta$  is a disconnected simplicial complex, then  $\text{depth } k[\Delta] = 1$ , [1, Chapter 5, Ex. 5.1.26]. This implies that if  $\text{depth } k[\Delta] > 1$ , then  $\Delta$  is connected. In particular, every Cohen-Macaulay simplicial complex of positive dimension is connected.

A satisfactory classification of all Cohen-Macaulay graphs over a field  $k$  has been standing open for some time. However, as pointed out by Herzog et al [6, Introduction], this is equivalent to a classification of all Cohen-Macaulay simplicial complexes over  $k$  which is clearly a hard problem. Accordingly, it is natural to

---

2000 *Mathematics Subject Classification.* 05C75, 13H10.

*Key words and phrases.* Bipartite graph, Cohen-Macaulay graph, Serre's condition, chordal graph.

Siamak Yassemi was supported in part by a grant from IPM No. 870130211.

Emails: haghghi@kntu.ac.ir, yassemi@ipm.ir, rahimzn@ut.ac.ir.

study special families of Cohen-Macaulay graphs. Recall that a graph  $G$  on the vertex set  $[n]$  is bipartite if there exists a partition  $[n] = V \cup W$  with  $V \cap W = \emptyset$  such that each edge of  $G$  is of the form  $\{i, j\}$  with  $i \in V$  and  $j \in W$ . It is easy to see that a graph  $G$  is bipartite if and only if it has no cycle of odd length. For a Cohen-Macaulay bipartite graph  $G$ , Estrada and Villarreal [2] showed that  $G \setminus \{\nu\}$  is Cohen-Macaulay for some vertex  $\nu \in V(G)$ . In [10] it is shown that the cyclic graph  $C_n$  is Cohen-Macaulay if and only if  $n \in \{3, 5\}$ . Herzog and Hibi gave a graph-theoretic characterization of all bipartite Cohen-Macaulay graphs. Due to our direct application, we state their result.

**Theorem** [5, Theorem 3.4]. Let  $G$  be a bipartite graph with vertex partition  $V \cup W$ . Then the following conditions are equivalent:

- (a)  $G$  is a Cohen-Macaulay graph;
- (b)  $|V| = |W|$  and the vertices  $V = \{x_1, \dots, x_n\}$  and  $W = \{y_1, \dots, y_n\}$  can be labeled such that:
  - (i)  $\{x_i, y_i\}$  are edges for  $i = 1, \dots, n$ ;
  - (ii) if  $\{x_i, y_j\}$  is an edge, then  $i \leq j$ ;
  - (iii) if  $\{x_i, y_j\}$  and  $\{x_j, y_k\}$  are edges, then  $\{x_i, y_k\}$  is also an edge.

Note that this result is characteristic-free.

Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . A subset  $C \subseteq V(G)$  is a *minimal vertex cover* of  $G$  if: (1) every edge of  $G$  is incident with a vertex in  $C$ , and (2) there is no proper subset of  $C$  with the first property. Observe that a minimal vertex cover is the set of indeterminates which generate a minimal prime ideal in the prime decomposition of  $I(G)$ . Also note that  $C$  is a minimal vertex cover if and only if  $V(G) \setminus C$  is a maximal independent set, i.e., a facet of  $\Delta_G$ .

A graph  $G$  is called *unmixed* if all minimal vertex covers of  $G$  have the same number of elements, i.e.,  $\Delta_G$  is pure. It is well known that every Cohen-Macaulay graph  $G$  is unmixed. A graph is called *chordal* if every cycle of length  $> 3$  has a chord. Recall that a chord of a cycle is an edge which joins two vertices of the cycle but is not itself an edge of the cycle.

Recall that a finitely generated graded module  $M$  over a Noetherian graded  $k$ -algebra  $R$  is said to satisfy the Serre's condition  $S_n$  if

$$\text{depth } M_{\mathfrak{p}} \geq \min(n, \dim M_{\mathfrak{p}}),$$

for all  $\mathfrak{p} \in \text{Spec}(R)$ . Thus,  $M$  is Cohen-Macaulay if and only if it satisfies the Serre's condition  $S_n$  for all  $n$ . A graph is said to satisfy the Serre's condition  $S_n$ , or simply is an  $S_n$  graph, if its edge ring satisfies this condition. Using [7, Lemma 3.2.1] and Hochster's formula on local cohomology modules, a pure  $d$ -dimensional Stanley-Reisner ring  $k[\Delta]$  satisfies  $S_2$  property if and only if  $\tilde{H}_0(\text{link}_{\Delta}(F); k) = 0$  for all  $F \in \Delta$  with  $|F| \leq d - 2$  (see [8, page 4]).

The main result of this paper is to prove that if  $G$  is a bipartite  $S_2$  graph, then  $G$  is Cohen-Macaulay (see Theorem 1.3). Consequently, the characterization of Cohen-Macaulay bipartite graphs by Herzog and Hibi carries over bipartite  $S_2$  graphs. It is shown that not only for bipartite graphs but also for chordal graphs Cohen-Macaulay property and the condition  $S_2$  are equivalent. To see an example of a non-Cohen-Macaulay  $S_2$  graph, it is shown that the cyclic graph  $C_n$  of length  $n \geq 3$  is  $S_2$  if and only if  $n = 3, 5$  or  $7$ . In particular,  $C_7$  is the only cyclic graph which is  $S_2$  but not Cohen-Macaulay. Finally, we reprove some known results

on certain bipartite Cohen-Macaulay graphs by providing rather simpler proofs compared to the existing ones.

## 1. THE MAIN RESULT

Our results are inspired by the aforementioned theorem of Herzog and Hibi [5, Theorem 3.4].

**Proposition 1.1.** *Let  $G$  be an unmixed bipartite graph with bipartition  $V = \{x_1, \dots, x_n\}$  and  $W = \{y_1, \dots, y_n\}$  such that  $\{x_i, y_i\}$  is an edge of  $G$  for all  $i = 1, \dots, n$ . Then  $V$  and  $W$  can be simultaneously relabeled such that the following statements are equivalent:*

- (a) *There exists a linear order  $V = F_0, \dots, F_n = W$  on some of the facets of  $\Delta_G$  such that  $F_i$  and  $F_{i+1}$  intersect in codimension one for  $i = 0, \dots, n-1$ .*
- (b) *If  $\{x_i, y_j\}$  is an edge, then  $i \leq j$ .*

By a simultaneous relabeling we mean that for all  $i$ ,  $x_i$  and  $y_i$  receive the same relabeling. In particular, under the assumptions of Proposition 1.1, with the new labeling,  $\{x_i, y_i\}$  is an edge of  $G$  for all  $i = 1, \dots, n$ .

Before proceeding on the proof of this Proposition note that the condition (a) is weaker than strongly connectedness of  $\Delta_G$ . Recall that a simplicial complex  $\Delta$  is strongly connected if for any two facets  $V$  and  $W$  of  $\Delta$  there exists a chain of facets satisfying (a). Here we only need this sequence just for the two specific facets  $V$  and  $W$ .

*Proof.* (a) $\Rightarrow$ (b): We have  $|F_1 \setminus F_0| = 1$ , say  $F_1 \setminus F_0 = \{y_1\}$ . Then  $F_1 = \{y_1, x_2, \dots, x_n\}$  because  $\{x_1, y_1\}$  is not a face of  $\Delta_G$ . Similarly,  $|F_2 \setminus F_1| = 1$ , say  $F_2 \setminus F_1 = \{y_2\}$ . Thus  $F_2 = \{y_1, y_2, x_3, \dots, x_n\}$  because again  $\{x_2, y_2\}$  is not a face of  $\Delta_G$ . Hence by induction we may assume that  $F_i = \{y_1, \dots, y_i, x_{i+1}, \dots, x_n\}$  for  $i = 0, \dots, n$ . In particular, if  $i > j$ , then  $\{x_i, y_j\}$  is a face of  $\Delta_G$ , and hence it is not an edge of  $G$ .

(b) $\Rightarrow$ (a): Set  $F_i = \{y_1, \dots, y_i, x_{i+1}, \dots, x_n\}$ . It is easy to see that for any  $i$ ,  $F_i$  is a maximal independent set and hence a facet of  $\Delta_G$ . Moreover  $F_i$  and  $F_{i+1}$  intersect in codimension one. □

**Lemma 1.2.** *Let  $G$  be a bipartite graph. Then  $G$  is a non-complete bipartite graph if and only if  $\Delta_G$  is connected.*

*Proof.* Let  $V_1 \cup V_2$  be the bipartition of  $G$ . Then  $G$  fails to be a complete bipartite graph if and only if there are two vertices  $x \in V_1$  and  $y \in V_2$  which are not adjacent, that is,  $\{x, y\}$  is an independent set of  $G$ , i.e.,  $\Delta_G$  is connected. □

Now we may state the main result which in particular provides a characterization of bipartite  $S_2$  graphs.

**Theorem 1.3.** *Let  $G$  be a bipartite graph with at least four vertices and with vertex partition  $V$  and  $W$ . Then the following are equivalent:*

- (a)  *$G$  is unmixed and  $V$  and  $W$  can be labeled such that there exists an order  $V = F_0, \dots, F_n = W$  of the facets of  $\Delta_G$  where  $F_i$  and  $F_{i+1}$  intersect in codimension one for  $i = 0, \dots, n-1$ .*
- (b)  *$G$  is a Cohen-Macaulay graph.*

- (c)  $G$  is a Buchsbaum non-complete bipartite graph.
- (d)  $G$  is an  $S_2$  graph.

*Proof.* We prove  $(a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) \Rightarrow (a)$ .

(a) $\Rightarrow$ (b): Since  $G$  is unmixed, by König's Theorem there is a bipartition  $V = \{x_1, \dots, x_n\}$  and  $W = \{y_1, \dots, y_n\}$  such that  $\{x_i, y_i\}$  is an edge of  $G$  for all  $i$ . By Proposition 1.1,  $V$  and  $W$  can be relabeled such that  $\{x_i, y_i\}$  is an edge of  $G$  for all  $i$  and if  $\{x_i, y_j\}$  is an edge in  $G$ , then  $i \leq j$ . We fix such a labeling. Let  $\{x_i, y_j\}$  and  $\{x_j, y_k\}$  be edges of  $G$  with  $i < j < k$ , and suppose that  $\{x_i, y_k\}$  is not an edge of  $G$ . Since  $\{x_i, y_k\}$  is a face of  $\Delta_G$  and  $G$  is unmixed,  $\Delta_G$  is pure, hence there exists a facet  $F$  of  $\Delta_G$  with  $|F| = n$  and  $\{x_i, y_k\} \subset F$ . Since  $F$  is a facet of  $\Delta_G$ , any 2-element subset of  $F$  is a non-edge of  $G$ . We have  $y_j \notin F$  since  $\{x_i, y_j\}$  is an edge of  $G$ . Similarly  $x_j \notin F$  since  $\{x_j, y_k\}$  is an edge of  $G$ . On the other hand, since  $\{x_t, y_t\}$  is an edge of  $G$  for all  $t$ , the facet  $F$  can not contain both  $x_t$  and  $y_t$ . Hence  $F$  is of the form  $F = \{z_1, \dots, z_n\}$ , where  $z_t = x_t$  or  $y_t$  for  $t = 1, \dots, n$ . Thus either  $y_j$  or  $x_j$  belongs to  $F$ , which is a contradiction. Consequently,  $G$  is Cohen-Macaulay by the theorem of Herzog and Hibi.

(b) $\Rightarrow$ (c): Since every Cohen-Macaulay ring is a Buchsbaum ring,  $G$  is also Buchsbaum. By definition, the ideal of the simplicial complex  $\Delta_G$  is equal to edge ideal of  $G$ . Hence  $\Delta_G$  is also Cohen-Macaulay and in particular,  $\Delta_G$  is connected. Therefore, by Lemma 1.2  $G$  is non-complete.

(c) $\Rightarrow$ (d): By [11, Corollary 2.7] the localization of every Buchsbaum ring at any of its prime ideals which is not equal to  $(x_1, \dots, x_n, y_1, \dots, y_n)$ , is Cohen-Macaulay. Therefore  $G$  satisfies the  $S_2$  condition.

(d) $\Rightarrow$ (a): Since  $\Delta_G$  satisfies the  $S_2$  condition, by [4, Corollary 2.4] for any two facets  $F$  and  $H$  of  $\Delta_G$ , there exist a positive integer  $m$  and a sequence  $F = F_0, \dots, F_m = H$  of facets of  $\Delta_G$  such that  $F_i$  intersects  $F_{i+1}$  in codimension one for all  $i = 0, \dots, m-1$ . Hence  $\Delta_G$  is strongly connected. In particular, since the partitions  $V$  and  $W$  of the vertices of  $G$  can be considered as two facets of  $\Delta_G$  and  $\Delta_G$  is strongly connected, the required sequence exists. Furthermore,  $|F_i| = |F_i \cap F_{i+1}| + 1 = |F_{i+1}|$  for all  $i = 0, \dots, m-1$ . This implies that any two facets of  $\Delta_G$  have the same number of elements and hence  $G$  is unmixed.  $\square$

**Remark 1.4.** The implication (b) $\Rightarrow$ (a) in the above theorem does not depend on the bipartite assumption of  $G$  and is valid in a more general setting. In fact a stronger implication is valid. More precisely, every Cohen-Macaulay simplicial complex is strongly connected. This follows, for example, by an argument similar to the implication (d) $\Rightarrow$ (a).

**Remark 1.5.** Theorem 1.3 reveals that for bipartite graphs Cohen-Macaulay and  $S_2$  properties are equivalent. This raises the question whether there are other families of graphs for which these two properties are equivalent. Here, we show that,

- (1) Every chordal  $S_2$  graph is Cohen-Macaulay.
- (2) The cyclic graph  $C_7$  is  $S_2$  but not Cohen-Macaulay.

In fact, chordal graphs are shellable [9, Theorem 2.13]. But any  $S_2$  graph is unmixed (see [3, Corollary 5.10.9], or [4, Remark 2.4.1]). Therefore, for chordal graphs Cohen-Macaulay and  $S_2$  properties are equivalent.

To establish (2) we classify all cyclic graphs  $C_n$  with respect to  $S_2$  property.

**Proposition 1.6.** *The cyclic graph  $C_n$  of length  $n \geq 3$  is  $S_2$  if and only if  $n = 3, 5$  or  $7$ . In particular,  $C_7$  is the only cyclic graph which is  $S_2$  but not Cohen-Macaulay.*

*Proof.* It is known that  $C_n$  is Cohen-Macaulay if and only if  $n = 3, 5$  [10, Corollary 6.3.6]. On the other hand,  $C_n$  of length  $n \geq 3$  is unmixed if and only if  $n = 3, 4, 5, 7$  [10, Exercise 6.2.15]. Accordingly,  $C_3$  and  $C_5$  are  $S_2$ . Since  $C_4$  is bipartite but not Cohen-Macaulay, by Theorem 1.3 it is not  $S_2$ . Furthermore, as mentioned before, every  $S_2$  graph is unmixed. Thus, the only cyclic graph which remains to be checked is  $G = C_7$ . To settle this, we apply the cohomological criterion for  $S_2$  property mentioned in the introduction. In fact, we need to check that for all  $F \in \Delta_G$  with  $|F| \leq 1$ ,  $\tilde{H}_0(\text{link}_{\Delta_G}(F); k) = 0$ . This condition is satisfied if  $\text{link}_{\Delta_G}(F)$  is connected which can easily be checked by direct inspection.  $\square$

In light of Theorem 1.3, we consider some known results on certain bipartite Cohen-Macaulay graphs and we provide rather simpler proofs compared to the existing ones.

As a consequence of Theorem 1.3(b) we may state the following result on the structure of trees satisfying the condition  $S_2$ .

**Corollary 1.7.** [10, Theorem 6.3.4] *Let  $G$  be a tree with at least four vertices. Then the following are equivalent:*

- (a)  *$G$  satisfies the condition  $S_2$ .*
- (b) *There is a bipartition  $V = \{x_1, \dots, x_n\}$ ,  $W = \{y_1, \dots, y_n\}$  of  $G$  such that*
  - (i)  *$\{x_i, y_i\} \in E(G)$  for all  $i$ .*
  - (ii) *for each  $i$  either  $\deg(x_i) = 1$  or  $\deg(y_i) = 1$ , exclusively.*
  - (iii)  *$V$  and  $W$  can be simultaneously relabeled such that there exists an order  $V = F_0, \dots, F_n = W$  of the facets of  $\Delta_G$  where  $F_i$  and  $F_{i+1}$  intersect in codimension one for  $i = 0, \dots, n-1$ .*

From part (b)(ii) of Corollary 1.7 it follows that every tree with  $2n$  vertices which satisfies the condition  $S_2$ , has precisely  $n$  vertices of degree one.

**Corollary 1.8.** *Every path of length greater than four does not satisfy the condition  $S_2$  and hence it is not Cohen-Macaulay.*

By Corollary 1.7 every bipartite  $S_2$  graph has at least two vertices of degree one. From this fact and Theorem 1.3 we get the following result which is a special case of [10, Proposition 6.2.1].

**Proposition 1.9.** *Let  $G$  be a bipartite  $S_2$  graph. Let  $y$  be a vertex of degree one of  $G$  and  $x$  its adjacent vertex. Then  $G \setminus \{x, y\}$  is still an  $S_2$  graph.*

*Proof.* Since  $G$  is bipartite, there exists an order  $V = F_0, \dots, F_n = W$  of facets of  $\Delta_G$  such that for each  $i = 0, \dots, n-1$ ,  $F_i$  intersects  $F_{i+1}$  in codimension one. Since for each  $i$ ,  $V \cup W \setminus F_i$  is a minimal vertex cover of  $G$ , it contains exactly one of the vertices  $x$  or  $y$ . Thus  $F_i$  contains  $y$  or  $x$  respectively. Again since any facet of  $\Delta_G$  is an independent set, none of these facets can contain both of these elements. Thus, if we delete both of these elements from  $V(G)$ , then they will be deleted from each element of the sequence  $V = F_0, \dots, F_n = W$ . By construction  $F_0 \setminus \{x\} = F_1 \setminus \{y\}$ , and hence we obtain a sequence of length  $n-1$  of facets of  $\Delta_{G \setminus \{x, y\}}$  such that each two consecutive members of this sequence intersect each other in codimension one. Now the claim follows from Theorem 1.3(b).  $\square$

**Remark 1.10.** A careful inspection of the proof of Proposition 1.9 reveals that every edge  $\{x, y\}$  where  $y$  is an arbitrary degree one vertex of  $G$ , intersects every member of the sequence  $F_0, \dots, F_n$ . Conversely, if we add a new vertex  $x_{n+1}$  to  $V$  and a new vertex  $y_{n+1}$  to  $W$  and the edge  $\{x_{n+1}, y_{n+1}\}$  to  $G$ , then the bipartite graph  $G_1 = V_1 \cup W_1$ , where  $V_1 = V \cup \{x_{n+1}\}$  and  $W_1 = W \cup \{y_{n+1}\}$ , has the sequence  $F_0 \cup \{x_{n+1}\}, F_1 \cup \{x_{n+1}\}, \dots, F_n \cup \{x_{n+1}\}, F_{n+1} = F_n \cup \{y_{n+1}\}$  as a subsequence of its facets which satisfies the assumption of Theorem 1.3(b), hence  $G_1$  is an  $S_2$  graph.

We end this paper with the following immediate result which is again a special case of [10, Proposition 6.2.1].

**Corollary 1.11.** *Let  $G$  be a tree with more than two vertices which is  $S_2$ . Let  $x$  be a degree one vertex of  $G$  and  $y$  its adjacent vertex. Then  $G \setminus \{x, y\}$  is an  $S_2$  graph.*

#### ACKNOWLEDGMENTS

This paper was initiated during Yassemi's visit of the Max-Planck Institute fur Mathematics (MPIM). He would like to thank the authorities of MPIM for their hospitality during his stay there. The authors are grateful to the referee for a careful review of this paper and for several corrections and remarks.

#### REFERENCES

1. W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, Cambridge University Press, 1993.
2. M. Estrada and R. H. Villarreal, *Cohen-Macaulay bipartite graphs*, Arch. Math. **68** (1997), 124-128.
3. A. Grothendieck, *Éléments de géométrie algébrique IV. Étude locale des schémas et des morphismes deschémas. II*. Inst. Hautes Études Sci. Publ. Math., No. 24, 1965.
4. R. Hartshorne, *Complete intersection and connectedness*, Amer. J. Math. **84** (1962), 497-508.
5. J. Herzog and T. Hibi, *Distributive lattices, bipartite graphs and Alexander duality*, J. Algebraic Combin. **22** (2005), no. 3, 289-302.
6. J. Herzog, T. Hibi and X. Zheng, *Cohen-Macaulay chordal graphs*, J. Combin. Theory Series A **113** (2006), 911-916.
7. P. Schenzel, *Dualisierende Komplexe in der lokalen Algebra und Buchsbaum Ringe*, LNM **907**, Springer, 1982.
8. N. Terai, *Alexander duality in Stanley-Reisner rings*, in "Affine Algebraic Geometry (T. Hibi, ed.)", Osaka University Press, Osaka 2007, 449-462.
9. A. Van Tuyl and R. H. Villarreal, *Shellable graphs and sequentially bipartite graphs*, J. Combin. Theory Ser. A **115** (2008), no. 5, 799-814, arXiv:math/0701296.
10. R. H. Villarreal, *Monomial Algebra*, Dekker, New York, NY, 2001.
11. K. Yanagawa, *Alexander duality for Stanley-Reisner rings and squarefree  $\mathbb{N}^n$ -graded modules*, J. Alg. **225** (2000), 630-645.

HASSAN HAGHIGHI, DEPARTMENT OF MATHEMATICS, K. N. TOOSI UNIVERSITY OF TECHNOLOGY, TEHRAN, IRAN.

SIAMAK YASSEMI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEHRAN, TEHRAN, IRAN AND SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), TEHRAN IRAN.

RAHIM ZAARE-NAHANDI, SCHOOL OF MATHEMATICS, STATISTICS & COMPUTER SCIENCE, UNIVERSITY OF TEHRAN, TEHRAN, IRAN.