



eCOMMONS

Loyola University Chicago
Loyola eCommons

Bioinformatics Faculty Publications

Faculty Publications

2-8-2018

Draft Genome Sequence of the Siderophore-Degrading Soil Bacterium *Mesorhizobium loti* Strain LU

Domenic Castignetti
Loyola University Chicago

Nathaniel Polley
Loyola University Chicago

Catherine Putonti
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub

 Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation


Castignetti, Domenic; Polley, Nathaniel; and Putonti, Catherine. Draft Genome Sequence of the Siderophore-Degrading Soil Bacterium *Mesorhizobium loti* Strain LU. *Prokaryotes: Genome Announcements*, , : 2, 2018. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/genomeA.00029-18>

This Article is brought to you for free and open access by the Faculty Publications at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

© The Authors 2018



Draft Genome Sequence of the Siderophore-Degrading Soil Bacterium *Mesorhizobium loti* Strain LU

Domenic Castignetti,^a Nathaniel Polley,^{a,b}  Catherine Putonti^{a,b,c}

^aDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

ABSTRACT Here, we present the draft genome of *Mesorhizobium loti* strain LU, a soil bacterium capable of degrading the trihydroxamate siderophore deferrioxamine B to its constituent monohydroxamic acids. Genome size was 6,399,828 bp, with a GC content of 61.5%. This draft genome consists of 35 scaffolds, with an N_{50} of 389,921 bp.

The bacterial symbionts, known as the rhizobia, fix atmospheric nitrogen in association with leguminous plants (1). Rhizobia have a significant role in the nitrogen fertility of soils and in agriculture, and the legumes are considered among the most important of agricultural species, second only to the grasses. As crops, the legumes supply the largest source of plant protein in the diets of both humans and livestock (2). In more recent times, the genus *Rhizobium* has been refined and divided into genera such as *Bradyrhizobium*, *Sinorhizobium*, and *Mesorhizobium* (1).

Siderophore degradation has been noted only in a select group of microbes (3). *Mesorhizobium loti* strain LU, isolated from a sample of garden soil in Glenview, IL, USA, is a siderophore-degrading bacterium that synthesizes the constituent monohydroxamic acids from the trihydroxamate siderophore deferrioxamine B and is able to use the siderophore as its sole source of carbon for growth (3). Experimental evidence indicates that the enzyme responsible for deferrioxamine B degradation is most likely a serine peptidase-like enzyme (4).

M. loti LU genomic DNA was isolated following growth in enriched medium, such as nutrient broth or tryptic soy broth, using the Wizard genomic DNA purification kit (Promega, Madison, WI) as described by the manufacturer. Genomic DNA concentration was determined using the Qubit fluorimeter. Library preparation for Illumina sequencing was performed by the Loyola University Chicago Genomics Facility; the Nextera XT DNA library preparation kit was used. The library was sequenced on the MiSeq Sequencer (Illumina) using the MiSeq reagent kit v2 (500 cycles). The run produced 1,603,054 paired-end reads in total.

Reads were trimmed using the tool sickle (<https://github.com/najoshi/sickle>) and assembled using SPAdes (v3.11.1) (5), producing 126 contigs. Coverage was assessed using BBMap (<http://sourceforge.net/projects/bbmap/>). Contigs with a coverage of less than 1 were removed from further consideration. This resulted in a final set of 35 scaffolds varying in size from 1,001 bp to 707,823 bp (N_{50} , 389,921 bp) with a coverage of 67×. The genome size was 6,399,828 bp with a GC content of 61.53%. Annotations were produced by the NCBI Prokaryotic Genome Annotation Pipeline using GeneMarkS+ (6). Five rRNA genes, 44 tRNA genes, and 5,989 protein-coding sequences were detected. The annotation pipeline identified 79 pseudogenes within the assembly.

Received 10 January 2018 **Accepted** 12 January 2018 **Published** 8 February 2018

Citation Castignetti D, Polley N, Putonti C. 2018. Draft genome sequence of the siderophore-degrading soil bacterium *Mesorhizobium loti* strain LU. Genome Announc 6:e00029-18. <https://doi.org/10.1128/genomeA.00029-18>.

Copyright © 2018 Castignetti et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Domenic Castignetti, dcastig@luc.edu.

Accession number(s). The draft whole-genome project for *Mesorhizobium loti* strain LU has been deposited at DDBJ/EMBL/GenBank under accession number [PJRO00000000](https://doi.org/10.1016/j.micres.2013.09.012). Raw sequence reads are deposited at DDBJ/EMBL/GenBank under accession number SRR6382439.

ACKNOWLEDGMENT

This work was supported by Loyola University Chicago.

REFERENCES

1. Laranjo M, Alexandre A, Oliveira S. 2014. Legume growth-promoting rhizobia: an overview on the *Mesorhizobium* genus. *Microbiol Res* 169: 2–17. <https://doi.org/10.1016/j.micres.2013.09.012>.
2. Garg N, Geetanjali N. 2007. Symbiotic nitrogen fixation in legume nodules: process and signaling. A review. *Agron Sustain Dev* 27:59–68. <https://doi.org/10.1051/agro:2006030>.
3. Pierwola A, Krupinski T, Zalupski P, Chiarelli M, Castignetti D. 2004. Degradation pathway and generation of monohydroxamic acids from the trihydroxamate siderophore deferrioxamine B. *Appl Environ Microbiol* 70:831–836. <https://doi.org/10.1128/AEM.70.2.831-836.2004>.
4. Zaya N, Roginsky A, Williams J, Castignetti D. 1998. Evidence that a deferrioxamine B degrading enzyme is a serine protease. *Can J Microbiol* 44:521–527. <https://doi.org/10.1139/w98-031>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.