Computational investigations on the unexpected extrusion of molecular iodine in Pd(II) σ -butadienyl complexes.

<u>Thomas Scattolin</u>^a, Fabiano Visentin^a, Luciano Canovese^a and Claudio Santo^a *Dipartimento di Scienze Molecolari e Nanosistemi, Universita` Ca' Foscari, Venezia*e-mail: thomas.scattolin@unive.it

We have experimentally and theoretically studied the stoichiometric addition of halogens or interhalogens to σ -butadienyl Pd(II) complexes bearing thioquinolines as spectator ligands. The observed reactions do not involve the expected elimination of the butadienyl fragment but rather the unpredictable extrusion of molecular iodine (Schemes 1-2).

We have explained this peculiar reactivity with a mechanistic hypothesis (Scheme 3) involving Pd(IV) intermediates (to save computer time, the COOMe was substituted with CN group). In the case of the reaction between complex Ia and IBr, it is apparent from the computational output (Scheme 4) that I_2 and complex Ib represent the favored reaction products from both kinetic and thermodynamic points of view (the energy values are expressed as ΔG° at 298K).

The geometrical optimization of the complexes was carried out using the hyper-GGA functional MO6 3 in combination with the LAN2TZ(f) 4 basis set for the Pd atoms, the LANL2DZdp basis set for the halogen atoms and the 6-31G(d,p) basis set for the other elements. Solvent effects (dichloromethane, ε =8.93) were included using CPCM 6 . The thermodynamic parameters were obtained by means of the stationary points characterized by IR simulation.

References

- 1. T.Scattolin, F.Visentin, C.Santo, V.Bertolasi, L.Canovese, Dalton Trans., 2016, 45, 11560-11567
- 2. R.V.Belzen, C.J.Elsevier, A.Dedieu, N.Veldman, A.L.Spek, Organometallics, 2003, 22, 722-736
- 3. Y.Zhao, D.G.Truhlar, Theor. Chem. Acc., 2008, 120, 215-241
- 4. L.E.Roy, P.J.Hay, R.L.Martin, J. Chem. Theory Comput., 2008, 4, 1029-1031
- 5. C.E.Check, T.Faust, J.M.Bailey, B.J.Wright, T.Gilbert, J. Phys. Chem. A, 2001, 105, 8111-8116
- 6. V.Barone, M.Cossi, J.Tomasi, J. Chem. Phys. 1997, 107, 3210-3221